1175

PILOT RANDOMIZED STUDY ON THE RISK OF INTRAVENTRICULAR HEMORRHAGE WITH MIDLINE VERSUS LATERAL HEAD POSITIONS

S. Al-Abdi¹, D.M. Nojoom², H. Alshaalan²,
M. Al-Aamri¹

¹*Pediatrics, King Abdulaziz Hospital (KAH), Al-Ahsa, ²Medical Imaging Department, King Abdulaziz Medical City, Riyadh, Saudi Arabia*

Objective: To study a hypothesis that keeping preterm infants' heads in midline positions reduces their risk of intraventricular hemorrhage (IVH).

Methods: We conducted a pilot randomized trial to study this hypothesis by comparing the IVH risk associated with flat midline (FM) and flat lateral (FL) head positions throughout the first seven days of life. Randomization was stratified based on gender and gestational age (< 27 or 27-29.9 weeks).

Results: Twenty-three infants younger than 30 weeks gestation were randomly assigned to the FM group, and 21 infants of similar age were assigned to the FL group (right-tilted: n = 9; left-tilted: n = 12). Important baseline characteristics were similar in both groups, except for multiple births which were significantly higher in the FM group. The risk of IVH was higher in the FM group than in the FL group (6 vs. 4; risk ratio (RR) 1.4; 95% confidence interval (CI): 0.4-4.2; p = 0.8). A secondary analysis showed that the risk of IVH was two times higher in the FM group than in the right-tilted FL group (6 vs. 1; RR 2.3; 95% CI: 0.3-16.9; p = 0.6) and was the same in the FM and left-tilted FL groups (6 vs. 3; RR 1.0; 95% CI: 0.3-3.5; p > 0.99).

Conclusions: Keeping a preterm infant's head in the FM position throughout the first seven days of life does not decrease the risk of IVH; however, a right-tilted FL position may reduce this risk. A larger study is needed for a definitive conclusion.

1176

RADIOGRAPHIC EXPOSURE IN THE FIRST WEEKS OF LIFE IN PRETERM INFANTS

S. Sanka, A. Curley

Neonatal Intensive Care Unit, Addenbrookes Hospital NHS Trust, Cambridge, UK

Aim: To determine the number of x rays performed on preterm infants (< 30 weeks gestation) admitted to a tertiary centre during the first seven weeks of life.

Methods: We calculated the number of chest, abdominal (AP and lateral), combined chest/abdominal x rays and x rays taken for contrast studies during the first seven weeks of life in preterm infants under 30 weeks gestation admitted to a single tertiary centre during 2008.

Results:

Jan-Dec 2008	CXR Median (range)	AXR Median (range)	CAXR Median (range)	Contrast study images. Pt= patient	No. of babies (No. of weeks stay)
24W	8 (1-22)	1(0-3)	3(0-10)	none	11(50)
25W	3.5(0-13)	0.5(0-9)	4(0-11)	24 (1 pt)	14(70)
26W	3(1-18)	0(0-9)	3.5(1-7)	0	12(46)
27W	2(1-14)	1(0-5)	2(0-13)	0	25(88)
28W	2(0-8)	0(0-6)	2(0-11)	8(1pt)	23(78)
29W	2(0-12)	0(0-7)	2(0-10)	9(1 pt)	21(71)
30W	1(0-16)	0(0-1)	0(0-4)	0	15(58)

[Results]

Conclusions: Preterm infants are exposed to a number of radiographs in the early weeks of life due to associated morbidities and also have diagnostic procedures that may involve X ray radiation. Repeated handling for radiographs can cause clinical instability and accumulation of a radiation dose that can have serious health consequences. Careful consideration should be given before requesting a radiograph in this vulnerable group.

1177

THE EFFECTS OF A DESIGNED PROGRAM ON OXYGEN SATURATION AND HEART RATE OF PREMATURE INFANTS HOSPITALIZED IN NICU IN -2009

P. Taheri, E. Abbasi, Z. Abdeyazdan,
N. Fathizadeh

Paediatric Nursing Dept, University Hospital Paediatric, Isfahan, Iran

Background: Prematurity is the main cause of death in infants under one years of age and is the main reason for NICU hospitalization. The stressful environment of NICU exposes preterm infants to

inappropriate stimuli. This study aimed to determine and compare the mean of rate and oxygen saturation of premature infants before and during a designed program in NICU.

Methods: In a clinical trial study of before and after intervention on a single group, 31 hospitalized premature newborn in NICU of Hospital in Isfahan were selected by simple continuous sampling method. Data were collected through interview, observation and records of a checklist. The data was analyzed using SPSS and descriptive and inferential statistics.

Results: Out of 31 premature infants in the study, 60% were boys and 35% were girls. The mean of arterial oxygen before and during the designed program were 92.80 ± 2.54 and 94.22 ± 2.59 , respectively. The results of paired t test showed a significant difference between the mean of arterial oxygen saturation of the infants before and during the program ($p=0.048$), but there was no significant difference between the mean of the infants' heart beat before and during the intervention.

Conclusions: The findings showed that applying daily silence periods can greatly help to increase oxygen saturation and can improve the growth of premature infants. Therefore, by programs to reduce light and noise in these wards would be possible.

Keywords: Programmed instruction, prematurity, neonatal, pulse oximetry, noise, light.

1178

IMMEDIATE MANAGEMENT OF GROWTH RESTRICTED INFANT

E. Petkovska, S. Jancevska

Department of Neonatal Intensive Care and Therapy, University Clinic of Gynecology and Obstetrics, Skopje, FYR Macedonia

Background and aims: Intrauterine growth restriction (IUGR) is characterized by fetal growth less than normal for the population and growth potential of a given infant. The decreased fetal growth rate in IUGR is an adaptation to an unfavourable intrauterine environment and may result in permanent alterations in metabolism, growth, and development. The acute neonatal consequences of IUGR are perinatal asphyxia and neonatal adaptive problems.

Methods: During the period of 3 years, immediate management of 160 infants from fetal growth restricted pregnancies were studied. We used SPSS 1.1 and Statgraf for Win 2.1 statistical programs. Results were compared by Person Chi-Square (< 0.05) and logistic regression analyses.

Results: Compared with normally grown infants, those who were growth restricted had increased risk for fetal distress (OR 16.47; 95%CI 6.86-39.55), operative way of delivery (OR 4.25; 95%CI 2.72-6.64), perinatal asphyxia (OR 3.26; 95%CI 1.96-5.43), need for resuscitation (OR 2.81; 95%CI 1.83-4.32) and transfer to NICU (OR 2.38; 95%CI 1.56-3.65).

Conclusions: Perinatal asphyxia is the initial concern in the IUGR fetus. Careful obstetric surveillance and timely delivery can prevent perinatal asphyxia and its clinical consequences. A neonatal resuscitative team should be available at delivery to improve neonatal outcome. Immediate management in delivery room should focus on the anticipation of a depressed infant, adequate resuscitation and insuring a normal physiologic transition.

1179

NEONATAL MORTALITY HISTORICAL SERIES AT A HOSPITAL IN THE SOUTH AREA OF THE SÃO PAULO CITY

J.R. Bertagnon¹, D.P. Escalhão²,
G.N.M. Almeida², H.A. Sayeg², R.T. Migliari²,
Y. Juliano³

¹Neonatologia, Hospital Geral do Grajaú - UNISA.,

²UNISA - CENEPS, ³Pós Graduação em Saúde Materno Infantil, UNISA, São Paulo, Brazil

Objective: To achieve the neonatal mortality coefficient (NMC) from 2005 to 2009 at the referred facilities and compare it to the 2002 data, when this hospital was not regarded yet as a risk maternity.

Method: A retrospective cross study based on the patients' records data for all live births (NB) during the 2005-2009 period ($N=9,966$) and compared by weight mortality specific coefficients to 2002 data, applying the specific coefficients for 2005 to 2009 using the Chi-Square test tool.

Results: There were 2,694 live births in 2002, out of which 32 NB had died ($NMC = 1.1.8\%$). For the period 2005 to 2009 these rates were 15.7% ; 19.55% ; 17.67% ; 17.0% and 14.2% , respectively.