n=6). Laser treatment was offered at a gestational age of 21 (17-26) weeks and delivery occurred at 29 (19-41) weeks.

Results: In 76%, at least one twin was live-born. The rate of fetal survival in donors was 60% and in recipients 66% ($p < 0.001$). 1 year-survival was 37% in donors and 44% in recipients ($p = 0.01$) and in 57% at least one twin was alive. There was no significant difference in survival when adjusted for Quintero stages, parity, maternal BMI or smoking whereas higher maternal age was associated with higher survival of both twins ($p < 0.05$).

Conclusions: Survival of at least one twin after FLOC was comparable to corresponding data of other international centra. Recipients are more likely to survive than donors. The finding of an increased survival rate with maternal age needs further investigation.

1215

EVALUATION OF BREASTFED INFANTS WITH RESPECT TO WHO GROWTH STANDARD: A META-ANALYSIS

D. Grathwohl¹, K. Macé², M.C. Fichot³, E. Spivey-Krobath³, P. Steenhout³

¹Bioanalytical Science, ²Nutrition & Health, Nestlé Research Center, Lausanne, ³Nestlé Nutrition, Vevey, Switzerland

Aim: Product development of infant formulas is done by randomized clinical trials, which are often containing an observational breastfeeding group for comparison purposes. These breastfeeding groups are tightly followed up in a clinical trial setting which was also the case when developing the WHO standard. Aim of this meta-analysis was to assess heterogeneity of the breastfed infants of the clinical trials with respect to the WHO standard.

Methods: Four studies were included in the meta-analysis, over all 192 infants. Population under investigation was healthy term born breastfed infants. The studies were conducted in Italy, Chile and two in France. Main outcome was length for age z-scores at 4 months of age, because WHO has communicated on length for age z-scores in their Multicentre Growth Reference Study (Brazil, Ghana, India, Norway, Oman and USA). The presented statistics is the random effect total and the 95% confidence interval which can interpret as a mean over all studies considering the heterogeneity of the contributing studies.

Results: The random effect total in length for age z-scores of the breastfed infants of the clinical trials was 0.04, [-0.19, 0.26]. For the WHO multicenter growth reference study the random effect total was estimated as -0.0164, [-0.1366, 0.1038].

Conclusion: We attribute the difference in totals and the increased heterogeneity to randomness.

1216

GROWTH OF INFANTS FED WITH NAN IS IN GOOD AGREEMENT WITH THE WHO GROWTH STANDARD: A META-ANALYSIS

D. Grathwohl¹, K. Macé², M.C. Fichot³, E. Spivey-Krobath³, P. Steenhout³

¹Bioanalytical Science, ²Nutrition & Health, Nestlé Research Center, Lausanne, ³Nestlé Nutrition, Vevey, Switzerland

Background and aims: The current NAN formula has an amino acid composition close to that of human breast milk. The objective of the meta-analysis was to evaluate growth of infants fed NAN with respect to WHO growth standard.

Methods: Five studies were included in the meta-analysis, over all 208 infants. Population under investigation was healthy term born infants fed with NAN formula. The studies were conducted in Australia, France and Italy. Main outcomes were z-score at 4 months of age. The American Academy of pediatrics is recommending 3 g/days as a clinically relevant treatment difference for weight gain. 3g/day corresponds to 0.5 z-scores at 4 months. The presented statistics is the random effect total and the 95% confidence interval which can interpret as a mean over all studies considering the heterogeneity of the contributing studies.

Results: The random effect total in weight, length, head circumference, BMI for age z-scores at 4 months was -0.11, [-0.29, 0.08], 0.13, [-0.13, 0.40], 0.29, [0.04, 0.54], -0.24, [-0.57, 0.10], respectively. This is within a boundary of ± 0.5 z-scores for weight and length, slightly above for head circumference and below for BMI.

Conclusion: Growth of infants fed with NAN is in good agreement with the WHO growth standard.