83

SUBARACHNOID SPACE QUANTIFICATION AS A SURROGATE MARKER OF BRAIN TISSUE INTEGRITY IN PRETERM INFANTS

G. Lodygensky¹, R. Recker², A. Merglen²,
M. Benders³, F. Lazeyras⁴, P.S. Hüppi⁵

¹Pediatrics, NICU, ²Pediatrics, University of Geneva, Geneva, Switzerland, ³Neonatology, Wilhelmina Children's Hospital, UMC Utrecht, Utrecht, The Netherlands, ⁴Radiology, ⁵Pediatrics, Division of Development and Growth, University of Geneva, Geneva, Switzerland

Background: Clinicians when evaluating MRI of preterm infants are in need for simple measurements that reflect actual brain tissue volumes. Subarachnoid space (SAS) measurement might be a reflection of white and grey matter integrity.

Methods: A cohort of 43 preterm infants was imaged at term equivalent by MRI (gestational age 29 ± 2 wks, birth weight 1295 ± 460 g). SAS was measured on a single T2-weighted coronal section at the level of the basilar truncus. SAS was then compared to actual brain tissue volumes quantified by the knn-algorithm using the same MRI.

Results: SAS correlated significantly with cerebrospinal fluid volume quantification (Pearson correlation 0.595, $p < 0.001$) but not with cortical grey matter, cerebral white matter or basal ganglia quantification.

Conclusion: SAS can't be used as a surrogate marker of brain tissue integrity.

Objective: To determine the effect of prolonged mechanical ventilation on cell-cycle regulation in a newborn rat model.

Methods: 7-day old Wistar rats were ventilated with low/moderate tidal volume and room air for 24 h. Following ventilation arterial blood gas analysis was performed and the animal was killed. Lung tissues were processed for histological or molecular/protein analysis to determine (1) number of proliferating cells by BrdU and phospho-Histone H3 staining and (2) gene and protein expression of cell cycle regulatory proteins by real-time PCR and Western blotting.

Results: After 24 h of ventilation arterial blood gas values were within normal range but alveolar formation was reduced. The number of proliferating cells was decreased (BrdU incorporation decreased from 10 to 3 % and phospho-histone H3 positive cells from 0.8 to 0.1% . Expression of CyclinD1 and E1 mRNA and protein was markedly down-regulated. Instead, protein expression of the CDK inhibitor p27^{Kip1} was significantly increased while its phosphorylation at site T157, T198 and T187 was decreased. Interestingly, CDK inhibitor p16^{Ink4a} levels were decreased by ventilation .

Conclusions: Prolonged mechanical ventilation causes cycle arrest of lung cells in newborn rats by increased transcription and nuclear localization of the CDK inhibitor p27^{Kip1}. These data may in part explain the arrest in alveolarization by mechanical ventilation of the neonatal lung.

85

BULLYING IN SCHOOLS - BELGRADE

S. Tomic

Health Centar Savski Venac, Belgrade, Serbia

Introduction: Contemporary world happenings followed by the financial and social crisis have resulted the increase of misery and poverty at the majority of the population, and especially of the most vulnerable groups as children - the young. All this together resulted the identity crisis, the value disarrangement that led to various types of battery and neglect and led to other forms of risky behavior. We witness every day the increase of all kinds of mistreating and especially of the bullying among the young. The aim is always the same - to hurt the victim.

84

PROLONGED VENTILATION CAUSES P27^{KIP1} INDUCED CELL CYCLE ARREST IN NEWBORN RAT LUNG

A.A. Kroon^{1,2}, J. Wang², Z. Huang²,
M. Kuliszewski², M. Post²

¹Pediatrics, Erasmus MC - Sophia Children's Hospital, Rotterdam, The Netherlands, ²Lung Biology Research, Physiology and Experimental Medicine Program, Hospital for Sick Children, Toronto, ON, Canada

Background: *In vitro* studies have demonstrated that mechanical stretch can inhibit lung cell proliferation. This inhibition of cell growth may lead to reduction of alveolar formation and contribute to the development of bronchopulmonary dysplasia.

Aim: Realizing the bullying in a school environment in one central Belgrade borough - Savski venac, trying to prevent it.

Methods: Realizing and analyzing the results of the survey that contained questions regarding bullying in a school environment. The survey covered 882 high school students in Savski venac (borough) - Belgrade.

Results and conclusions: The research showed that bullying happened among 39,6% of the young population. The bullying, no matter what kind of it, never happened to 60,4-95,1% of children. The most common form of bullying was in a form of quarrel try with friends 36,6%, it happened once 19,8%, 2-3 times 12,6%, and 4 or more times happened in 6% of situations. The most frequent bullying as a physical hurt happened in 95,1% of situations, of which once in 2% and in 0,9% two or more times. The frequent bullying was also in a form of mockery 36,8%, of which once in 17%, 2-3 times 10,8%, and over 4 times in 7,4%

86

NASAL CONTINUOUS AIRWAY PRESSURE VERSUS HELMET CONTINUOUS AIRWAY PRESSURE IN INFANTS WITH BRONCHIOLITIS: A CROSSOVER STUDY

J. Mayordomo-Colunga, C. Rey, A. Medina, M. Los Arcos, A. Concha, S. Menéndez, A. Vivanco

Pediatric Intensive Care Unit, Hospital Universitario Central de Asturias. Universidad de Oviedo, Oviedo, Spain

Objective: To compare the efficacy and safety of CPAP delivered by helmet interface (HI) and CPAP delivered by Infant-Flow® system (IF) with nasal prongs.

Methods: Prospective crossover study including infants < 3 months of age with refractory bronchiolitis. Patients with a modified Wood's clinical asthma score (M-WCAS) ≥ 4 or venous $\text{PCO}_2 \geq 60$ mmHg despite nebulized adrenalin were eligible. Infants were randomly allocated to receive CPAP by HI or IF during one hour, and then switched to the other CPAP system. Measurements were taken at baseline and after 60 and 120 minutes. After this period, CPAP was maintained with the system used in the second phase of the crossover period.

Results: Fifteen patients fulfilled inclusion criteria: 8 males, age 50.8 ± 15.7 (mean $\pm$ SD) days, weight 4.4 ± 0.7 kg. Nine infants received HI-CPAP in the first crossover phase and 6 received IF-CPAP initially. Baseline mean values were: M-WCAS 4.6 ± 1 points; venous PCO_2 57 ± 10.4 mmHg, heart rate (HR) 173 ± 15.2 beats/minute; respiratory rate (RR) 56.2 ± 11.2 breaths/minute. After 120 minutes of CPAP, clinical score, HR and RR fell 1.9 points, 15 beats/minute, and 8 breaths/minute, respectively ($p < 0.05$). At 60 minutes, improvement in clinical score (3 ± 0.9 vs. 2.8 ± 1), HR (166 ± 21.5 vs. 160.2 ± 13.3) and RR (47.3 ± 18.5 vs. 47.3 ± 11.1) was similar in both groups ($p > 0.05$). No adverse effects were recorded during the crossover period; later, two patients required intubation: one belonged to IF group and the other one to HI group.

Conclusions: HI-CPAP was as effective as IF-CPAP in infants with acute bronchiolitis. Safety of both systems seems to be similar.

87

AEROSOLIZED PFC IMPROVES PULMONARY FUNCTION AND GAS EXCHANGE IN A PRETERM LAMB MODEL

V. Mielgo¹, X. Murguía¹, C. Rey-Santano¹, E. Gastiasoro¹, E. Ruiz-del-Yerro¹, H. Lafuente¹, B. Loureiro², M.A. Gomez-Solaetxe³, F.J. Alvarez-Díaz¹, A. Valls-i-Soler²

¹Research Unit of Experimental Physiology,

²Neonatal Unit, Cruces Hospital, Vizcaya,

³Electronics and Electrotechnics High Technical School of Maritime Studies, University of Basque Country, Portugalete, Spain

Background: The aerosolization of PFC has emerged as a feasible alternative to the instillation for the experimental treatment of Respiratory Distress Syndrome.

Aim: To determine the effects of aerosolized perfluorocarbon (PFC) on pulmonary gas exchange and lung function in preterm lambs.

Methods: 12 preterm lambs (80-90%GE) were used. Lambs were randomly assigned to receive aerosolized PFC (10 ml/kg/h for 2h), delivered via an inhalation catheter, coupled to a jet nebuliser, and placed at the distal end of the endotracheal tube connected to a ventilator (PFC-aerosol) or not (CONTROL), both groups were maintained during 6h on IMV. Cardiovascular parameters, arterial partial pressure of CO_2 ($P_{a\text{CO}_2}$), pH, oxygenation