regions. A rabbit model of GMH/IVH demonstrated an increase in activated MG in the adjacent PVWM, which may cause or exacerbate WM injury.

Aim: To use post mortem (PM) brain MRI and immunohistochemistry to investigate MG distributions in the immature brain in the presence of GMH/IVH and to compare to controls with normally appearing brain.

Methods: Preterm (< 30 wk GA) PM cases were recruited with parental consent and ethics approval. Images were acquired at 3T with T1 and T2 weighted sequences. Iba1 and CD45 MG were counted in the anterior and posterior PVWM areas, centrum semiovale, germinal matrix (GM), subplate and cortex.

Results: 6 brains showed isolated GMH/IVH and 6 showed normal appearances. Total number of MG in the GM (Iba1=14.21±1.39 p< 0.05; CD45=11±4.70) and PVWM (Iba1=12.30±0.91 p< 0.05; CD45=4.8±1.52) were higher than other regions in the GMH/IVH group. GMH/IVH PVWM MG numbers were higher than in the normal group (p=0.005).

In the presence of IVH/GMH there was a marked increase in PVWM MG as seen in experimental models of IVH/GMH. These may be responsible for initiating or exacerbating an injurious process.

98

DOPAMINE AND CIRCULATION IN THE BRAIN: A STUDY IN NEWBORN PIGLETS

G.H. Hahn¹, C. Heiring², O. Pryds³, G. Greisen¹

¹Department of Neonatology, National University Hospital, Rigshospitalet, Copenhagen,

²Department of Paediatrics, Hillerød Hospital, Hillerød, ³Department of Neonatology, National University Hospital, Hvidovre Hospital, Copenhagen, Denmark

BACKGROUND AND Aims: Dopamine is widely used to treat hypotension in newborns. Data regarding its effect on the circulation in the brain, however, are ambiguous. To test for a direct effect on the brain, we estimated the magnitude of variability in arterial blood pressure (ABP) passing from systemic to cerebral circulation (i.e. frequency gain) during an ABP-challenge with dopamine in piglets.

Methods: Piglets (n=7, age 1-2 days) were anaesthetized and monitored with NIRS (cerebral oxygenation) during normo- and hypovolaemia. In both states, ABP was challenged with dopamine during a 20-minute stepwise dose-increment (20, 30, 40 and 50 µg/kg/min). Each infusion lasted 2½ minutes. Pauses lasting 2½ minutes were interpolated. Before and after the dopamine-challenge, ABP was challenged with six inflations of a thoracic aorta balloon. Each inflation lasted 30 seconds and pauses lasting 30 seconds were interpolated. Subsequently, frequency gain between ABP and cerebral oxygenation was calculated (frequency range: 0.003-0.04 Hz).

Results: Mean variability in ABP was 36±4.3 mm Hg during balloon challenge and 17±3.2 mm Hg during dopamine challenge (mean ±SEM). Mean gain between ABP and cerebral NIRS was 0.11±0.019 during ABP-challenge with the aorta balloon and 0.12±0.029 µM/mm Hg during ABP-challenge with dopamine (P=0.86). The mean difference was 0.01 [95%-CI: -0.07 to +0.09]

Conclusions: ABP-variability induced by dopamine and by inflating an aortic balloon had similar effects on cerebral oxygenation. This suggests that dopamine has no direct constricting or dilating effect on the cerebral blood vessels or that such effects are counterbalanced by changes in cerebral oxygen uptake.

99

CORRELATION OF NON-INVASIVE SYSTOLIC AND MEAN BLOOD PRESSURE (BP) MEASUREMENTS WITH ECHOCARDIOGRAPHIC HAEMODYNAMIC ASSESSMENT

S. Gupta¹, J.P. Wyllie²

¹Neonatal Paediatrics, University Hospital of North Tees, Stockton on Tees, ²Neonatal Paediatrics, The James Cook University Hospital, Middlesbrough, UK

Objective: Systolic and mean BP values are routinely utilised to define hypotension on neonatal units. There is however no data correlating systolic or mean BP with cardiac haemodynamics in newborn infants.

Methods: In a randomised trial¹, 71 infants received either two transfusions of 10ml/kg or single transfusion of 20ml/kg. Clinical and echocardiographic haemodynamic measurements