

Results: S100B maternal and neonatal blood concentrations were significantly higher at all monitoring time-points in the SSRI ($P < 0.001$, for all), whilst proteins' concentrations in healthy mothers and newborns were within normality ranges.

Conclusions: The elevated S100B protein concentrations in maternal and newborn bloodstreams suggest that SSRI exposure, in agreement to adults' findings, can exert CNS side-effects both in intrauterine and in post-natal periods. Further investigations aimed at investigating short/long term neurological sequelae in these patients are needed.

148

BRANCHED CHAIN AMINO ACID REQUIREMENTS FOR TERM NEONATES

F. Maingay-de Groof¹, L. Huang¹, G.J. Voortman¹, C. Chao², Y. Huang², J.B. van Goudoever¹

¹Neonatology, Pediatrics, Erasmus MC Sophia, Rotterdam, The Netherlands, ²Neonatology, Pediatrics, Fudan Children's Hospital, Shanghai, China

Background: Dietary intake should meet the requirement to obtain an optimal growth and neurodevelopment in the neonate. The essential branched chain amino acids (BCAAs), leucine, isoleucine and valine, are mainly used for incorporation into body protein. Current recommended BCAA requirements for infants 0-6 months (respectively 156, 88 and 87 mg·kg⁻¹·d⁻¹, ratio 1.8:1:1) are based on the amino acid content of human milk. However, human milk fluctuates in composition during lactation and even during a feeding, while milk consumption rate varies widely as well. Questions remain on the validity to use mean amino acid composition of human milk to determine requirements.

Objectives: To quantify the requirement of leucine, isoleucine and valine in term neonates using the Indicator Amino Acid Oxidation method.

Design: Enterally fed term infants received randomly graded intakes of leucine (15-500 mg·kg⁻¹·d⁻¹), isoleucine (5-216 mg·kg⁻¹·d⁻¹) and valine (5-236 mg·kg⁻¹·d⁻¹). Breath samples containing ¹³CO₂ were collected during L-[1-¹³C]phenylalanine (indicator amino acid) administration, measured by isotope ratio mass spectrometry and analysed using a biphasic regression crossover analysis.

Results: 83 term Asian neonates (birth weight: 3.29 ± 0.4 kg, gestational age: 39.4 ± 1.3 wks, postnatal age: 12.6 ± 5.1 d) were included. The mean requirement (at breakpoint) for leucine, isoleucine and valine was respectively 140, 105 and 110 mg·kg⁻¹·d⁻¹ and the upper 95% confidence interval was 240, 152 and 165 mg·kg⁻¹·d⁻¹.

Conclusion: The requirements of the individual BCAAs are almost twice the current recommendations. A Leu:Ile:Val ratio of 1.3:1:1 is more appropriate in term formula.

149

GESTATIONAL AGE PATTERNS OF FETAL AND NEONATAL MORTALITY RATES: THE EURO-PERISTAT PROJECT

A. Mohangoo¹, S. Buitendijk¹, J. Zeitlin², and the EURO-PERISTAT Network

¹Prevention and Care, TNO Quality of Life, Leiden, The Netherlands, ²Epidemiological Research Unit on Perinatal and Women's Health, INSERM UMR S16, Paris, France

Background: The recently published European Perinatal Health Report showed wide variability in perinatal mortality rates between European countries. We investigated the gestational age patterns of mortality in order to better understand differences between low versus high mortality countries.

Setting: The Euro-Peristat project developed a list of valid and reliable indicators for monitoring and evaluating perinatal health, including fetal and neonatal mortality. Data from 2004 on 29 countries/regions were analyzed.

Results: The fetal mortality rate ranged from 2.6 per 1000 births in Slovakia to 9.1 in France (weighted average of 5.4 per 1000 births) and the neonatal mortality rate ranged from 1.6 per 1000 live births in Cyprus to 5.7 in Latvia (weighted average of 3.0 per 1000 live births). In some countries, fetal mortality rates declined dramatically after excluding extremely preterm births (< 28 weeks), while elsewhere rates stayed stable. The exclusion of the extremely preterm births hardly influenced the variability in neonatal mortality rates, although a large decline was observed for the Netherlands, where active intervention is very conservative before 26 weeks of pregnancy. We did not find that countries with low mortality rates had higher proportions of extremely preterm births (which could be considered less