Conclusions: Prediction of neonatal neurological morbidity might be greatly improved by fetal cardiovascular evaluation, influencing prenatal and neonatal management. The association of prenatal cardiovascular parameters with long-term neurodevelopment merits further investigation.

201

ASSESSMENT OF GESTATIONAL AGE IN VERY PRETERM NEONATES USING CEREBELLAR MEASUREMENTS AT CRANIAL ULTRASOUND - WHAT IS THE BEST APPROACH?

A. Graca¹, K. Cardoso¹, F. Cowan²

¹Neonatology - Department of Paediatrics, Hospital de Santa Maria / University of Lisbon, Lisbon, Portugal, ²Department of Paediatrics and Neonatal Medicine, Imperial College School of Medicine / Hammersmith Hospital, London, UK

Background and aims: Clinical assessment of gestational age (GA) can be challenging. Several ultrasound approaches to estimate GA using cerebellar measurements have been reported, claiming to be simpler and more accurate than clinical assessment. We compare the accuracy of 3 previously described cerebellar measurements for estimating the GA in VLBW infants.

Methods: We studied prospectively VLBW infants under 32 weeks GA defined by certain dates, IVF date or early fetal ultrasound. We excluded infants with IUGR, neurological problems or scan abnormality except isolated GLH/transient flares. Ultrasound acquisition and cerebellar measurements were made by author AG. Measurements of the vermis anterior-posterior diameter (APD, Cuddihy 1999), transverse cerebellar diameter via anterior (TCDa, Makhoul 2000) and mastoid fontanelles (TCDm, Davies 2001) were obtained. Estimated GA was calculated using published equations, and compared to known GA using intraclass correlation coefficient (ICC).

Results: We studied 60 infants. Mean birth GA was 28.4 weeks and mean postnatal age at scan was 1 week. ICC was 0.801 for APD, 0.123 for TCDa and 0.802 for TCDm ($p < 0.001$). Best estimates of GA were obtained with TCDm (difference SD 1 week) and APD (difference SD 1.3 weeks).

Conclusions: APD and TCD gave good estimates of GA and can be recommended. We obtained similar measurements of TCD via the two approaches, but only the equation for the mastoid fontanelle gave

a good estimate of GA with excellent ICC. Studies assessing the use of cerebellar measurements for estimating GA in growth restricted, neurologically abnormal and older infants are needed.

202

POST-HAEMORRHAGIC VENTRICULAR DILATATION AND ADC MEASUREMENTS IN THE WHITE MATTER IN PRETERM BORN INFANTS AT TERM EQUIVALENT AGE

E. van Mierlo, M. Brouwer, B. van Kooij, F. Groenendaal, L. de Vries, M. Benders

Neonatology, Wilhelmina Children's Hospital / UMC, Utrecht, The Netherlands

Objective: Post-haemorrhagic ventricular dilatation (PHVD) might cause additional white matter (WM) injury due to increased pressure on surrounding tissue. The extent of associated WM injury seems to be the main predictor for an adverse neurodevelopmental outcome.

Aim: Assessing WM injury in preterm born neonates with PHVD at term-equivalent age (TEA) by measuring the ADC on diffusion-weighted MRI (DW-MRI).

Methods: In this retrospective patient-controlled study, 23 preterm infants with PHVD (median GA 27.3 weeks (range 25.6-30.6)), admitted to our neonatal intensive care unit (NICU), were matched to 23 control patients for gender and GA (median GA 27.4 (range 25.3-30.9)). DW-MRI was performed in all neonates at TEA. Regions of interest were drawn manually on the ADC-map with equal size and location in the frontal, parietal and occipital WM bilaterally.

Results: Results are presented in table 1. PHVD was associated with slightly higher ADC-values in the occipital WM ($p < 0.05$). No significant differences in ADC-values of the frontal and parietal WM were observed between both groups.

ADC $\times 10^{-3} \text{mm}^2/\text{s}$ (mean $\pm$ SE)	PHVD N=23	No PHVD N=23	p-value
Frontal	1.566 $\pm$ 0.020	1.572 $\pm$ 0.023	0.830
Parietal	1.632 $\pm$ 0.022	1.594 $\pm$ 0.025	0.288
Occipital	1.549 $\pm$ 0.024	1.468 $\pm$ 0.024	<0.05

[Table 1]