later. Data were analysed by multi-level analysis and Mann-Whitney-U-test.

Results: Nine neonates (median age: 1 day [range:0-18]) underwent BAS and four neonates (median age: 4 days [range:0-6]) did not. R_cSO_2 increased after BAS (from mean 37% to 49% to 64% [$p=0.000$]), as did $tcSaO_2$ (from mean 68% to 81% to 84% [$p=0.000$]). FTOE decreased after BAS (from mean 0.46 to 0.40 to 0.25 [$p=0.000$]). Neonates who did not need BAS showed higher baseline r_cSO_2 and $tcSaO_2$ compared to neonates who did (median 57% vs 35% [$p=0.020$] and 88% vs 77% [$p=0.034$], respectively). After BAS, there were no differences between both groups.

Conclusion: In neonates with TGA in need of BAS, r_cSO_2 increased and FTOE decreased following BAS. This suggests improved cerebral oxygenation, which, possibly, protects against hypoxic-ischemic brain-injury.

222

SECOND COURSE OF IBUPROFEN FOR PDA : COMPARISON OF 10-5-5 VS 20-10-10 MG/KG IN INBORN INFANTS < 28 WKS GA

X. Durrmeyer, E. Millot, F. Decobert, C. Danan,
I. Layouni, G. Dassieu

*Réanimation Néonatale, Centre Hospitalier
Intercommunal de Créteil, Créteil, France*

Background and aims: Recent pharmacological data suggest increased ibuprofen doses for treating patent ductus arteriosus (PDA). We evaluated the effectiveness and tolerance of a double dosed second ibuprofen course in our population.

Methods: In our unit PDA management relies on systematic and repeated echocardiographic exams seeking strict treatment criteria (ductal diameter, left atrium/aortic root, left pulmonary artery diastolic blood flow). Persistent echocardiographic criteria after a first ibuprofen course induce a second course if not contra-indicated.

We conducted a monocenter retrospective study by comparing two time periods: November 2004 to March 2006 and April 2006 to November 2009 where the second ibuprofen course doses respectively were 10-5-5 (IBU1) and 20-10-10 (IBU2) mg/kg. Baseline characteristics, treatment incidence, ductal closure and complications were compared between IBU1 and IBU2. Term-based subgroup analysis was performed (< or $\geq$ 26 wks GA).

Results: IBU1 and IBU2 populations were comparable. Ductal closure was more frequent for IBU2 (23/55,42%) than for IBU1 (3/23,13%) ($p=0,02$). Subgroup analysis confirmed this result only for infants $\geq$ 26 wks GA.

Incidence of mortality and common morbidities were comparable between IBU1 and IBU2. Among infants < 26 wks GA we observed a higher, although not significant, incidence of "death or BPD" for IBU2 vs IBU1 (37% vs 9%).

Conclusions: This is the largest report of increased ibuprofen doses in infants < 28 wks GA with systematic echocardiographic assessment. In this study risk/benefit balance supports the use of a double-dosed second course of ibuprofen for PDA treatment only in infants $\geq$ 26 wks GA.

223

THE DEFINITION OF A HEMODYNAMIC SIGNIFICANT DUCT IN RANDOMISED CONTROLLED TRIALS, A SYSTEMATIC LITERATURE REVIEW

I. Zonnenberg¹, K. de Waal²

*¹Neonatology, VU Medical Centre, Amsterdam,
The Netherlands, ²Neonatology, John Hunter
Hospital, Newcastle, NSW, Australia*

The ductus arteriosus is associated with morbidity in preterm infants. Treatment is prescribed for a hemodynamically significant duct (HSDA), but its definition varies. We systematically reviewed published clinical and ultrasound definitions (US) of a HSDA.

Methods: Pubmed and the Cochrane library were searched for randomised trials on ductal closure or trials evaluating the prevention of a symptomatic duct developing.

Results: Fifty nine trials were included in our review. Thirty nine trials used clinical and US criteria to define a HSDA, 5 trials clinical criteria only, 10 trials US criteria only, 2 trials clinical or US criteria and in 3 trials the criteria were not mentioned. Re-entry criteria for ductal assessment was clinical only (2), clinical followed by ultrasound (39), US only (3) or it was not mentioned (15). Clinical criteria were defined in 41 trials, with murmur (29), bounding pulses (27), hyperdynamic precordium (24), cardiomegaly (19) and respiratory status (19) most reported on. US criteria were defined in 50 trials, with LA/Ao ratio > 1.15 to 1.7 being most reported (33). Other US

criteria used were flow pattern in the descending aorta (12) and in the pulmonary artery (13), or any left-right shunt (9). Ductal diameter > 1.5 mm was used in 8 more recent trials, but in only one trial as solitary US criteria.

Conclusion: A large variation in HSDA definition exists. We should try reaching an international consensus on the definition of a HSDA, and evaluate if clinical signs are correlated with the mentioned US parameters or DAd.

224

NOVEL ULTRASOUND DILUTION TECHNIQUE DETECTS LEFT-TO-RIGHT SHUNTS WITH HIGH ACCURACY IN CHILDREN

M. Boehne¹, M. Baustert¹, C.M. Happel¹, S. Schoff¹, D. Hohmann¹, V. Paetzel¹, W.A. Osthaus², M. Sasse¹, A. Wessel¹, H. Bertram¹

¹Paed. Cardiology and Intensive Care Medicine,

²Clinic for Anaesthesiology and Intensive Care Medicine, Hannover Medical School, Hannover, Germany

Objectives: Recently introduced paediatric cardiac output monitor (COstatus, Transonic Systems Inc. Ithaca, NY) also measures blood volumes and detects cardiac shunts. The current study assessed its ability to identify left-to-right shunts and measured associated blood volumes during cardiac catheterisation in children.

Patients and Methods: Cardiac catheterisation was performed in 20 children and adolescents (3.5 kg- 102 kg). Shunts and pulmonary (Qp) versus systemic blood flow (Qs) ratio were determined by routine oximetry. Subsequently three measurements with COstatus were performed while the monitor software announces the recognition of left-to-right shunt. Dilutions curves were recorded and validated.

Results: Left-to-right shunts due to various types of cardiac defects were detected in 6 patients by oximetry. Applying COstatus diagnostic accuracy was very convincing, with a sensitivity of 100% (6/6) and a specificity of 92.8% (13/14). The threshold for determination a left-to-right shunt was an Qp/Qs ratio of 1.3. In one case of a hypertrophic obstructive cardiomyopathy the distortion of the dilution curve led to a false positive result. A significant difference was detected for means of central blood volume index between patients with left-to-right shunts

and controls with no cardiac shunts (95% CI; 24.9 versus 18.2 ml/kg; $P < 0.05$).

Conclusion: Novel ultrasound dilution technique identifies left-to-right shunts in children with a high sensitivity and detects even small shunts with an Qp/Qs ratio of 1.3. Raised central blood volume index corresponds well with clinical features in patients with left-to-right shunts and may become helpful a diagnostic parameter in the future.

225

ORAL VS INTRAVENOUS IBUPROFEN FOR TREATMENT OF PATENT DUCTUS ARTERIOSUS

I. Mungan Akin¹, E. Okulu¹, B. Atasay¹, S. Arsan¹, E. Tutar², S. Atalay², T. Turmen¹

¹Department of Pediatrics, Division of Neonatology,

²Department of Pediatrics, Division of Cardiology, Ankara University Medical Faculty, Ankara, Turkey

Background and aims: The aim is to evaluate and compare the efficacy and side effects of oral and intravenous(IV) ibuprofen for treatment of PDA in preterm infants.

Methods: A prospective, randomized controlled trial was conducted in premature neonates with gestational ages(GA) < 32 weeks, birth weight(BW) < 1500 g, at Ankara University NICU between January 2008 to January 2010. Infants who had defined symptoms for PDA together with echocardiographically demonstrated ductus and presymptomatic infants with a ductus diameter/kg $>1,5$ or with a left atrium/aorta ratio $>1,5$ detected on routine echocardiogram in the 2nd day of life were included in the study. Three doses of either oral or IV ibuprofen were randomly given at a dosage of 10,5,5mg/kg every 24 hours. Daily physical examination, serial laboratory evaluation and echocardiogram were used to evaluate symptomatic PDA, treatment outcomes, complications and side effects.

Results: Thirty-three infants were recruited during two years period and randomly assigned into the groups. Twenty of them received IV ibuprofen. The GA, BW, presence of antenatal corticosteroid, APGAR scores, use of surfactant, ductus diameters, La/Ao ratio of the two groups were similar. Closure, reopening and need for surgical ligation didn't differ between groups. There were no significant difference in the prevalence of NEC, renal dysfunction and mortality between two groups.