criteria used were flow pattern in the descending aorta (12) and in the pulmonary artery (13), or any left-right shunt (9). Ductal diameter > 1.5 mm was used in 8 more recent trials, but in only one trial as solitary US criteria.

Conclusion: A large variation in HSDA definition exists. We should try reaching an international consensus on the definition of a HSDA, and evaluate if clinical signs are correlated with the mentioned US parameters or DAd.

224

NOVEL ULTRASOUND DILUTION TECHNIQUE DETECTS LEFT-TO-RIGHT SHUNTS WITH HIGH ACCURACY IN CHILDREN

M. Boehne¹, M. Baustert¹, C.M. Happel¹, S. Schoff¹, D. Hohmann¹, V. Paetzel¹, W.A. Osthaus², M. Sasse¹, A. Wessel¹, H. Bertram¹

¹Paed. Cardiology and Intensive Care Medicine,

²Clinic for Anaesthesiology and Intensive Care Medicine, Hannover Medical School, Hannover, Germany

Objectives: Recently introduced paediatric cardiac output monitor (COstatus, Transonic Systems Inc. Ithaca, NY) also measures blood volumes and detects cardiac shunts. The current study assessed its ability to identify left-to-right shunts and measured associated blood volumes during cardiac catheterisation in children.

Patients and Methods: Cardiac catheterisation was performed in 20 children and adolescents (3.5 kg- 102 kg). Shunts and pulmonary (Qp) versus systemic blood flow (Qs) ratio were determined by routine oximetry. Subsequently three measurements with COstatus were performed while the monitor software announces the recognition of left-to-right shunt. Dilutions curves were recorded and validated.

Results: Left-to-right shunts due to various types of cardiac defects were detected in 6 patients by oximetry. Applying COstatus diagnostic accuracy was very convincing, with a sensitivity of 100% (6/6) and a specificity of 92.8% (13/14). The threshold for determination a left-to-right shunt was an Qp/Qs ratio of 1.3. In one case of a hypertrophic obstructive cardiomyopathy the distortion of the dilution curve led to a false positive result. A significant difference was detected for means of central blood volume index between patients with left-to-right shunts

and controls with no cardiac shunts (95% CI; 24.9 versus 18.2 ml/kg; P< 0.05).

Conclusion: Novel ultrasound dilution technique identifies left-to-right shunts in children with a high sensitivity and detects even small shunts with an Qp/Qs ratio of 1.3. Raised central blood volume index corresponds well with clinical features in patients with left-to-right shunts and may become helpful a diagnostic parameter in the future.

225

ORAL VS INTRAVENOUS IBUPROFEN FOR TREATMENT OF PATENT DUCTUS ARTERIOSUS

I. Mungan Akin¹, E. Okulu¹, B. Atasay¹, S. Arsan¹, E. Tutar², S. Atalay², T. Turmen¹

¹Department of Pediatrics, Division of Neonatology,

²Department of Pediatrics, Division of Cardiology, Ankara University Medical Faculty, Ankara, Turkey

Background and aims: The aim is to evaluate and compare the efficacy and side effects of oral and intravenous(IV) ibuprofen for treatment of PDA in preterm infants.

Methods: A prospective, randomized controlled trial was conducted in premature neonates with gestational ages(GA) < 32weeks, birth weight(BW) < 1500g, at Ankara University NICU between January 2008 to January 2010. Infants who had defined symptoms for PDA together with echocardiographically demonstrated ductus and presymptomatic infants with a ductus diameter/kg>1,5 or with a left atrium/aorta ratio>1,5 detected on routine echocardiogram in the 2nd day of life were included in the study. Three doses of either oral or IV ibuprofen were randomly given at a dosage of 10,5,5mg/kg every 24 hours. Daily physical examination, serial laboratory evaluation and echocardiogram were used to evaluate symptomatic PDA, treatment outcomes, complications and side effects.

Results: Thirty-three infants were recruited during two years period and randomly assigned into the groups. Twenty of them received IV ibuprofen. The GA, BW, presence of antenatal corticosteroid, APGAR scores, use of surfactant, ductus diameters, La/Ao ratio of the two groups were similar. Closure, reopening and need for surgical ligation didn't differ between groups. There were no significant difference in the prevalence of NEC, renal dysfunction and mortality between two groups.