(Z-score). Female DM1 patients had a significant higher BMI Z-score (0.51 ± 0.84 vs. -0.11 ± 0.95 ; $p=0.01$) and a significant lower trabecular BMD (175.7 ± 30.6 vs. $195.2 \pm 32.7 \text{ mg/cm}^2$; $p=0.03$) and Z-score (-0.55 ± 0.87 vs. 0.03 ± 0.92 ; $p=0.02$). Only 2 DM1 females and none of the males had a trabecular BMD Z-score < -2 . Mean HbA1c was significantly higher in female patients (8.68 ± 1.19 vs. $7.83 \pm 1.24\%$; $p=0.02$). HbA1c ranged between 6.0-10.8% and was negatively correlated with trabecular BMD Z-score ($r=-0.15$; $p=0.3$), but correlation was not significant.

Conclusions: Whereas diabetes type 1 females, compared to healthy controls, have a lower trabecular bone mineralization at the end of puberty, the majority of adolescent DM1 patients have a trabecular BMD within normal range.

	Control Group	Group A	Group B	Group C
No	19790	40	131	681
Primipara (%)	45.9	39.5	45.4	47.5
Maternal age (years, mean $\pm$ SD)	27.36 ± 5.72	$30 \pm 3.8^*$	$33.3 \pm 5.36^*$	$30.4 \pm 5.7^*$
Male sex (%)	52.1	57.5	53.4	51.7
GA < 32 wks (%)	6.1	2.5	16*	5
GA 33-36 wks (%)	10.7	22.5*	29*	12.9
GA 37-40 wks (%)	83.2	75	55	82.1
Caesarian section (%)	32.8	72.5*	70.2*	46.4*
IVF (%)	3.5	0	14.5*	6
Hypertensive disorders (%)	3.3	12.5*	11.5*	5.5*
Term neonates: GA (wks) (mean $\pm$ SD)	39.1 ± 1.1	$37.9 \pm 0.85^*$	$37.8 \pm 0.84^*$	$38.5 \pm 1^*$
BW (g) (mean $\pm$ SD)	3307 ± 461	$3666 \pm 726^*$	3300 ± 458	3279 ± 469

261

THE EPIDEMIOLOGY OF DIABETES IN PREGNANCY AND MATERNAL CHARACTERISTICS IN NORTHERN GREECE

E. Diamanti, K. Sarafidis, V. Soubasi, K. Agoropoulou, K. Anegronou, E. Agakidou, A. Marioglou, V. Drossou-Agakidou

1st Dept. of Neonatology, Aristotle University of Thessaloniki, Thessaloniki, Greece

Aims: To determine the prevalence of diabetes in pregnancy and evaluate maternal characteristics according to the type of diabetes.

Methods: We prospectively studied the maternal characteristics of all neonates who were born in our perinatal center and were treated in our department during the period 1996-2007. Infants of diabetic mothers were classified in three groups: Group A: mothers with preexisting diabetes, Group B: mothers with gestational diabetes, managed with insulin, Group C: mothers with gestational diabetes, managed with diet. Non diabetic mothers served as control.

Results: Overall, 20.643 live neonates from 19.791 pregnancies were studied. Of them, 852 were complicated by diabetes. The incidence of DP was 1.86% from 1996 to 2000 and 5.57% from 2001 to 2007. Maternal-neonatal characteristics are shown in the table (* $p < 0.001$).

[Maternal-neonatal characteristics]

Conclusions: The incidence of DP increased during the last 6 years. The prevalence of IVF is significantly elevated only in gestational diabetes treated with insulin. Also, in term neonates, birth weight was significantly higher only in women with preexisting diabetes. These results, however, need to be confirmed in other studies.

262

MEDICAL AND NURSE'S PRACTICES REGARDING ON END OF LIFE, IN A SOUTHERN BRAZILIAN PEDIATRIC INTENSIVE CARE UNIT

P. Lago, J. Piva, C. Nilson

Pediatric, PUCRS, Porto Alegre, Brazil

Objective: Evaluate medical and nurse's practices offered in last 24 hours of life of children that died in 2 Brazilian PICU.

Methods: The prospective study was based on: 1) medical chart's analysis of children that died between April- September/2008; 2) Semi- structured interview with the nurses (17). Researchers fielded out 3 protocols looking for data about interventions and care offered in the last 24 hours of life as well as the nurse's participations on EOL decision. According to the treatment offered in the last moments of life, children were classified in 2 groups: "full CPR" and "no CPR". The groups were compared.

Results: 37 deaths were identified in this period (excluded 3 cases- brain death). The CPR was not performed in 50% of children. In 10 of 17 patients not reanimated, the plan for LSL was not recorded in the medical charts. In 80% patients of the “no CPR” group, the vasoactive drugs infusion and ventilator parameters were maintained/increased without differences when compared to “full CPR” ($p=1,0$ e $p=0,6$). The nurse’s practices evaluated using NAS score in the last 24 hours of life didn’t differ even when the patient wasn’t reanimated. When nurses were asked about participation on EOL decision, only 30% confirmed it.

Conclusion: Brazilian studies have demonstrated that EOL practice has been a process centralized on the medical perspectives. We observed that nurses and medical attitudes before death didn’t change even when the patients were considered terminal illness and the nurses rarely participated on the decision making process.

263

BENEFITS OF STRUCTURED MULTI DISCIPLINARY MEDICAL ETHICAL DECISION MAKING

G. van Blijderveen, C. de Boer, M. Williams

Pediatrics / Neonatology, Erasmus MC Sophia Childrens Hospital, Rotterdam, The Netherlands

Background/aim: Until 2008, medical ethical decision making (MEDM) in the NICU lacked structure. Moral deliberation and effecting ethical decisions were not up to standard. A guideline to improve MEDM was developed, and evaluated one year after implementation.

Method: The guideline consisted of multidisciplinary meetings presided by an independent chair, and uniform preparation and report. Meetings were structured following a five phase model: 1) exploration, 2) agreement on the ethical question, 3) analysis, 4) evaluating possible solutions, decision-making, 5) planning actions. Initially, health care workers (HCW) of the NICU were trained in ethical/legal aspects and practiced the method with a historical patient case. Before the training, HCW completed a 15-item questionnaire on ‘structure’, ‘role participants’, ‘parental representation’, ‘problems, solutions and deliberation’, and ‘decisions and documentation’. One year and 22 structured meetings after introduction, the same questionnaire was completed again.

Results: Response before and after introduction of the guideline was 91/114 (80%) and 85/123 (69%) respectively. Nurses and physicians were equally represented in the two samples ($N=63/63$, and $N=19/14$; $X^2=.601$; $p=.438$). Evaluation of the guideline showed improvement in ‘structure of MEDM’ ($p<.000$ for all 5 items), ‘role participants’ ($p<.000$ for all 3 items), ‘parental representation’ ($p<.000$, for 1 item), and ‘problems, solutions and deliberation’ ($p<.000$, for 2 items; the 3rd item treatment options did not improve, $p=.932$). ‘Decisions and documentation’ showed no improvement (3 items, $p=.215$, $p=.412$ and $p=.066$ respectively).

Conclusions: Ethical deliberation significantly improved after implementation of structured MEDM. Ongoing attention will be paid to ‘documentation of the decisions’.

264

SEMI-STRUCTURED INTERVIEWS WITH GRIEVING PARENTS AFTER THE DEATH OF THEIR NEWBORN INFANT IN THE INTENSIVE CARE UNIT

I. Wermuth, A. Schulze

Division of Neonatology, Center for Perinatal Medicine Grosshadern, Dr. von Hauner Children’s Hospital, Munich, Germany

Background/aims: To obtain empiric data on parent’s experience of death of their newborn infant and to identify variables influencing the grief reaction.

Methods: Parents of all infants who died during a 5-year-period were asked to complete a standardized questionnaire and/or to participate in a semi-structured interview. Interviews were recorded and later transcribed. The questionnaire (242 questions, 21 pages) included the Perinatal Grief Scale (PGS).

Results: 31 mothers and 19 fathers of 48 children participated. The interviews lasted 2.6 hrs on average. PGS-Scores were higher (i) for mothers vs. fathers, (ii) if the couple had previous children, (iii) with increasing time after the infant’s death.

Grief intensity was not higher if there was parental involvement in the decision to withdraw life-sustaining therapy. 95% of parents involved in a decision to restrict intensive care felt their involvement was adequately handled by medical staff. 92% of these