Conclusion: Childhood AAC may be secondary to a variety of etiologies, especially during the course of infectious diseases. Presence of shock and a low value of fibrinogen are predictable factors for mortality in childhood AAC.

275

NON-ALCOHOLIC FATTY LIVER DISEASE AND STEATOHEPATITIS IN TAIWANESE CHILDREN: THE PREVALENCE AND SERUM RETINOL BINDING PROTEIN 4 LEVEL

Y.-J. Yang¹, S.-C. Huang²

¹Department of Pediatrics, National Cheng Kung University and Hospital, ²Department of Pediatrics, Kuo General Hospital, Tainan, Taiwan R.O.C.

Subjective: The aims of this study were to investigate the prevalence of non-alcoholic fatty liver disease (NAFLD) and non-alcoholic steatohepatitis (NASH) in schoolchildren with various body mass index (BMI). Moreover, the correlation between serum RBP4 level and liver injury was investigated.

Methods: 847 schoolchildren aged 4 to 12 years were evaluated by anthropometric measurements. The serum glucose, aspartate aminotransferase (AST), alanine aminotransferase (ALT), triglycerides and cholesterol levels were measured after an 8-hour overnight fast. NAFLD was diagnosed as fatty infiltrates of liver in sonogram and NASH was defined as NAFLD with an elevated ALT level.

Results: The prevalence of overweight (85th % < BMI < 95th %) and obesity (BMI > 95th %) in schoolchildren was 14.8% and 11.4%, respectively. The mean age and male-to-female ratio were significantly higher in children with obesity than those with normal weight (P=0.001). The obese children had a higher rate of abnormal AST and ALT (P< 0.001) and higher fasting triglyceride (P< 0.001) and glucose (P=0.04) levels than those with normal weight. Among 245 children with ultrasound examinations, fatty liver disease was identified in 2.5%, 20.6%, and 68.2% of children with normal weight, overweight, and obesity, respectively. Nine (40.9%) subjects were diagnosed as NASH in children with obesity. The serum RBP4 level was significantly higher in children with NAFLD and NASH than that in normal (P< 0.05).

Conclusions: Childhood obesity is closely related to male gender, increase of age, fasting triglycerides, and glucose levels. Moreover, the serum RBP4 level was parallel to the liver injury in children.

276

GASTROESOPHAGEAL REFLUX DISEASE IN INFANT: RELATIONSHIP BETWEEN MULTICHANNEL INTRALUMINAL IMPEDANCE-PH MONITORING AND CLINICAL OUTCOME

F. Cresi, E. Locatelli, C. Marinaccio, G. Grasso, E. Castagno, L. Silvestro

Pediatrics, University of Turin, Turin, Italy

Background and aims: combined multichannel intraluminal impedance and pH-monitoring (MII/pH) is a new technique identifying refluxes irrespective of acidity and detecting their duration, proximal extent and pH. These features are important when studying gastroesophageal-reflux-disease (GERD) in infants, in which weakly acid refluxes are prevalent. Clinical application of MII/pH is uncommon yet, owing to lack of reference values and difficult interpretation of the variables obtained. Aim of the present study was to investigate the relationship between MII-pH results and clinical outcome on a sample of GERD infants.

Methods: infants (age: 0-3 months) with GERD symptoms were studied with MII/pH and submitted to a follow-up consisting in clinical examinations and structured interviews to parents at 3-6-9-12-18-24-36 months to evaluate the presence of symptoms and the therapy effects.

Results: 54 patients completed the follow-up (32M; age 33.69±21.78 days; weight 3465.13±791.08 gr.; length 52.08±3.55 cm). 22 and 15 patients were still symptomatic at the age of 6 and 9 months, without difference in MII/pH values between symptomatic and non symptomatic infants. The 12 infants that were still symptomatic at the age of 12 months showed a higher bolus exposure index (BEI) with respect to the healed patients (2.63±1.72% vs 1.72±1.05; p=0.029).

Conclusions: MII-pH in first months of life could be useful to identify patients with high risk of presenting GER symptoms for more than 1 year. BEI is a pH independent variable with a significant relationship with symptoms duration. Our data highlight the clinical relevance of MII/pH-detected weakly acidic refluxes in GERD infant.