no alarm messages (low IQ signal, low perfusion, sensor off, ambient light). Infants were excluded if they received oxygen treatment or assisted ventilation in this time or they had a congenital anomaly that could interfere with transition. We used these data to construct centile charts.

Results: Data from 468 infants, of 25 to 42 weeks gestation, were used. At two minutes after birth the 10th, 50th and 90th centiles were 46%, 73% and 91% and at five minutes 73%, 89%, and 97%. Deciding the appropriate centile to target for oxygen treatment is a balance between treating those who need it and not giving it to those who don't. If the 50th centile was used this would lead to half of the normal population receiving oxygen unnecessarily. Choosing a lower centile (e.g. 10th) may be more appropriate.

Conclusions: The appropriate saturation target for infants receiving oxygen during resuscitation remains uncertain. Future trials could compare outcomes of infants resuscitated using different SpO₂ targets.

342

CONGENITAL ANOMALIES AND PERINATAL RESOURCE UTILIZATION: A POPULATION STUDY

A.M. Moore^{1,2}, W. Yoon³, P.C. Kim⁴, Canadian Neonatal Network

¹Division of Neonatology, Hospital for Sick Children, ²University of Toronto, ³Mount Sinai Hospital, ⁴Surgery, Hospital for Sick Children, Toronto, ON, Canada

Background and aims: Congenital anomalies pose significant health, social and economic challenges to society. The Fetal Alert Network (FAN) integrates antenatal care for pregnancies with anomalies in Ontario (population >13 million) and allows us examine the impact of antenatal diagnosis on maternal and fetal outcomes.

Methods: Retrospective analysis of all infants admitted to Level III NICUs in the Province of Ontario Canada, between Jan 2005 and Dec 2008, using CNN and FAN data. Infants were stratified into anomaly (based on ICD codes), versus non-anomaly groups and pre or postnatal diagnosis of anomaly. Data were analyzed using descriptive statistics and bivariate and multivariate analyses were used to assess fetal and maternal outcomes.

Results: Over the four year period 15,237 infants were admitted to regionalized tertiary level NICUs (2.8% of all births) and 4,400 (29%) had at least 1 congenital anomaly. There was significant difference in maternal age between the 2 groups $p < 0.0001$, but not in maternal diabetes or mode of delivery. There were significantly more male infants in the anomaly group $p < 0.005$. Birth weight (mean 2.3kg) and GA (mean 34 weeks) although low, were greater in the anomaly group. Illness severity as measured by SNAP-II scores, duration of ventilation and length of hospital stay were all greater in the anomaly group, $p < 0.0001$.

Conclusions: This study provides new insights into the impact of congenital anomalies on perinatal health care provision in a regionalised setting and the rationale for continued efforts to decrease the prevalence of these defects.

343

HYPERBILIRUBINEMIA IN A NATIONAL COHORT OF EXTREMELY PREMATURE INFANTS

E. Fladmark¹, G. Haram¹, D. Bratlid^{1,2}, and The Norwegian Extreme Premature Study Group

¹Department of Laboratory Medicine, Children's and Women's Health, Faculty of Medicine,

Norwegian University of Science and Technology,

²Department of Pediatrics, St. Olavs University Hospital, Trondheim, Norway

Background and aims: Hyperbilirubinemia is a possible risk factor for adverse neurodevelopmental outcome in extremely premature infants. The study was undertaken to analyse the frequency, severity and treatment of hyperbilirubinemia in such infants.

Methods: Hyperbilirubinemia was studied retrospectively in a national cohort of 422 infants with gestational age (GA) of 22⁰ to 27⁶ completed weeks and/or birth weight (BW) of 500 to 999 g who survived beyond the 2nd day and had bilirubin measured. Group differences were analysed by t-test.

Results: Mean maximum bilirubin levels (TsB) were 176±43 µmol/L (range 72-391) and did not differ between GA. TsB ≥ 200 µmol/L and ≥ 250 µmol/L was seen in 24.2 % and 4.3 % of infants, respectively. Mean duration of TsB ≥ 200 µmol/L was 1.9±1.7 days (range 1-10) and did not differ between GA, but was more frequent in boys ($p=0.048$). 85.4 % of infants received phototherapy for 2.3±1.7 days (range

0-12). One infant received exchange transfusion, blood group immunization was diagnosed in only one infant (AB0). Prenatal steroids caused lower TsB ($p=0.003$) while initial acidosis increased TsB ($p=0.015$). Increased TsB was also associated with development of persistent ductus arteriosus ($p=0.03$), the degree of intraventricular hemorrhage ($p=0.015$), death ($p=0.025$), and possible neurologic sequelae at discharge ($p=0.02$).

Conclusions: Significant hyperbilirubinemia is common in extremely premature infants in spite of treatment. TsB seems to be associated with both prenatal and neonatal conditions as well as final outcome. Blood group immunization seems to be significantly underdiagnosed in these infants.

344

PRONGS OR MASK FOR NASAL CONTINUOUS POSITIVE AIRWAYS PRESSURE IN PRETERM INFANTS (THE POM TRIAL): A RANDOMISED TRIAL

E.A. Kieran¹, E.J. Molloy^{1,2}, A.R. Twomey¹, J.F. Murphy¹, C.P. O'Donnell^{1,2,3}

¹Neonatology, National Maternity Hospital, Holles St, ²The Children's Research Centre, Our Lady's Children's Hospital, Crumlin, ³School of Medicine and Medical Science, University College Dublin, Dublin, Ireland

Background: Preterm infants frequently receive support with nasal continuous positive airways pressure (NCPAP) via nasal prongs or a nasal mask. While both prongs and masks may cause trauma to the nose, it is not clear which is more effective.

Objective: To compare the effectiveness of NCPAP given via prongs and mask to preterm infants. We hypothesise that the use of prongs compared to masks will reduce the rate of intubation and ventilation within 72 hours of starting NCPAP.

Methods: Infants < 30 weeks gestation receiving NCPAP with the Infant Flow Driver or SiPAP system (Viasys, USA) are randomised to either prongs or mask. Infants are block randomised stratified for gestational age (< 28 weeks, 28-30 weeks); and according to whether NCPAP was started as a primary treatment for respiratory distress or post-extubation. Infants are intubated and ventilated if they reach pre-determined criteria for respiratory failure (2 or more of worsening distress; $FiO_2 > 0.4$; $pH < 7.2$; $PaCO_2 > 9$ kPa; recurrent apnoea) < 72 hours after commencing NCPAP. All other aspects

of treatment are the same between the groups. Relevant secondary outcomes are recorded.

Results: Since enrolment began (02/08/2009) 64 infants have been recruited and had the primary outcome determined. We expect that the primary outcome will be determinable for the total sample (120 infants) by early October 2010.

Conclusion: This randomised controlled trial will provide valuable information about the optimal interface to use when giving NCPAP to preterm infants.

345

COMPARISON OF CEREBRAL TISSUE OXYGENATION INDEX AND CARDIAC OUTPUT IN INFANTS WITH BIRTH WEIGHT LESS THAN 1250 GRAMS

J. Sirc^{1,2}, E.M. Dempsey^{3,4}, J. Miletin^{1,2}

¹Department of Paediatric and Newborn Medicine, Coombe Women and Infants University Hospital, Dublin, Ireland, ²Neonatal Intensive Care Unit, Institute for the Care of Mother and Child, Prague, Czech Republic, ³Department of Neonatology, Cork University Maternity Hospital, ⁴Department of Paediatrics and Child Health, University College Cork, Cork, Ireland

Background and Aims:

(1) To correlate cerebral tissue oxygenation index (cTOI) and cardiac output measurements in infants with birth weight less than 1250 g during first 48 hours of life

(2) To correlate these measurements with clinical parameters

Methods: A prospective observational study. Newborns with birth weight < 1250g were eligible for enrolment. Superior vena cava (SVC) flow, right and left ventricular outputs were measured. Patent ductus arteriosus was assessed and cTOI was measured. All measurements were done at 6, 12, 24 and 48 hours of age. Various clinical parameters (including mean blood pressure, mean airway pressure, pH, pO_2 , pCO_2 , O_2 saturations) were also recorded.

Results: 10 neonates were enrolled following parental consent. The mean birth weight was 767 g ($SD \pm 184$), mean gestational age was 25.3 weeks ($SD \pm 1.4$). Mean SVC flow at 6 hours of age was