59.6 ml/kg/min and increased to 76.2 ml/kg/min at 48 hours of age. Mean cTOI was 64.9% at 6 hours of age, 63.3% at 12 hours of age, 67.5% at 24 hours of age and 67.7% at 48 hours of age. Mean cerebral fractional tissue oxygen extraction (cFTOE) did not change significantly and was 0.28, 0.31, 0.28 and 0.27 at 6, 12, 24 and 48 hours respectively. There was no significant correlation between SVC flow and cTOI.

Conclusion: Superior vena cava flow increased during first 48 hours of life. There was a slight increase in cTOI and no change in cFTOE. This may reflect increased cerebral metabolism during the first 48 hrs.

346

NEONATAL TREATMENT TOXICITY ON DAY 7 AND THROUGHOUT THE FIRST MONTH OF LIFE

R. Vieux^{1,2,3}, J. Fresson⁴, F. Guillemin^{2,3}, J.-M. Hascoet¹

¹Department of Neonatology, Maternité Regionale Universitaire, Nancy Cedex, ²CIC-EC CIE6, INSERM Nancy, ³EA 4360 APEMAC, Nancy University, ⁴Biostatistics Department, Maternité Regionale Universitaire, Nancy, France

Background and aims: Preterm infants are at high risk of drug-induced acute kidney injury. We aimed to determine the impact on GFR during the first month of life, of the drugs prescribed shortly after birth in the Neonatal Intensive Care Units

Methods: multicenter study nested in a cohort of 27-31 weeks GA infants. Infants parted into two groups: "Low GFR": GFR day7 < median GFR for GA; "High GFR": GFR day7 ≥ median GFR for GA. GFR was measured weekly for one month and analyzed with a logistic regression model and an ANOVA with repeated measures (drug and time effects throughout the first month).

Results: 269 infants included. Nephrotoxic drugs prescribed before day7 were mainly aminoglycosides, glycopeptides, and Cox-inhibitors. Cox-inhibitors were far the most nephrotoxic drugs, and GFR remained low for the first month of life after Cox-inhibitor administration. Even high plasmatic levels of aminoglycosides or of glycopeptides did not reach the nephrotoxic effect of Cox-inhibitors:

	"Low GFR" (n=183)	"High GFR" (n=86)	p
Cox inhibitors	55 (30.0%)	15 (17.4%)	0.03
Aminoglycosides >2 days	48 (26.2%)	26 (30.2%)	0.49
High serum level of aminoglycosides	28 (15.3%)	9 (10.5%)	0.28
Glycopeptides >2 days	18 (9.8%)	10 (11.6%)	0.65
Cox inhibitors + aminoglycosides	11 (6.0%)	4 (4.7%)	0.65
Cox inhibitors + glycopeptides	11 (6.0%)	3 (3.5%)	0.38

[Nephrotoxicity and GFR group]

Conclusion: If GFR day7 < median reference value, and infants have received ibuprofen, blood levels of other potentially nephrotoxic drugs should be closely monitored.

347

BRAIN TISSUE OXYGENATION IN VERY LOW BIRTH WEIGHT INFANTS (VLBW) IMMEDIATELY AFTER DELIVERY DURING RESUSCITATION USING SUSTAINED INFLATIONS

H. Fuchs, W. Lindner, A. Leiprecht, T. Trischberger, H. Hummler

Children's Hospital, University of Ulm, Ulm, Germany

Background: Near-Infrared Spectroscopy (NIRS) has been used to study cerebral oxygenation during various conditions in the NICU. However, essentially no information is available on brain tissue oxygenation in VLBWI immediately after delivery.

Objective: To study the timely relationship of brain tissue oxygenation (StO₂) as measured by NIRS in VLBWI immediately after delivery.

Methods: A Foresight laser diode NIRS device was attached to the head of 14 VLBWI (GA 28.2 ± 3.1 wks, BW 907 ± 301 g) immediately after delivery. Preductal arterial oxygenation (SpO₂) was measured using pulse oxymetry and infants were exposed to CPAP 5 cmH₂O, an initial FiO₂ = 0.40 and a sustained inflation (15 s) with a pressure of 20 cmH₂O. Depending on the response of the heart rate and SpO₂ a second inflation with 25 cmH₂O and a third inflation with 30 cmH₂O was allowed. FiO₂ was adjusted according to pulse oximetry