(for both boys and girls 8.1 years, SD 0.15). When compared with the general population, children with DS had more emotional and behavioral problems ($p < 0.001$), except for the anxious/depressed scale on which they scored more favorable ($p < 0.001$). Significantly lower HRQoL scores for the TACQOL scales gross motor skills, autonomy, social functioning and cognitive functioning were found (p -values < 0.001). Hardly any differences were found with regard to the scales physical complaints, positive and negative emotions.

Conclusions: Eight-year-old children with DS have an average developmental delay of four years, more emotional and behavioral problems, and a less favorable HRQoL compared to children from the general population.

446

HIGH FREQUENCY OF LOW SEVERITY DISABILITIES IN VERY PREMATURE INFANTS AT AGE FIVE

E.S. Potharst-Sirag^{1,2,3}, A.G. van Wassenae³,
B.A. Houtzager¹, J.W.P. van Hus⁴, B.F. Last^{2,5}, J.H.
Kok³

¹Pediatric Psychosocial Department, Emma Children's Hospital - Academic Medical Center,
²Developmental Psychology, Free University,
³Neonatology, Emma Children's Hospital - Academic Medical Center, ⁴Rehabilitation, Academic Medical Center, ⁵Pediatric Psychosocial Department, Emma Children's Hospital - Academic Medical Center', Amsterdam, The Netherlands

Background: Children born preterm run an increased risk of problems on multiple domains. The co-existence of these problems needs more attention.

Aim: To find out how many preterm born children have disabilities in comparison with term controls.

Methods: Children (N = 127) born at < 30 weeks GA or BW < 1000 g, born in our hospital, reaching the corrected age of 5 in a 18 month period were eligible. Healthy term born children, mostly recruited via the school of each preterm child, served as controls. Motor development was assessed using the M-ABC II, neurological functioning using the Touwen neurological examination, cognitive functioning (total, verbal and performat IQ, and processing speed quotient) using the WPPSI-III-NL, and behaviour using the SDQ. (parents and teachers). Scores > -1 SD below mean were defined as disability.

Results: 104 preterm born children (mean GA 28.8 weeks (SD 1.6), mean BW 1043 grams (SD 250)) and 91 term controls (mean GA 39.8 weeks (SD 1.8), mean BW 3224 (SD 528) participated. In the preterm group, 50% had one or more abnormal IQ score and 44% had minor neurological dysfunction, while 5% had CP. In the term controls this was 22%, 7% and 0% respectively. In the preterm group, 75% had ≥ 1 disability and 52% ≥ 2 disability and this was 27 resp. 10% in the term controls.

Conclusions: Of very premature children, able to undergo an age appropriate IQ test, only 25 % compared to 73% of term controls is without disability on all domains.

447

PARENTAL EDUCATION DOES NOT INFLUENCE OUTCOME OF VERY PRETERM CHILDREN ON ALL DEVELOPMENTAL DOMAINS

E.S. Potharst-Sirag^{1,2,3}, A. van Wassenae²,
B.A. Houtzager^{1,2}, J.W.P. van Hus⁴, B.F. Last^{1,3},
J.H. Kok²

¹Pediatric Psychosocial Department, ²Neonatology, Emma Children's Hospital - Academic Medical Center, ³Developmental Psychology, Free University, ⁴Rehabilitation, Academic Medical Center, Amsterdam, The Netherlands

Background: Children born preterm run an increased risk of problems on multiple domains. It is not clear how SES influences these different outcomes.

Aims: To study the relation between parental education and multiple domain outcomes at age five in very preterm born children in comparison with term controls.

Methods: 104 children born < 30 weeks infants GA and/or < 1000 gr at birth and 95 term controls were assessed at the (corrected) age of five years. Motor, neurological cognitive (total, verbal and performat IQ, and processing speed quotient) and behavioural functioning were assessed using the M-ABC II, the Touwen examination, the WPPSI-III-NL, and the SDQ. (parents and teachers). The differences between preterm and term children were studied in three strata of parental education level (high, medium, low).

Results: The mean difference in IQ score between 104 preterm and 95 term children was 10 points.

In the high education stratum this difference was 5 compared to 14 in the low stratum. The same pattern was found for verbal, performal IQ, processing speed and SDQ scores. Neurological outcome of preterm born children was not normal in 49% especially due to minor neurological dysfunction (MND), which occurred in 44%. MND nor M ABC scores were influenced by parental education, with comparable preterm-term differences in each parental education stratum.

Conclusions: The biological risk of preterm children is highlighted by their neurological outcome. The social risk factor modifies this biological risk and determines especially cognitive and behavioral outcome.

448

OBSTRUCTIVE PULMONARY DISEASE IN OLD AGE AMONG INDIVIDUALS BORN PRETERM

E. Berggren Broström

*Karolinska University Hospital Neonatology,
Karolinska Institutet, Stockholm, Sweden*

Background: There are only few studies of the association between preterm birth and risk of chronic lung disease in old age. The aim of this study was to assess the association between poor fetal growth, preterm birth, sex and risk of asthma and Chronic Obstructive Pulmonary Disorder (COPD) in adulthood.

Method: We have followed up a cohort of all infants born preterm (< 35 weeks) or with low birth weight (< 2,000 grams and < 2,100 grams for girls and boys, respectively) in a source population of 250,000 individuals born from 1925 through 1949 in Sweden and an equal number of controls. Follow-up started in January 1st, 1987 and continued to December 31, 2006. At start of follow up, there were 6,425 subjects in the cohort. There were 2,176 subjects in the cohort with low birth weight (< 2,500 grams), and 151 had a birth weight of less than 1,500 grams. In all, 2,931 subjects were born preterm (< 37 weeks), and 986 had a gestational duration of 32 weeks or less. We considered cohort subjects as cases if they had a main discharge diagnosis of asthma or COPD.

Results: For any obstructive airways disease, there was a statistically significant increase in risk with decreasing birth weight when males and females were analyzed together.

Compared to women born at term, women born before 33 weeks of gestation had an almost threefold increase in risk for any obstructive airways disease. For fetal growth, the association was u-shaped rather than linear.

449

LOW MATERNAL LEVELS OF FIRST-TRIMESTER PAPP-A AND FREE β -HCG ARE ASSOCIATED WITH AN INCREASED RISK OF ADVERSE NEONATAL OUTCOME

I. Kirkegaard^{1,2}, N. Uldbjerg², T.B. Henriksen^{1,3}

¹Perinatal Epidemiology Research Unit, ²Dept. Obstetrics and Gynecology, ³Dept. Pediatrics, Aarhus University Hospital, Skejby, Aarhus, Denmark

Background and aim: Pregnancy-associated plasma protein-A (PAPP-A) and free beta-human chorionic gonadotropin (β -hCG) are highly efficient in first-trimester screening for Down syndrome. In addition, studies have found them associated to preterm delivery, pre-eclampsia and low birth weight, but the association to neonatal outcomes is only sparsely elucidated. The aim of the current study was evaluate PAPP-A and free β -hCG in relation to adverse neonatal outcomes.

Methods: 9450 singleton pregnant women at Aarhus University Hospital were included between January 2005 and December 2007. PAPP-A and free β -hCG were measured in the first trimester. Neonatal outcomes were obtained from specific registration forms filled out by the doctors responsible for admission and discharge from the neonatal intensive care unit (NICU). The outcomes explored in this study were admission to NICU, neonatal jaundice and hypoglycemia.

Results: PAPP-A below 0.4 MoM was significantly associated with admission to NICU (OR 1.6; 95% CI, 1.2 - 2.0), and the same applied to free β -hCG below 0.4 MoM (OR 1.5; 95% CI, 1.1 - 2.1). The results could not be explained by preterm delivery or low birth weight as the results remained unchanged when restricting to deliveries in gestational weeks 39 and 40 and when adjusting for birth weight. Furthermore, we found significant associations between PAPP-A and both hypoglycemia and jaundice and between free β -hCG and jaundice.

Conclusion: Both low PAPP-A and free β -hCG were associated with an increased risk of admission to NICU. The results could not be explained by either preterm delivery or low birth weight.