Method: Fifty-two children and adolescents (26 females) aged 9.0-18.0 years (median: 14.0 years) completed validated measures of QoL, locus of control and self-concept 12-192 months after heart transplantation for cardiomyopathy (n=38) or congenital heart disease (n=14).

Results: QoL scores for the total transplant group were lower than those of healthy norms ($t=3.45$, $p=.001$) but those transplanted for cardiomyopathy did not differ from the healthy population and had better perceived QoL than those with a pre-transplant diagnosis of congenital heart disease ($p=.003$). There were no differences in self-concept or locus of control between these two groups but transplanted children had a more positive self-concept than healthy children ($t=2.079$; $p=.043$). Higher QoL scores were correlated with positive self-concept ($r=.583$; $p<.001$), an internal locus of control ($r=.560$; $p=.003$) and older age ($r=.506$; $p<.001$) but there was no relationship between time since transplant and QoL.

Conclusion: Many children and adolescents have an excellent QoL after transplant, particularly those transplanted for cardiomyopathy. The relationship between QoL, self-concept and locus of control requires further investigation longitudinally to determine causality and to inform interventions to maximise QoL for all patients.

527

ON YOUR OWN FEET AHEAD: INNOVATIONS IN DUTCH TRANSITIONAL CARE FOR ADOLESCENTS WITH CHRONIC CONDITIONS

A. van Staa^{1,2}, J. Havers³, H. Sonneveld², On Your Own Feet

¹Expertise Centre Transitions of Care, Rotterdam University, ²Institute Health Policy & Management, Erasmus University Rotterdam, Rotterdam,

³Dutch Institute for Healthcare Improvement CBO, Utrecht, The Netherlands

Background: In the Netherlands, adolescents with chronic conditions are often ill prepared for their transition to adult care. Research revealed that services are not geared toward adolescents' needs and preferences. To bridge these gaps, a four-year innovation programme, On Your Own Feet Ahead, was started in 2008.

Aims:

1) To develop and execute a national programme aimed at providing more integrative health care services and appropriate transitional care to adolescents (12 -25 years) with somatic chronic conditions in the Netherlands.

2) To monitor the changes by an independent evaluation study.

Methods: Using the Breakthrough method, multidisciplinary pilot teams from hospitals and rehabilitation centres in the Netherlands implement a number of interventions and policy changes to improve transitional care. Effects are evaluated on team, parent and patient level.

Results: The programme focuses on the establishment of independent health behaviours, increase self-management skills and improvement of the organisation of transitional care. Interventions include developing a joint policy and protocol for care transfers, appointing a transition coordinator and setting up joint clinics between paediatric and adult care. To enhance self-management and self-reliance, most teams have started implementing the newly developed Individual Transition Plan (ITP).

Conclusions: The evaluation study demonstrated that after a year most teams reported that the programme has helped them to improve on the major goals they had set to achieve. They are making progress in making care more patient-centred. A Toolkit of suitable interventions was developed and the programme will be disseminated nation-wide in the coming years.

528

TRANSITION FROM CHILD TO ADULT CARE FOR YOUNG PEOPLE WITH MENTAL HEALTH PROBLEMS: A PROCESS OF TRANSITION

S. Buckley

Nursing & Midwifery, Trinity College Dublin, Dublin, Ireland

The transition of young people from child to adult healthcare services has become increasingly important. Evidence suggests that the process is often poorly managed and many young people with mental health problems experience abrupt ad hoc transitions from child to adult healthcare services. A well planned transition is viewed as leading to improved clinical, educational and social outcomes

for young people. Whilst a poorly planned transition is associated with increased risk of non-adherence to treatment and decrease of follow up which can lead to serious consequences for the young persons' health and well-being. There is limited empirical evidence in Ireland on how the process is experienced by young people with mental health problems; how it impacts on their health and well-being and how the transition process is managed by all stakeholders.

This study proposes a mixed methods longitudinal design to explore the views of young people, their parents and healthcare professionals (Child Psychiatrist, Registrars, Nurses, Psychologist and Social Workers) on the experiences of transition. The data will be obtained using questionnaires and interviews at five different time periods:

- (1) in the child mental health services
- (2) interviews with participants before transition (in child services)
- (3) distribution of questionnaire to young people-after transition
- (4 & 5) interviews with participants after transition (adult services) Placing the young persons experience as central to the development of services and ensuring continuity of care are key priorities of National Children's Strategy (2000) and directly impact the quality of service provision.

529

**AUDIT ON SWINE FLU VACCINATION
UPTAKE IN STAFF WORKING IN PEDIATRIC
DEPARTMENT AT IPSWICH GENERAL
HOSPITAL**

M. Kumar, D. Agarwal
NHS Suffolk, Ipswich, UK

Background: H1N1 influenza pandemic management was the most challenging scenario faced by health authorities worldwide recently. Department of health (DOH) in UK after doing epidemiological study of first wave of swine flu (April to Sep 2009) gave recommendations for management of predicted Second wave. One of the important recommendations was to prioritize vaccination of frontline health and social care workers. Previous studies have shown that the uptake of seasonal flu vaccine among healthcare workers has consistently been low.

Objective: To evaluate the uptake rate and reasons for non-uptake of H1N1 Influenza vaccine in frontline healthcare staff working in a busy Pediatric and Neonatal Unit in a Hospital in East Anglia, U.K.

Method: Prospective study conducted over 4 week period during Second Wave of Swine Flu Pandemic. A simple anonymised questionnaire was equally distributed throughout various areas of pediatric and Neonatal department. Staff was randomly informed verbally and actively encouraged to ensure maximum responses.

Results: 46(62%) out of 74 respondents were not vaccinated. 30 (65%) out of 46 staff were not vaccinated because of Lack of information either relating to vaccine itself (safety, efficacy, side effects); its availability and accessibility .16 (35%) out of 46 said they were "too busy to get vaccinated". Other reasons stated were - unpleasant reaction to seasonal flu vaccine, needle phobia and people being on annual leave.

Conclusion: Despite the pandemic nature of disease and robust mechanisms in place, Swine flu vaccination uptake continued to be poor. Onsite departmental information and vaccination sessions can be helpful to improve uptake.

530

**PERSISTENCE OF IMMUNE RESPONSE TO
CANDIDATE MENINGOCOCCAL SEROGROUPS
A,C,W-135,Y TETANUS TOXOID-CONJUGATED
VACCINE (MENACWY-TT) UP TO 42 MONTHS
FOLLOWING PRIMARY VACCINATION**

**L. Østergaard¹, V. Bianco², M. Van der Wielen²,
J. Miller³**

¹Department of Infectious Diseases, Skejby
Sygehus, Aarhus University Hospital, Aarhus,
Denmark, ²GSK Biologicals, Wavre, Belgium,
³GSK Biologicals, King of Prussia, PA, USA

Background and aims: A primary dose of the candidate meningococcal A,C,W-135,Y tetanus toxoid-conjugated vaccine (MenACWY-TT) was immunogenic and well tolerated in subjects aged 15-19 years (NCT00126945). Here, the persistence of the immune response to MenACWY-TT was compared with the MenA,C,W-135,Y polysaccharide vaccine *Mencevax*[®] ACWY (MenPS), and was evaluated up to 42 months after primary vaccination.

Methods: This series of phase II, open, controlled persistence studies included subjects who received