limited to hypothermia and treatment of seizures. The principal function of the GABA system in mature brain is inhibition, however in neonatal brain GABA provides much of the excitatory drive. Thus whilst anticonvulsants augment GABA's inhibitory actions in mature brain, administration of GABAergic drugs to neonates may potentially exacerbate seizures and worsen HI brain injury. Furthermore changes in GABA_A receptor expression will influence receptor pharmacology. We aimed to assess changes in protein expression of the GABA_A receptor in the neonatal HI piglet.

Methods: Newborn piglets were subjected to a 30 min HI insult and euthanased at 72 h. Brain tissue was collected from several brain regions and GABA_A receptor α_3 protein expression levels analysed by western blot.

Results: Expression of α_3 was significantly elevated in frontal cortex of HI animals without seizure; animals with seizure trended toward lower α_3 expression. In occipital cortex, α_3 expression was found to be diminished in all HI animals however presence of seizures was associated with significantly lower α_3 expression.

Conclusions: GABA_A receptor α_3 expression was significantly altered following neonatal HI; presence of seizures further changed this expression. Efficacy of anticonvulsants in neonatal brain may not only depend on regional and temporal maturation of GABAergic inhibitory function but also on GABA_A receptor subunit expression following both HI and seizures. There is a critical need to develop effective treatment strategies specific to the neonatal HI brain.

572

ROUTINE MAGNETIC RESONANCE IMAGING OF THE BRAIN AT TERM CORRECTED AGE IN ALL OUR VERY LOW BIRTH WEIGHT INFANTS

B. Geier¹, S. Otto², N. Utzig³, R. Stenger¹, M. Zygmunt⁴, C. Fusch⁵, N. Hosten²

¹Neonatology, ²Neuroradiology, ³Paediatric Neurology, ⁴Gynaecology, Universitätsklinikum Greifswald, Greifswald, Germany, ⁵Neonatology, McMaster University, Hamilton, ON, Canada

Background and aims: To compare findings of cranial ultrasound (US) and magnetic resonance imaging (MRI) of the neonatal brain.

Methods: A retrospective population based cohort study of premature infants with a very low birth weight (VLBW) of less than 1500 grams who were admitted to our neonatal unit on day 1 (period: 2007 to 2009).

All infants underwent cranial US on day 1, 3, 7, 14, 28, 42 monthly thereafter and at term corrected age. Infants who were eligible for this study obtained a MRI at term (37 to 42 weeks gestational age) corrected age.

Retrospectively, we compared the cranial US findings as a predictor of a wide spectrum of pathology on MRI.

Results: Paired MRI and US studies were performed in (n=140) VLBW infants who were born at a median gestational age of 28 (range: 22+1 to 34+5) weeks and a median birth weight of 1020 (range: 335 to 1495) grams.

US predicted some MRI findings accurately: germinal layer haemorrhage (GLH), cystic lesions, intraventricular haemorrhage (IVH) and severe white matter (WM) echogenicity on US for the presence of WM haemorrhagic parenchymal infarction on MRI.

Other MRI changes were less well-predicted: delay in maturation and myelination, reduced cortical folding, congenital malformations, punctate lesions and mild or no WM echogenicity on US for the presence of normal (n=112) WM signal intensity on MRI.

Conclusions: MRI of the neonatal brain might shed light on the origin of brain lesions causing long-term neurodevelopmental sequelae and might change our perinatal and neonatal management.

573

NEUROMUSCULAR ACTIVATION DURING REVERSE AND NON-REVERSE CHEWING CYCLES IN CHILDREN WITH UNILATERAL POSTERIOR CROSSBITE

M.G. Piancino¹, D. Farina², F. Talpone¹, A. Merlo¹, T. Valledonga¹, G. Frongia¹, P. Bracco¹

¹Department of Orthodontics and Gnathology, University of Turin, Turin, Italy, ²Department of Health Science and Technology, Center for Sensory-Motor Interaction (SMI), Aalborg University, Aalborg, Denmark

Introduction: Posterior unilateral crossbite is an asymmetric malocclusion developing early, during