and December 2009 with the diagnosis of severe TBI. KOSCHI was used to assess recovery.

Results: Of 92 admissions for trauma, 47 had severe TBI. The median age was 8 years, PRISM score was 13 and PTS was 6. The leading cause was road traffic accident (57%). Specialized medical teams provided pre-hospital care in 68% of cases. Raised intracranial pressure (ICP) signs were present in 34 cases. Neurosurgical intervention was necessary in 16 children and 10 had invasive ICP monitoring. All children were ventilated, 42 had hyperosmolar therapy (mannitol or NaCl 3%), 5 barbiturate-induced coma and 12 vasoactive drugs. Transcranial doppler (TCD) was performed in 27 children, 18 had raised pulsatility index. There was a good correlation between TCD information and ICP readings. Eight children (17%) died. In the follow-up period 8 children were in KOSCHI category 1, 2 in category 2, 5 in category 3, 11 in category 4 and 21 in category 5.

Conclusions: KOSCHI provides a practical scale for paediatric head injury outcome. A high percentage of patients were discharged without any sequelae and mortality rate was low. Treatment in a PICU and early neurosurgical consultation were determinant for the good outcome.

576

RECOVERY OF SLEEP-WAKE CYCLING ON AMPLITUDE-INTEGRATED ELECTROENCEPHALOGRAPHY IN POST-SURGICAL TERM NEONATES

M. Olischar¹, A. Davidson², R.W. Hunt³

¹Neonatology, University Children's Hospital Vienna, Vienna, Austria, ²Anaesthesia and Pain Management, ³Neonatology, The Royal Children's Hospital, Melbourne, VIC, Australia

Background: Sleep characteristics have been used for prediction of neurodevelopmental outcome and may also be useful to determine quality of recovery after surgery. Cerebral monitoring has not previously been routinely used to observe effects of anaesthesia and surgery on the developing brain.

The aim of this prospective study was to evaluate the influence of anesthesia on the recovery of SWC in term newborns after surgery.

Methods: Term neonates admitted to NICU who underwent major intra-abdominal surgery were studied. The Brainz® monitor was applied post-

operatively using the standard C3, C4 and P3, P4 positions. Subjects were monitored until mature SWC were observed and the time of onset and quality of SWC together with the background pattern were then assessed.

Results: Forty-seven patients were recruited. Surgery was performed for congenital diaphragmatic hernia (15), esophageal atresia (12), duodenal and jejunal atresia (7), ileal obstruction (3), exomphalos (3), gastroschisis (4), malrotation (1), meconium ileus (1) and a patent urachus (1). When patients returned to the NICU morphine and midazolam infusions were commenced. Emergence of SWC was observed at a median of 13 hours after surgery (range 1 to 136 hours). SWC became mature at a median of 30 hours (range 4 to 175 hours). All patients were still on morphine and midazolam infusions at this stage.

Conclusion: The emergence of sleep-wake cycling on aEEG after surgery can be observed as early as one hour post-operatively and appears not to be affected by high dose infusions of morphine and midazolam.

577

THE INFLUENCE OF MUSIC ON AEEG-ACTIVITY IN NEUROLOGICALLY HEALTHY NEWBORNS ≥ 32 WEEKS GESTATIONAL AGE - A PILOT STUDY

M. Olischar¹, H. Shoemark², T. Holton², R.W. Hunt²

¹Neonatology, University Children's Hospital Vienna, Vienna, Austria, ²Neonatology, The Royal Children's Hospital, Melbourne, VIC, Australia

Background: Music is a valuable resource in the NICU. In pursuit of further evidence aEEG may provide a clear picture of the impact of recorded music on newborn brain activity.

Aim: To evaluate the influence of music on aEEG-activity in neurologically healthy neonates born ≥ 32 weeks' gestational age.

Methods: Twenty neurologically healthy infants admitted to the NICU at The Royal Children's Hospital, Melbourne with minor abnormalities requiring further investigation were included in this study. Ten subjects were randomized in the assessment group (music stimulation) and 10 subjects were only monitored by aEEG (without music stimulation). The source of music was Music

for Dreaming, (Sound Impressions, Pty. Ltd.) and was administered using micro-speakers 20cm from the infant's head at 50-55 dbA. The aEEG was recorded on the BrainZ Monitor (BRM2, Version 8.0, BrainZ Instruments, New Zealand). Traces were evaluated offline for sleep-wake cycles and background patterns.

Results: aEEGs were recorded for a mean duration of 4.0 hours. All 20 patients showed continuous background patterns. SWC in the intervention group had a mean duration of 19 minutes, minimum amplitudes of 6mcV and maximum amplitudes of 25mcV whereas SWC in the control group had a mean length of 19min, min. ampl. of 8mcV and max. ampl. of 25mcV. The duration between SWC was 50 minutes in the intervention group and 53 minutes in controls.

Conclusion: There is a trend in SWC becoming more discontinuous in subjects exposed to music. There seems to be an effect of music on the behaviour of SWC.

578

SOME OCCASIONAL INTERVENTIONAL DIFFICULTIES IN CHILDHOOD-DIABETIC-KETOACIDOTIC(DKA)-HYPERGLYCEMIC-HYPEROSMOLAR(HHS) PRECOMA-COMAS STATES WITH UNUSUAL FEATURES/ ASSOCIATIONS: THE ARGUMENT FOR A COMPREHENSIVE-SYSTEMATIC-EVALUATION OF CASES FOR A DIRECTED- TIMELY- APPOSITE-INTERVENTIONS

E.U. Onyekwelu, Neurology

Paediatrics, Royal Victoria Teaching Hospital, Banjul, The Gambia

Background/purpose: The 4Ps-polydipsia-polyuria-polyphagia-progressive weight loss/ associated complications heralds the onset-of-adult-Type-I-DM. However, in childhood most cases were brought to notice by the abrupt onset of DKA-HHS-precoma/comas, occasionally other-unusual-complications/strikingly uncommon-associations, overwhelms-the-classic-features-/ posing-diagnostic/management -enigmas/delays. Some children with Diabetic coma continue to have physiologic-metabolic-instability/deepening-coma despite-aggressive-interventions.Although the most likely causes may be DKA-cerebral oedema-HHS, -other unusual -aetiology/pathological-associations-such as hypercoagulability,

-erythrocytopaenia, -meningoencephalitis, -malaria-, intracerebral suppuration,-acute-leucocytoclastic-vasculitis,-thrombocytosis,-reduced red-cell-plasticity,other-causes of metabolic-acidosis,-ATHI,-MRDM,-tumefactions should be considered/-excluded-especially in those cases-presentinig-unusually/-atypically-with- rare/-remarkable associations or/were therapeutically-difficult.

Methods/cases: An observational-prospective-analysis-of-cases-of-relevance.

Results: 66-cases of childhood-DM,with-precoma/-coma needed admissions,/closer monitoring ,/Intensive-care-support.Typically-classical-features-were-observed in 43(65.15%), while-the-remaining 23(34.85%) presented-atypically-unusually or with-remarkably-rare associations.Manifestations-ranging-from-extreme-fatigue,-exhaustion,-kaussmauls-breathing,-air hunger,-staggering,-unsteady/ abnormal-gait,-drowsiness,-somnolence,-stupor,-delirium,-confusion,acute-cognitive-defects,-disorientation,-altered mental status,-slurred speech,-obtundation,-poorly-responsive-status,-focal-neurologic-deficit,-sudden uncounciouness,-frank coma- were remarkably accompanying features in both subsets.The unusual,/atypical presentations/rare associations were subclassified systemically:These were DKA-HHS -with neurological-complications:Dural-sinus/-Lateral-sellar-compartment-thrombosis(n=1),Jacksonsonian-like-seizures(n=1)-encephalitis(n=1),-Severe-malaria/hyperparasitaemia(n=1),

D K A - w i t h - c i r c u l a t o r y - complications:Dysautonomia/-Internal warmth-syncope(n=1)-erythrocytopaenia(n=2),-acute-gastroenteritis(n=1),Cardiogenic-Hypovolumeic-shock-syndrome(n=1),necrobiosis-lipoidica-diabeticorum.(n=1),perforating-granuloma-annulare(n=1)-

DKA-with-renalcomplications:Azotaemia(n=1)-(enuresis n=1),-Haematuria-Urethral-syndrome(n=2),-hypocalcaemia(n=1)-Lactic-acidosis(n=1)-Deranged-LFTS(n=2),

DKA-with-multiple-complications:Multisystemic-infections-suggestive of an immunosuppressive-state such as suppurative-pneumonias-UTIs-GETs(n=1). DKA with other specified associations- Mariucs syndrome(n=1),MRDM-HHS(n=1),MRDM-DKA(n=1), Cases had very-