649

PARATHYROID HORMONE (PTH) AND MARKERS OF CALCIUM METABOLISM IN CHILDREN WITH CONGESTIVE HEART FAILURE

A.A. Fattouh¹, **R.N. Said**², A.M. Nour El Din¹,
R.I. Ammar², D.A. Ah³

¹Pediatrics, National Research Center, ²Pediatrics, Faculty of Medicine, Cairo University, ³Clinical and Chemical Pathology, National Research Center, Cairo, Egypt

Background: Parathyroid hormone (PTH) regulates the content of calcium and exerts an effect on myocardial function. Abnormal secretion of PTH has been reported with depressed mechanical performance of the heart muscle.

Aim is to evaluate parathyroid function in patients with CHF due to dilated cardiomyopathy (DCM), measuring some different parameters of calcium metabolism.

Patients and Methods: A case-control study carried on 30 cases (20 males and 10 females) with congestive heart failure (CHF) having a mean age of 6.69 ± 3.94 years. They were recruited from Cardiomyopathy Clinic in Children Hospital Cairo University. Cases were further classified into (i) Decompensated HF (n=18) (NYHA III, IV) (ii) Compensated HF (n=12) (NYHA II). Thirty healthy children (matched age and sex) were enrolled to serve as control. Parathyroid hormone (PTH) and Osteocalcin (OC) were measured, in addition to total serum calcium, phosphorus, alkaline phosphatase, sodium and potassium.

Results: There is statistically significant lower serum calcium, osteocalcin and higher serum phosphorus levels in DCM cases compared to controls ($P=0.0001$, 0.0001 , 0.002 respectively). There was no statistically significant difference between cases and controls regarding serum PTH. i.e. normal PTH despite depressed calcium in DCM

Conclusion: Although three decompensated CHF cases showed elevated Parathyroid hormone levels, (NS) PTH levels were normal in rest of cases. Significant hypocalcemia, hyperphosphatemia and reduction in serum Osteocalcin levels were noted in CHF patients. Whether, or not, these factors contribute to systemic illness that accompanies CHF, including bone wasting, mandates further probing

650

INVESTIGATION OF THE PROBABLE CAUSES OF SPECIFIC CHILDHOOD DISABILITIES IN EASTERN AFGHANISTAN (PRELIMINARY REPORT)

J.A. Nasir

Environmental Health Protection Unit, Public Health Institution of Canton Sarajevo-Pediatrics Department Division, National Institute of Health Islamabad, Islamabad, Pakistan

Summary: To identify risk factors likely to cause developmental disabilities and to generate hypotheses for a further study of risk factors predisposing to disability and slow mental and physical development. Mothers and children attending/visiting the Sandy Gall's Afghanistan Appeal (SGAA) clinics in the East Zone of Afghanistan. The main outcome measures, developmental disabilities, cerebral palsy (CP), club foot, CDH and polio. Results, 37.5% of the disabilities were present at birth, 46% of the disabled children were born from parents who are 1st cousin which was confirmed by computing χ^2 value, which is 10.87 with one degree of freedom. 58.3% of the disabled children were born from parents who lack antenatal care. 22.4% of the pregnant women in the defined population had antenatal checkups, and 97% of the mothers in the defined population are illiterate, 52% of the children found with CP specific and mental retardation. 25% were with delayed physical and mental development. 13% were with CP specific and 10% were club foot. 2% of the children were visually handicapped, 9% with hearing impairment. 61% of children were from Nangarhar, 25% from Laghman and 14% from Kunar.

651

THE SPECTRUM OF COMPATIBLE LANDRY-GUILLAIN-BARRE SYNDROME IN CHILDHOOD. A SINGLE REFERRAL CENTRES EXPERIENCE: AN ANALYSIS OF 48 CASES

E.U. Onyekwelu, Neurology

Paediatrics, Royal Victoria Teaching Hospital, Banjul, The Gambia

Background / purpose: Although the association of neuropathies/infections was suggested historically, the first scholarly description was made by Landry in 1859.

In 1892, Osler classically described acute febrile polyneuritis.

Guillain/Barre/Strohl in 1916 characterized Albumino/cytologic/dissociation(ACD)

In 1956, Miller/Fisher(MF) described an ataxic/pseudo/internuclear/opthalmoplegic[PINO]clinico/pathologic/immunologic variant.

LGBS is a post-infective/immune-related/symmetrical/motor,/occasionally sensory/areflexic/flaccid/ascending/demyelinating/polyneuropathy. Clinico/pathologic/ geographic/molecular/epidemiologic distinct variants were described. Presumptively diagnosed, in appropriate clinical settings, by ACD/F-wave/conduction/latency /immunological/ MRI perturbations. However its chronology/ temporality/ clinico/patho/radiological/ patterns may be occasionally non-specific/characterless. Relative to inflammatory-demyelinopathies, axonopathies implicated longer coursed nerves most. Structural/functional recoveries occurred most rapidly in inflammatory-demyelinopathies. In otherwise previously healthy children, a rich anastomic/microvasculatic/meshworked vaso-nervorum made immuno/vasculitic/end arteritic obliterative/ischaemic/polyneuropathies unlikely epi-phenomena. Previously, most investigations targeted adults.

This paper addressed a single referral centres experience with childhood/

LGBS to illuminate these aspects.

Methods: LGBS cases were analyzed for historical/clinical/pathological /radiological details.

Results: 48 cases were concerned (n=31) presented in dry season/ M: F= 5:7/ mean age = 7.8 years. In (n=35) an associated infective antecedent 2-4 weeks previously, ARIs(n=25)/acute gastroenteritis(n=9). Predominant features were acute progressive ambulatory difficulties /cephalgia/myalgia/sensory defects/flaccidity/ areflexia/hyperhydrosis/PINO.

Objective-subjective /laboratory/ functionality recovery phenomenologies inferred AIDP(n=17)/AMAN(n=16)/AMSAN(n=9)PC (n=6) / M-F(n=1) DN (n=3) respectively.

On ambulatory follow ups, suboptimal /tardively resolving cases were asymptomatic. Gastroenteritic precedents were commonly associated with atypicalities.

Antimicrobial applications for co-existent RTIs/ UTIs/ physico-therapy, carbamazepine for neuro-myalgias expedited recovery/ precluded invasive investigations/interventions

/ventilatory support/opioids/plasmapheresis/immunoglobulin expedited timely discharge.

Conclusions/importance: Experientially, LGBS presented tardively.

Its putative antecedent/epidemiology / features/ periodicity/ laboratory profile could suggest the likely neuropathologic subsets in an individual case. The poliovirus must not be considered an antecedent in LGBS.

652

THE ANTECEDENTS, NATURAL HISTORY AND INTERVENTIONS OF BRACHIAL PLEXOPATHIES (BPS); WITH IMPLICATIONS FOR PATTERNS OF ACCIDENTAL AND NON-ACCIDENTAL INJURIES

E.U. Onyekwelu, Neurology

Paediatrics, Royal Victoria Teaching Hospital, Banjul, The Gambia

Background / purpose: BPs are defects in the anterior rami of C5 to C6 or C7 -Erbs-Duchenne-(ED), C8 to Th1-Dejerine-Klumpkes-(DK) spinal nerves which impulsates the upper limbs musculatures due to perilesional oedema, haemorrhages, neuro-axonotemesis, contusions or Wallerian degenerations. Occasionally BPs-acute traumatic injury (ATI) are radiologically indistinct epi-phenomena in infancy. Some of its aspects were either unaddressed/uncharacterized/contentiously discussed or not uniformly adaptable.

Method: A prospective observational study of cases in a single referral centre.

Results: Of the 64 cases, (n=48) were perinatally related.

Appropriately supervised deliveries(n=17)/delayed supervision(n=19)/absolutely

unsupervised(n=12)Localizations: right (n=38). Putatively principal, antecedents'

were macrosomia-dystocia/assisted-breech .Limited range of motion,(ROM)/neuro-