the neuroinflammatory component of autism. Neuroimaging of intranetworking connections has rapidly gained momentum within the field and an understanding of not only how they function as a unit but also how the innate anatomy can endure stress could play a vital role in identifying a therapeutic agent for the condition.

671

BEHAVIOURAL, STRUCTURAL AND CHEMICAL CHANGES SURROUNDING BRAIN INFLAMMATION; THE COMMON PATHWAY UNDERLYING AUTISM

D. Rasoul¹, E.C. Campbell², S. Lynch², G.H. Doherty², M.H. Dunn³, S.J. Powis², A.M.H. Young⁴

¹*School of Medicine, University of Manchester, Manchester*, ²*Bute Medical School*, ³*Department of Physics and Astronomy, University of St Andrews, St Andrews*, ⁴*School of Clinical Medicine, University of Cambridge, Cambridge, UK*

Introduction: A causation model for Autistic Spectrum Disorder (ASD) has not yet been firmly established. Characterised by a triad of impairments in social skills, verbal communication and behaviour, the condition has been recognised as having a strong inflammatory component. Historically, the functional anatomy of the prefrontal cortex is well defined and its dysfunction in autism spectrum disorders has lead to the proposal of a number of psychological models identifying the impact of under development within the condition. However, only recently have we begun to understand the structural and molecular implications underpinning the disease.

Methods: Using ICD-10 and DSM-IV classifications we review the diagnostic characteristics which compound the condition together with the current thinking on the roles of cytokines such as IL-1 α , IL-1 β , IL-6 and TNF- α , their effect on intracellular signaling cascades and ultimately their effect on cell ultrastructure.

Results: Here we collaborate the current understanding of a novel autistic triad consisting of the behavioural, chemical and ultrastructural balances in the autistic brain. We demonstrate how they interact, concluding that these details are neither mutually exclusive nor mutually independent.

Conclusions: This model for autistic spectrum disorder demonstrates the requirement for a multidisciplinary approach to the treatment of

the condition and hope that the integration of this hypothesis will assist in the development of both diagnostic and therapeutic advances.

672

AN AUDIT OF THE MANAGEMENT OF A FIRST AFEBRILE SEIZURE IN CHILDHOOD

M. Boyle, J. Chukwu, A. Connolly, D. Webb

National Children's Hospital, Dublin, Ireland

Objective: To compare the management of first afebrile seizures with internationally recognized standards in an Irish tertiary paediatric setting.

Methods: Records of first afebrile seizure presenting to a Paediatric Emergency Department period between June 2007 and December 2008 were identified through interrogation of emergency department admission books and Paediatric ward ledgers. Management standards were devised from a compilation of British (NICE 2004) and North American (AAN 2003) guidelines. The standards comprised 3 principle areas: (i) the medical record and seizure documentation (ii) investigations recorded and (iii) documentation of management at discharge. Complete concordance with the standards would accrue a score of 21/21. Relevant patient charts were reviewed, assessing medical and nursing notes.

Summary: 26 children were identified over the 18 month period, 13 boys and 13 girls (age range from 1 month to 13 years). No case achieved full concordance with the devised standards. The average score was (13) 50% with a range from 5 (19%) to 17 (65%). Seizures were well documented 75% but compliance with guidelines on investigation of the first afebrile seizure in childhood was poor 63%.

Conclusion: Presentation with first afebrile seizures is common in childhood. We have demonstrated a lack of concordance with international guidelines on management of the first afebrile seizure in our institution. We now intend to provide further guidance on this topic and introduce an information pack at discharge before repeating the audit cycle.