47 (19%) of those reviewed had a final diagnosis of musculoskeletal chest pain, 41 (17%) had a diagnosis of normal child, 22 (9%) had a Lower Respiratory Tract Infection, 18(7.5%) had Gastro oesophageal reflux.

No child required immediate cardiology review.

Conclusions: Chest Pain is a common presentation to the ED, our series supports evidence that it is rarely cardiac in origin.

723

BELL'S PALSY: TO TREAT OR NOT TO TREAT

R. Mc Namara¹, P. Keenan¹, M. Mc Kay²

¹Paediatrics, ²Emergency Medicine, Children's University Hospital, Dublin, Ireland

Objective: To audit management of children presenting with Bell's Palsy to a paediatric Emergency Department (ED) in a tertiary referral centre.

Methods: Retrospective chart review of children presenting with Bell's palsy from October 2006 to February 2009. Information was gathered on investigations carried out, treatment prescribed, referral to neurology and follow up.

Results: In total 46 children presented with Bell's palsy from Oct 06 to Feb 09, F:M, 1.7:1.

The median age was 9.29 with an interquartile range of 6.26 - 12.2.

Of interest median weight was 39.2Kg which at 9.29 years is > 91st centile for both boys and girls.

16 of 46 (34%) patients received prednisolone. The duration of treatment varied from 3 days to two weeks.

Four children received acyclovir.

An MRI brain was carried out in seven children. All MRIs were normal.

Three children were referred to neurology, in two instances referral was sent for recurrent episodes, the third child had torticollis associated with facial palsy.

Follow up telephone contact with 35 of 46 parents revealed that in all of these patients the condition had resolved, including the 3 children referred to neurology. However interestingly 7 parents can still

notice slight facial asymmetry particularly when the child is tired.

Conclusion: In uncomplicated Bell's Palsy in a child with a normal examination, further investigations are generally not necessary. From our experience steroids and aciclovir are not required for Bell's Palsy in children.

724

HEAD INJURIES IN CHILDREN : IS IT NICE TO CT SCAN?

N. Ganjoo¹, D. Stanciu²

¹Department of Paediatrics,, Addenbrookes Hospital, Cambridge, ²Department of Paediatrics, Colchester University Hospitals NHS Trust, Colchester, UK

Introduction: Trauma is commonest cause of hospital admission in children and head injuries account for 75% of children admitted. Head injury is a common cause of acquired neurological and cognitive disability in childhood. Indications for CT scan in children with head injury are well documented in NICE Guidelines,UK.

Aim: To evaluate our practice for children attending A&E department with head injury at district general hospital in comparison with NICE guidelines.

Method: This was a retrospective audit of case notes of children over a 2months period, attending the emergency department with head injury. Case notes were analysed to record Glasgow coma scale, children admitted for neurology observation and whether the NICE guidelines for CT scan was followed.

Results: Of 120 children with head injury who attended A&E, 107 full case notes were available. In total 13/107 children (12%) met the criteria for CT scan. However 6/13 children (46%) were sent home after clinical assessment by a doctor. 7/13 (54%) were admitted for 24 hours neurology observation of which 3/13 underwent CT scan. Head injury leaflet was given to all children sent home.

Conclusions: This study reiterates continual awareness of NICE guidelines amongst physicians for highest quality care. However as evident in this study, it may require a four fold increase in the number of CT scans done. This may not be possible at a district general hospital due to local constraints of radiology and can be circumvented by proper