assignment of a diagnostic score in clinical practice can be useful to apply early environmental and dietary prevention measures.

770

THE STATE OF CYTOKINE LINKS OF IMMUNITY AT HEPATOGENIC ULCERS AND WAYS OF CORRECTION

O.M. Horlenko¹, O.M. Moskal¹, E.I. Arshiy²,
F.V. Horlenko³, B.M. Halay²

¹Pediatrics with Infectious Diseases, ²Internal Medicine, ³Surgical Diseases, Uzhgorod National University, Medical Faculty, Uzhgorod, Ukraine

Background and aims: The great number of researches is confirmed, that development of ulcerous illness and chronic gastritis is accompanied violations of cytokine station. They present are given about the change of products of proinflammatory and anti-inflammatory interleukins in the mucus shell of stomach and duodenum.

Methods: We study the levels of IL-1 β , IL-2, IL-4, IL-8 in the serum blood of patients with hepatogenic ulcers and erosions which complicated the chronic diffuse diseases of liver (CHDDL). We observed 120 patients, amonge them 72 patients with the cirrhosis of liver (CL) and 48 patients by chronic active hepatitis (CHAH) by alcoholic etiology. The I-st group of patients used the base treatment at CL with including - "Ursofalk" and "Dufalac", 1 month, II-nd group used the base treatment.

Result: In I-st group of patients levels of IL-1 β , IL-4 under used of treatment decreased to the reference dates. In the II-nd group of patients the levels of all of interleukins after treatment remained higher normal indexes. There were following reliable differential levels of interleukins of I-st and II-nd groups after treatment: concentration of IL-1 β ($p < 0,001$), IL-4 ($p < 0,01$). The indexes of interleukins in the serum blood of patients of the II group remain higher than indexes of I of group.

Conclusions: After used complex treatment concentration of IL-1 β decreased in 2 times comparatively with an initial level ($p < 0,01$), concentration of IL-2 in 1,7 times ($p < 0,05$), concentration of IL-4 in 1,2 times ($p > 0,05$), concentration of IL-8 in 1,4 times ($p > 0,05$).

771

THE MANAGEMENT OF ACUTE GASTROENTERITIS IN CHILDREN IN PORTUGAL

H. Antunes^{1,2}, L. Januário³, H. Flores⁴, J. Cabral⁴,
F. Santos⁴, I. Pó⁴, A.I. Lopes⁵

¹Life and Health Sciences Research Institute (ICVS), ²Gastroenterology, Hepatology and Nutrition Unit, Pediatric Dpt. on behalf of the Gastroenterology Section of the Paediatric Portuguese Society (SPP) and on behalf of the SPP, Braga Hospital, Braga, ³Pediatric Dp. on behalf of the SPP, Pediatric Hospital, Coimbra, ⁴Pediatric Dpt. on behalf of the Gastroenterology Section of the SPP, Estefânia Hospital, ⁵Pediatric Dpt. on behalf of the Gastroenterology Section of the SPP, S. Maria Hospital, Lisboa, Portugal

Aim: To evaluate the knowledge/practice in the AGE's management in children.

Methods: An anonymous semi-structured questionnaire(21 questions) was applied to paediatricians, paediatrician fellows(PF) and general practitioners(GP) from July 7 to September 25, 2009,in the site of SPP. The Zprobe was used to compare paediatricians/PF(1);paediatricians/GP(2).

Results: There were 430 answers:paediatricians 261(27.3% registered paediatricians in the national Board), PF 94, GP 83. They answered they knew ESPGHAN'recommendations for Oral Rehydration Solution(ORS) use: paediatricians 82.0%;PF 76.6%;GP 41.0%,(1) $p=0.220$;(2) $p < 0.001$. The sodium recommended ORS content was correctly answered by paediatricians:68.6%; PF:69.1%; GP:32.5%,(1) $p=0.397$;(2) $p < 0.001$. ORS was reported to be used in 86.3% of cases (paediatricians:88.5%; PF:95.7%;GP:68.7%,(1) $p=0.017$;(2) $p < 0.001$); and home made "ORS" in 12.8% (GP:"yes":25.3%;PF:"no":57.4%,(2) $p < 0.001$). Nineteen percent answered they would ever recommend stopping diet. The delay until reassuming normal diet was respectively:< 4hours(h)54.0%(normal diet interval); 4-6h:38.7%; >6h:7.9%. Maintenance of breastfeeding was reported in 96.3%; the use of lactose-free milk or diluted milk was not recommend in 53.0%(paediatricians:53.4%; PF:69.1%; GP:33.7%,(1) $p=0.008$;(2) $p < 0.001$) and 86.5%(pediatricians:85.8%; PF:94.7%; GP:79.5%,(1) $p=0.008$;(2) $p=0.176$), respectively. The recommendation to modify the diet was

assumed in 10.5% (paediatricians: 6.5%; PF: 1.1%; GP: 33.8%, (1) $p=0.005$; (2) $p<0.001$) and the reported use of juices/cola was respectively: juices: paediatricians: 2.7%; PF: 0%; GP: 3.6%, (2) $p=0.367$ and cola: paediatricians: 5.0%; PF: 0%; GP: 13.3%, (2) $p=0.045$. Drugs have been used in 26.8% (9.7% if we excluded racecadotril and probiotics). Personal preferences in the presentation of ORS included mostly pre-prepared solutions (paediatricians: 64.0%; PF: 66.0%; GP: 47.0%, (1) $p=0.377$; (2) $p=0.010$).

Conclusions: The practice in the AGE' management in children in Portugal is mostly based on ESPGHAN recommendations. The paediatricians/PF seemed to have a better knowledge/practice than GP, emphasizing the need to spread these recommendations to all professionals involved in paediatric care.

772

FIVE YEARS EXPERIENCE IN LIVER TRANSPLANTATION IN BULGARIA

R. Mitrova¹, H. Jelev², J. Uzunova¹,
R. Alamurova¹, D. Christova¹

¹University Hospital Lozenetz, ²Paediatric University Hospital, Sofia, Bulgaria

First successful liver transplantation in Bulgaria was performed in 2004. The first transplanted child was a seven months old girl with biliary atresia. It was a living donor transplantation, donor was the father.

Since then 17 transplantations were performed. We discuss the early post transplantation period, complications, mortality and survival.

773

PREDICTIVE FACTORS OF SURGERY IN NECROTIZING ENTEROCOLITIS

J. Rousseaux¹, J. Mcheik², J.P. Saulnier³, L. Bott¹,
K. Hussein¹

¹Neonatal Intensive Care Unit, ²Pediatric Surgery Department, ³Pediatric Intensive Care Unit, University Hospital of Poitiers, Poitiers, France

Background: Previous studies have shown controversy in surgical or medical management of necrotizing enterocolitis (NEC).

Objective: To study perinatal predictive factors of surgery in monocentric NEC population.

Material and methods: A retrospective study

conducted in Neonatal Intensive Care unit from 1st January 2003 to the 31st December 2008. Fetal and neonatal clinical characteristics were assessed for 33 NEC cases, stages II or more of Bell classification (mean GA $\pm$ SD = 28.9 $\pm$ 2.4 wks; birth weight = 1156 $\pm$ 416 g). Data were compared between A subgroup of 11 infants with surgical management and B subgroup of 22 infants with medical management.

Results: There was no significant difference in mean GA and birth-weight between A and B subgroups, respectively (27.6 $\pm$ 2.3 vs 28.9 $\pm$ 2.5 wks; $p=0.13$) and (1055 $\pm$ 355 g vs 1208 $\pm$ 439 g; $p=0.29$). NEC had occurred at 29 $\pm$ 15.8 and 29.8 $\pm$ 13.5 days respectively in A and B groups, leading to death in 4/11 vs 7/22. Surgery was associated with high bell classification stages [6/11 (3a) and 4/11 (3b), groupe A vs 4/22 (3a) and 1/22 (3b), groupe B; $p=0.02$]. Surgery was associated with significantly higher incidence of nosocomial infection (9/11 vs 12/22; $p=0.05$), antenatal mother's urinary tract infection (4/11 vs 1/22; $p=0.015$), dopamine use for hemodynamic disorders (8/11 vs 8/22; $p=0.02$) and more vaginal delivery (9/11 vs 9/22; $p=0.02$). Multivariate analysis showed vaginal delivery as an independent predictive factor of surgery in NEC odd ratio 25; IC 95% [1.2-520], $p=0.04$.

Conclusion: Our study shows that surgery management of NEC was associated with significantly higher incidence of vaginal birth.

774

ANTIBIOTICS THERAPY IN MILD NONTYPHOID SALMONELLOSIS CHILDREN PROLONGED HOSPITALIZATION AND FECAL EXCRETION TIME-A PRELIMINARY REPORT

I.-F. Huang^{1,2}

¹Department of Pediatrics, Kaohsiung Veterans General Hospital, Kaohsiung, ²National Yang-Ming University, Taipei, Taiwan R.O.C.

Objective: Antibiotics for nontyphoid salmonellosis has long been a controversial matter. Most studies didn't classify the salmonellosis with severity of disease. The purposes of the study were to investigate the outcomes of the mild nontyphoid salmonellosis children with antibiotics therapy.

Material and methods: This is a prospective observational study. Between 2005 and 2008, all pediatric patients with nontyphoid salmonellosis admitted to Kaohsiung Veterans General Hospital,