Bifidobacterium bifidum in a rat model of necrotizing enterocolitis: antimicrobial peptide and protein responses

Mark A. Underwood¹, Anchasa Kananurak², Christine F. Coursodon³, Camille K. Adkins-Reick³, Hiutung Chu², Stephen H. Bennett¹, Jan Wehkamp^{2,4}, Patricia A. Castillo², Brian C. Leonard², Daniel J. Tancredi¹, Michael P. Sherman^{1,4}, Bohuslav Dvorak³ and Charles L. Bevins²

INTRODUCTION: Necrotizing enterocolitis (NEC) is a devastating disease of premature infants. Probiotics decrease the risk of NEC in clinical and experimental studies. Antimicrobial peptides protect the gut against noxious microbes and shape the commensal microbiota, but their role in NEC remains unclear.

METHODS: To investigate the expression of antimicrobial peptides in experimental NEC and the impact of probiotics on their expression, premature rats were divided into three groups: dam fed (DF), hand fed with formula (FF), or hand fed with formula containing *Bifidobacterium bifidum* (FF + BIF). All groups were exposed to asphyxia and cold stress.

RESULTS: Like in human ontogeny, the rat pup has low expression of Paneth cell antimicrobials, which increases rapidly during normal development. The expression of lysozyme, secretory phospholipase A_2 (sPLA₂), pancreatic-associated proteins 1 and 3 mRNA was elevated in the FF group with a high incidence of NEC, as compared with the DF and FF + BIF groups where the disease was attenuated.

DISCUSSION: We conclude that induction of antimicrobial peptides occurs in experimental NEC similar to that reported in human disease and is attenuated when disease is averted by probiotic *B. bifidum*. The induction of antimicrobial peptides is likely an adaptive mucosal response that is often not sufficient to prevent disease in the premature gut.

Necrotizing enterocolitis (NEC) is common in premature infants, with an incidence of 3–10% and a mortality rate for severe disease of 25–40% (1,2). The pathophysiology of this devastating disease remains poorly understood (3). The major risk factors underscore the central role of host-microbe interactions: immaturity of the intestine (structural, functional, and immunologic), enteral feedings, and dysbiosis (imbalance in the intestinal microbiota) (3,4). Probiotics are dietary supplements containing live health-promoting bacteria. Several well-designed clinical trials have demonstrated a decreased incidence of NEC in premature infants receiving probiotic microorganisms (5). In the neonatal rat model of NEC, oral administration of the probiotic *Bifidobacterium bifidum* protects the small intestine against the disease (6,7). This protective effect is associated with the reduction of inflammation, stabilization of the mucus layer, improvement in intestinal integrity, and decreased epithelial apoptosis (7,8).

A variety of immune mechanisms protect the intestinal mucosa both by mediating homeostatic interactions with the lumenal microbiota and by protecting from pathogenic microbes. Central to the innate immune defenses are physical and chemical barrier components, multiple pathways for microbe detection and response, and a collection of constitutive and inducible antimicrobial peptides and small proteins (henceforth referred to simply as antimicrobial peptides). Antimicrobial peptides are important in the immune function of phagocytic leukocytes and many epithelial cells. In the small intestine, antimicrobial peptides are expressed by Paneth cells, enterocytes, and leukocytes of the lamina propria (9).

Paneth cells, located at the base of the crypts of Lieberkühn in the small intestine, sense bacteria that penetrate the mucus layer, and, in response, secrete large amounts of antimicrobial peptides that protect the intestinal mucosa, shape the intestinal microbiota, and protect the adjacent intestinal stem cells (9). Given the importance of the chemical barrier provided by Paneth cell antimicrobial peptides (4), and the involvement of intestinal microbes in disease pathogenesis, Paneth cell antimicrobial secretions likely play a role in protecting the intestinal mucosa from NEC (10).

In the human fetus, Paneth cells express antimicrobial peptides very early in gestation, long before exposure to bacteria, although at relatively low levels that increase with gestation (11). These observations generated the hypothesis that immaturity of Paneth cell function contributes to the increased risk of NEC in premature infants. Infants with NEC had higher

Received 23 May 2011; accepted 10 January 2012; advance online publication 7 March 2011. doi:10.1038/pr.2012.11

¹Department of Pediatrics, University of California–Davis, Sacramento, California; ²Department of Microbiology and Immunology, University of California–Davis, Davis, California; ³Department of Pediatrics, University of Arizona, Tucson, Arizona; ⁴Current affiliation: Dr. Margarete Fischer-Bosch-Institute of Clinical Pharmacology, Stuttgart, Germany (J.W.); Department of Child Health, University of Missouri–Columbia, Columbia, Missouri (M.P.S.). Correspondence: Mark A. Underwood (mark.underwood@ucdmc.ucdavis.edu)

Antimicrobial peptides in NEC

Articles

numbers of Paneth cells and greater antimicrobial peptide mRNA expression at the time of surgery than control infants with ileal atresia, suggesting that Paneth cell antimicrobials were increased in response to NEC (12), but that the fledgling response was inadequate in severe NEC. A contemporary study reported that some infants with NEC had very few lysozyme-positive Paneth cells, consistent with Paneth cell destruction, degranulation, or deficiency (13). A larger, more recent analysis demonstrated no change in Paneth cell numbers or expression of Paneth cell antimicrobials during acute NEC in premature infants, but reported marked increases in Paneth cell numbers and expression of human defensin (HD) 5 during recovery from NEC (14). Paneth cell metaplasia in the colon was also noted during recovery, a finding that has been reported previously in inflammatory bowel disease, but not NEC (14).

The aims of this study were: first, to define the development of intestinal antimicrobial molecule expression in the healthy rat pup, and, second, to analyze the expression of these products in a well-established rat model of NEC with and without the probiotic *B. bifidum*. We report that the rat pup, similar to premature infants, has low expression of Paneth cell antimicrobials that increases during development. Furthermore, we report an induction of antimicrobial peptide expression in experimental NEC that is attenuated when disease is averted by probiotic administration.

RESULTS

The rat pup is born with an immature intestinal tract that matures during the first 21 days of extrauterine life, modeling the changing intestinal tract of the premature infant. The rat small intestine contains Paneth cells and secretes antimicrobial peptides that are orthologous to those of the human (Table 1).

Table 1. Human and rat intestinal antimicrobial peptides and prot
--

Species	Defensins ^a	C-type lectins ^ь	Enzymes ^c
Human	HNP 1–4	HIP/PAP	Lysozyme
	HD 5–6	PSP/Reg	sPLA ₂
	BD 1-4		
Rat	NPs 1–4	PAP1	Lysozyme
	Cryptdins 5–6	PAP3	sPLA ₂
	rBDs 1–3	PSP/Reg	

BD, β -defensin; HD, human defensin; HIP, hepatocarcinoma-intestine-pancreas; HNP, human neutrophil peptide; NP, neutrophil peptide; PAP, pancreatic-associated protein; PSP, pancreatic secretory peptide; Reg, islet regenerating peptide; sPLA₂, secretory phospholipase A₂.

^aHumans produce α-defensins in neutrophils (HNPs 1–4) and in Paneth cells (HD5 and HD6) and BD in skin and all mucosal surfaces. The rat produces similar α-defensins in neutrophils (NP) and Paneth cells (cryptdins) and β-defensins (rBD) in all mucosal surfaces. In addition, the rat α-defensin NP3 is expressed in both Paneth cells and circulating neutrophils.^bA family of C-type lectins that bind to peptidoglycan, and thus are highly selective against Gram-positive bacteria, are expressed in the small intestine. Members of this family are named PSP/Reg and PAP/RegIII. In the human gut, PSP/Reg is expressed in Paneth cells, and PAP/RegIII (also known in the human as HIP/PAP) is expressed in both enterocytes and Paneth cells. The rat intestine expresses one PSP/Reg of Nameth cells. Lysozyme is also expressed in macrophages and dendritic cells.

Antimicrobial Gene Expression Increases Exponentially in the Developing Rat Pup

Figure 1 summarizes the mRNA expression of antimicrobial peptides in development from 3 days of age up to adulthood. Note that in order to display the wide range of expression (>10,000-fold), the *y*-axis is a log scale. For the enzymes lysozyme and secretory phospholipase A, (sPLA,), the a-defensins (cryptdins) 5 and 6, and the defensin-processing enzyme matrix metalloproteinase (MMP) 7, the rapid change approximated a logarithmic increase, with correlation coefficients that ranged from 0.71 for lysozyme to 0.94 for MMP7. For the C-type lectins pancreatic-associated proteins (PAP) 1 and 3 and the α -defensin neutrophil peptide (NP) 3, expression changed minimally in the rat pup but dramatically increased at the time of weaning (day 21) and remained high in the adult. The housekeeping gene GAPDH did not vary significantly during development. Expression of cathelicidin-related antimicrobial peptide (15), pancreatic secretory peptide/islet regenerating peptide (Reg) and BD2 (a β -defensin) (16) were at or below





Articles Underwood et al.



Figure 2. Antimicrobial peptides in development. Immunohistochemistry analyses of healthy rat ileum at 3, 5, 21, and 38 (adult) days utilizing antibodies against lysozyme (Ly) and PAP/RegIII (PAP) and at days 21 and 38 against NP3. Original magnification ×40 (for 3- and 5-day specimens) and ×20 (for 21- and 38-day specimens). Bar = 100 μ m. NP, neutrophil peptide; PAP, pancreatic-associated protein; RegIII, regenerating protein.

the level of reliable detection (100 mRNA copies/10 ng total RNA) throughout development (data not shown).

Antimicrobial Peptides Are Variably Expressed in the Developing Rat Pup

Sections of ileum from rat pups of varying ages stained with antibodies to lysozyme, NP3, and PAP/RegIII (stains both PAP1 and PAP3) are presented in **Figure 2**. Lysozyme and NP3 were strongly expressed in adult Paneth cells, but were not detectable in the youngest rat pups. There has been some controversy regarding the localization of PAP/RegIII in the small intestine. In the mouse, the orthologous molecule RegIII γ is expressed in the secretory granules of the Paneth cell (17). In the rat, early studies immunolocalized PAP/RegIII to the enterocytes of the lower villus (18). We found that PAP/RegIII was present in the enterocytes even in the youngest rats, with minimal signal noted in the lower villi and crypts.

Antimicrobial Gene Expression Is Increased in the NEC Rat Model and Attenuated by *B. bifidum*

As shown previously (7), the incidence of histologic NEC was 0% in the dam-fed (DF) group, 57% in the hand fed with formula (FF) group, and 17% in the hand fed with formula containing *Bifidobacterium bifidum* (FF + BIF) group ($P < 0.01 \chi^2$ analysis). Administration of *B. bifidum* reduced the median histologic score to 1.0 compared with 2.0 in the FF group ($P \le 0.01$) (7).

The small intestinal mRNA expression of antimicrobial enzymes, defensins, and C-type lectins is summarized for the three groups of rat pups in **Figure 3**. The comparisons between rat pups developing NEC (FF) and the other groups that demonstrated significant differences are denoted with asterisks (*P < 0.05, **P < 0.01, †P < 0.001). Significant increases in expression of lysozyme, sPLA₂, PAP1, and PAP3



Figure 3. Antimicrobial peptide mRNA in NEC. Fold increases above the DF group in mRNA expression of Paneth cell products: (**a**) enzymes, defensins, and the housekeeping gene *GAPDH*, (**b**) PAP1, and (**c**) PAP3. All rat pups were exposed to asphyxia and cold stress. White bars: damfed, black bars: formula fed, gray bars: fed with formula supplemented with *Bifidobacterium bifidum*. N = 8 for each group. Error bars are 95% confidence intervals for the fold change. *P < 0.05, **P < 0.01, $^{+}P < 0.001$. DF, dam fed; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; PAP, pancreatic-associated protein; sPLA₂, secretory phospholipase A₂.

occurred in the FF group compared with the DF group and of lysozyme, sPLA,, and PAP1 in the FF group compared with the FF + BIF group. Cryptdins 5 and 6 demonstrated a similar pattern that did not reach statistical significance. The error bars in Figure 3 represent the 95% confidence intervals for fold change of expression of the FF and FF + BIF groups over the DF group and allow an additional comparison (i.e., if the 5% confidence interval is >1, the difference is significant). Expression of pancreatic secretory peptide/Reg, MMP7, cathelicidin-related antimicrobial peptide, and NP3 was low in all three groups, with no statistically significant differences between groups (data not shown). Rat pups born by cesarean section 1 day early and fed by surrogate dams but not exposed to cold stress and asphyxia were similar to the dam-fed stressed group in expression of all antimicrobial molecules tested (data not shown).

Detection of Antimicrobial Molecules in the NEC Rat Model

Sections of ileum from rat pups in the DF and FF groups showed positive staining for lysozyme in discrete cells in the lamina propria, with no difference between groups, but no staining of epithelial cells (data not shown). We could not detect staining of NP3 in either group, suggesting that expression is too low for detection at this developmental stage. In spite of the dramatic increase in PAP1 and PAP3 in the FF group at the mRNA level, attempts to demonstrate differences between DF and FF tissue at the protein level with immunohistochemistry and western blotting using three different antibodies were unsuccessful. Whether this reflects no true change at the protein level or challenges specific to the antibodies selected is unclear.

DISCUSSION

NEC remains a devastating disease of premature infants. Immaturity of the integrity of the intestinal barrier (physical, chemical, and functional) and differences in the intestinal microbiota appear to be important predisposing factors (3,4,19). Clinical studies indicate that probiotics are currently among the most promising preventive interventions (5,20,21). Our previous studies with *B. bifidum* OLB6378 have shown decreased intestinal injury and improved integrity of the intestinal barrier (7) and downregulation of intestinal apoptosis (8) in the rat NEC model.

This study was designed to investigate a central component of the mucosal innate immune system of the rat pup: antimicrobial peptides in development and in the rat NEC model. Our results show that as the rat pup matures toward weaning, mRNA expression of several intestinal antimicrobials increases from 1,000- to 10,000-fold. These changes are most marked for lysozyme, sPLA₂, cryptdin 5, and cryptdin 6, all of which are produced in Paneth cells, but not enterocytes. PAP1 is highly expressed even in the very young rat pup; both PAP1 and PAP3 increase >100-fold near the time of weaning. NP3 is an α -defensin that is produced in both the neutrophil and the Paneth cell in the rat. MMP7 is a metalloproteinase that cleaves the inactive forms of the Paneth cell cryptdins to the active forms in rodents. Both of these Paneth cell products increase dramatically near the time of weaning.

Several differences in intestinal antimicrobial development between the mouse, the rat, and the human are noteworthy. In the human fetus, Paneth cells are identifiable by ultrastructural analysis and reverse transcriptase-PCR detection of HD5 and HD6 as early as 14-17 weeks gestation with increases in mRNA by northern analysis of 40-250fold between premature infants and adults (11). In mice, the appearance of lysozyme-staining Paneth cells between 4 and 7 days of life correlates with the development of resistance to infection with pathogenic Shigella (22). Expression of mouse intestinal defensins increases 100-1,000-fold with development from the newborn to adult (23). The low expression of cathelicidin-related antimicrobial peptide in the young rat pup is in striking contrast to the mouse, in which constitutive expression is observed for the first 2 weeks after birth and then gradually decreases (23). In the mouse, Paneth cells are first identifiable at 7-14 days of life (22,24). Our ability to identify the initial appearance of Paneth cells, the sources of antimicrobial peptides in the young rat pup, and changes in Paneth cell numbers in the rat NEC model is hampered by the lack of antibodies to Paneth cell–specific peptides such as cryptdin 5 and cryptdin 6.

It is likely that the expression of these different antimicrobial products in the rat reflect multiple mechanisms of control as reported in the mouse. RegIII γ (the mouse PAP/ RegIII ortholog) and MMP7 appear to be inducibly expressed with minimal expression in germ-free mice and marked increases upon exposure to the intestinal microbiota (17,25). In contrast, the mouse α -defensins (cryptdins) appear to be expressed constitutively with little or no differences between germ-free and conventionally raised mice (26). Our data in the rat support the notion that lysozyme, sPLA₂, cryptdin 5, and cryptdin 6 increase rapidly with development, whereas PAP1, PAP3, NP3, and MMP7 increase at about the time of weaning, likely in response to changes in the intestinal microbiota.

The dramatic increases during development of rat intestinal antimicrobial expression parallel the more limited observations in human fetuses and premature infants. These observations suggest a possible association between the diminished Paneth cell products in premature infants and the susceptibility of these infants to NEC. Specifically, low levels of antimicrobial peptide expression in premature infants (and young rat pups) may contribute to NEC susceptibility by mechanisms that involve either an inadequate response to opportunistic pathogens within the microbiota, and/or an inability to shape the developing microbiota in an *ex utero* environment that is not physiologic (the "normal" milieu for the premature intestinal mucosa is to be bathed by swallowed amniotic fluid that is generally free from bacteria and bacterial antigens (27,28)).

To further explore this hypothesis, we utilized a well-established rat model of NEC. In this model, experimental NEC was attenuated by concurrent administration of B. *bifidum* (6), and completely obviated in rats remaining with and fed by surrogate dams but exposed to the same levels of asphyxia and cold stress. Our analyses demonstrate that the mRNA expression of several antimicrobials is significantly increased in the FF rat pups that develop NEC, compared with DF controls exposed to the same asphyxia and cold stress that do not develop NEC. The pups that received formula supplemented with B. bifidum were relatively protected from NEC and had correspondingly minimal increases in expression of the antimicrobial peptides. Intestinal antimicrobial peptides are induced in NEC at the mRNA level, but the levels are low and difficult to detect at the protein level with available reagents, and thus may be inadequate to defend the immature gut from invasion and disruption by microorganisms.

It is possible that the increase in antimicrobial peptides in the FF pups is a response to mucosal inflammation. Indeed, in the colon, β -defensins 2 and 3 are induced by proinflammatory cytokines such as interleukin-1 β and tumor necrosis factor- α (29). Likewise, adults with ankylosing spondylitis have increased expression of the α -defensin HD5, lysozyme, and sPLA₂ in the presence of chronic inflammation, but not in the presence of acute inflammation or normal histology (30). On

Articles

the other hand, similar changes have not been demonstrated in other diseases of chronic intestinal inflammation (31). In ileal Crohn's disease, decreased expression of HD5 and HD6 is observed, and more significant reductions are closely related to a specific mutation in the nucleotide-binding oligomerization domain containing 2 receptor; these changes did not correlate with inflammation (32). In colonic Crohn's disease and ulcerative colitis, Paneth cell metaplasia results in increased expression of α -defensins that is isolated to the colon. In these various investigations there is a lack of evidence for induction of Paneth cell antimicrobial peptides in acute inflammation, therefore we favor the explanation that the noted increases observed in this study are a response to disease (albeit insufficient for prevention) rather than a response to inflammation.

One proposed mechanism of probiotic action is stimulation of innate antimicrobial production. The impact of probiotics on expression of antimicrobial peptides has been studied for one of the inducible β -defensins expressed in the colon with mixed results (33–35). Our results suggest that *B. bifidum* does not increase the expression of small intestinal antimicrobial products.

In conclusion, these studies show ontogenesis of antimicrobial peptides in the small intestine of neonatal rats under normal physiologic conditions. The development of experimental NEC is associated with increases in mRNA expression of several of these antimicrobial molecules at the site of injury. Oral administration of probiotic *B. bifidum* OLB6378 can avert both NEC and the associated increase in expression of antimicrobial peptides.

METHODS

Antimicrobial Expression in the Rat Pup

This protocol was approved by the Animal Care and Use Committee of the University of California–Davis. To evaluate the mRNA expression in the developing rat pup, healthy Sprague–Dawley rat pups were raised with their dams and killed at 3, 5, 10, 14, and 21 days of life (n = 6 at each time point) and compared with healthy adult rats (n = 2).

Experimental NEC Model

This protocol was approved by the Animal Care and Use Committee of the University of Arizona. Neonatal Sprague–Dawley rats (Charles River Labs, Pontage, MI) were collected by cesarean section 1 day before scheduled birth, and their first feeding started 2 h after delivery. Rat pups were hand fed every 6 h with rat milk substitute formula (FF, n = 30), formula containing 5×10^6 colony forming units per day of *B. bifidum* OLB6378 (NITE BP-31, Meiji Dairies, Odawara, Japan; FF + BIF, n = 30), or dam fed by surrogate mothers (DF, n = 16). All animals were exposed to asphyxia (breathing 100 % nitrogen gas for 60 s) and cold stress (4 °C for 10 min) twice daily (7,8). After 96 h, all of the surviving animals were killed by decapitation.

RNA Preparation and Real-Time PCR

For the healthy developmental rat studies (University of California-Davis), total RNA was isolated from ileal tissue using cesium chloridegradient ultracentrifugation (36). For the rat NEC studies (University of Arizona), total RNA was isolated from ileal tissue using the RNeasy Mini Kit (Qiagen, Santa Clarita, CA) (37). Real-time PCR for all specimens was performed using single-stranded cDNA from tissue with specific oligonucleotide primer pairs (Table 2) in a temperature cycler equipped with a fluorescence detection monitor (LightCycler; Roche Diagnostics, Mannheim, Germany) as previously described (38,39). A negative control reaction that omitted template cDNA was included with each set of reactions to check for possible contamination.

Table 2.	Primer s	ets for quant	itative reverse	-transcriptase PCR
----------	----------	---------------	-----------------	--------------------

Gene target	Sequence
Lysozyme sense	5'-CAAGCCATACAATGTGCGAAGAGAG-3'
Lysozyme antisense	5'-TGTTGGTTTGAGGGGAAAGCAAG-3'
sPLA ₂ sense	5'-CATTGTGGTGTGGGTGGCAGAG-3'
sPLA ₂ antisense	5'-TGGTTTGTAGAGCAGGAGATTTGGC-3'
Cryptdin 5 sense	5'-GACCAGGTTGTTTCTGTCTCCATTG-3
Cryptdin 5 antisense	5'-TGAGGCTTCCGTATCTCTTGTTGC-3'
Cryptdin 6 sense	5'-AGCAACCATCAGATGAGGACCAGG-3'
Cryptdin 6 antisense	5'-ACCTTGAGCACAGAACGCAGTGG-3'
PAP1 sense	5'-TGCCAGAAGAGACCTGAAGGACAC-3'
PAP1 antisense	5'-TTGTTACTCCACTCCATCCACCTC-3'
PAP3 sense	5'-CCAAGAACCCAACAGAGGTGGATG-3'
PAP3 antisense	5'-GGTCCCACAGTGACTTCCAGAGACAG-3'
PSP/Reg sense	5'-GTTTCTCTACAAATCCTGGGACACTGG-3'
PSP/Reg antisense	5'-TTGGGCATCACAACTGTTATCTCTCC-3'
NP3 sense	5'-TTTGGAGGGGATAAAGGCACTGC-3'
NP3 antisense	5'-TCAGCAACAGAGTCGGTAGATGCG-3'
MMP7 sense	5'-CGGCGGAGATGCTCACTTTGAC-3'
MMP7 antisense	5'-TGGCTCAGGAAGGGCGTTTGC-3'
CRAMP sense	5'-TGCCTCTAACCGTTTCCCAGACC-3'
CRAMP antisense	5'-TGCTCAGGTAACTGCTGTGATGCC-3'
BD2 sense	5'-TTTCTCCTGGTGCTGCTGTCGC-3'
BD2 antisense	5'-CCACAAGTGCCAATCTGTCGAAAAC-3'
GAPDH sense	5'-TGACCACAGTCCATGCCATCACTG-3'
GAPDH antisense	5'-ATGACCTTGCCCACAGCCTTGG-3'

BD, β -defensin; CRAMP, cathelicidin-related antimicrobial peptide; HD, human defensin; HIP, hepatocarcinoma-intestine-pancreas; HNP, human neutrophil peptide; NP, neutrophil peptide; PAP, pancreatic-associated protein; PSP, pancreatic secretory peptide; Reg, islet regenerating peptide; sPLA₂, secretory phospholipase A₃.

Gene-specific plasmid standards were included with each set of reactions. The PCR conditions were: initial denaturation at 95 °C for 10 min, followed by 45 cycles with each cycle consisting of denaturation (95 °C for 15 s), annealing (60 °C for 5 s), and extension (72 °C for 10 s).

NEC Evaluation

After termination, a 2-cm piece of distal ileum was removed and fixed in 70% ethanol, paraffin embedded, and stained with hematoxylin and eosin for histological evaluation of NEC. Pathological changes in intestinal architecture were evaluated by a blinded investigator, and a score of 0–4 was assigned according to a published NEC scoring system, with scores of 2–4 defined as NEC (40).

Immunohistochemistry

A 1–2 cm section of distal ileum was collected from each animal and fixed in either 70% ethanol or 10% formalin, paraffin embedded, sectioned at 4–6 μ m, and processed as described (12,41). The sections were blocked with 1.5% normal goat serum (Vector Laboratories, Burlingame, CA) in phosphate-buffered saline for 20 min, and then incubated with either rabbit anti-lysozyme polyclonal antibody (Zymed; Invitrogen, Carlsbad, CA) or mouse anti-rat RegIIIa monoclonal antibody (R&D System, Minneapolis, MN) in 1.5% blocking serum overnight at 4 °C, then washed with phosphate-buffered saline and incubated with either biotinylated anti-rabbit or anti-mouse secondary antibody (Vector Laboratories) in 1.5% blocking serum for 20 min at room temperature. Vectastain Elite ABC reagent (Vector Laboratories) was applied, and sections were incubated for another 20 min, followed by 3,3'-diaminobenzidine substrate for 5 min, then light green counterstain.

Antimicrobial peptides in NEC Articles



Statistics

Pearson correlations were used to determine how closely the rise of antimicrobial peptides approximated exponential growth. ANOVA was used to compare antimicrobial mRNA expression levels among experimental groups. For each animal and antimicrobial molecule, the mean of two replicate templates was used as the absolute expression level, which was then log-transformed to improve the validity of statistical inferences. To address the primary research hypothesis that expression levels are increased in NEC and that the probiotics can reduce these levels, the main ANOVA models included pairwise contrasts estimated between the FF group and each of the other two groups.

STATEMENT OF FINANCIAL SUPPORT

M.A.U. received funding from National Institutes of Health (NIH) HD059127 and the Children's Miracle Network; B.D. from NIH HD039657 and a gift from Meiji Dairies Corporation; C.L.B. from NIH AI32738; A.K. from the Development and Promotion of Science and Technology Talents Project, Thailand; H.C. and B.C.L. from NIH T32AI060555; and M.P.S. from NIH HD057744 and the Gerber Foundation. This study was also made possible by grant number UL1 RR024146 from the National Center for Research Resources, a component of the NIH, and NIH Roadmap for Medical Research

REFERENCES

- Eichenwald EC, Stark AR. Management and outcomes of very low birth 1. weight. N Engl J Med 2008;358:1700-11.
- 2. Henry MC, Moss RL. Necrotizing enterocolitis. Annu Rev Med 2009;60:111-24.
- Neu J, Walker WA. Necrotizing enterocolitis. N Engl J Med 3. 2011;364:255-64.
- Morowitz MJ, Poroyko V, Caplan M, Alverdy J, Liu DC. Redefining the 4 role of intestinal microbes in the pathogenesis of necrotizing enterocolitis. Pediatrics 2010;125:777-85.
- Guthmann F, Kluthe C, Bührer C. Probiotics for prevention of necrotising 5. enterocolitis: an updated meta-analysis. Klin Padiatr 2010;222:284-90.
- 6. Caplan MS, Miller-Catchpole R, Kaup S, et al. Bifidobacterial supplementation reduces the incidence of necrotizing enterocolitis in a neonatal rat model. Gastroenterology 1999;117:577-83.
- 7. Khailova L, Dvorak K, Arganbright KM, et al. Bifidobacterium bifidum improves intestinal integrity in a rat model of necrotizing enterocolitis. Am J Physiol Gastrointest Liver Physiol 2009;297:G940-9.
- 8. Khailova L, Mount Patrick SK, Arganbright KM, Halpern MD, Kinouchi T, Dvorak B. Bifidobacterium bifidum reduces apoptosis in the intestinal epithelium in necrotizing enterocolitis. Am J Physiol Gastrointest Liver Physiol 2010;299:G1118-27.
- 9 Bevins CL, Salzman NH. Paneth cells, antimicrobial peptides and maintenance of intestinal homeostasis. Nat Rev Microbiol 2011;9:356-68.
- Salzman NH, Underwood MA, Bevins CL. Paneth cells, defensins, and 10. the commensal microbiota: a hypothesis on intimate interplay at the intestinal mucosa. Semin Immunol 2007;19:70-83.
- 11. Mallow EB, Harris A, Salzman N, et al. Human enteric defensins. Gene structure and developmental expression. J Biol Chem 1996;271:4038-45.
- Salzman NH, Polin RA, Harris MC, et al. Enteric defensin expression in 12. necrotizing enterocolitis. Pediatr Res 1998;44:20-6.
- 13. Coutinho HB, da Mota HC, Coutinho VB, et al. Absence of lysozyme (muramidase) in the intestinal Paneth cells of newborn infants with necrotising enterocolitis. J Clin Pathol 1998;51:512-14.
- 14. Puiman PJ, Burger-Van Paassen N, Schaart MW, et al. Paneth cell hyperplasia and metaplasia in necrotizing enterocolitis. Pediatr Res 2011;69:217-23.
- 15. Termén S, Tollin M, Olsson B, Svenberg T, Agerberth B, Gudmundsson GH. Phylogeny, processing and expression of the rat cathelicidin rCRAMP: a model for innate antimicrobial peptides. Cell Mol Life Sci 2003;60:536-49.
- 16. Jia HP, Mills JN, Barahmand-Pour F, et al. Molecular cloning and characterization of rat genes encoding homologues of human β -defensins. Infect Immun 1999;67:4827-33.
- 17. Cash HL, Whitham CV, Behrendt CL, Hooper LV. Symbiotic bacteria direct expression of an intestinal bactericidal lectin. Science 2006;313:1126-30.

- 18. Iovanna JL, Keim V, Bosshard A, et al. PAP, a pancreatic secretory protein induced during acute pancreatitis, is expressed in rat intestine. Am J Physiol 1993;265:G611-8.
- 19. Sharma R, Tepas JJ 3rd. Microecology, intestinal epithelial barrier and necrotizing enterocolitis. Pediatr Surg Int 2010;26:11-21.
- 20. Szajewska H. Probiotics and prebiotics in preterm infants: where are we? Where are we going? Early Hum Dev 2010;86 Suppl 1:81-6.
- 21. Sherman MP. New concepts of microbial translocation in the neonatal intestine: mechanisms and prevention. Clin Perinatol 2010;37:565-79.
- 22. Fernandez MI, Regnault B, Mulet C, et al. Maturation of paneth cells induces the refractory state of newborn mice to Shigella infection. J Immunol 2008;180:4924-30.
- 23. Ménard S, Förster V, Lotz M, et al. Developmental switch of intestinal antimicrobial peptide expression. J Exp Med 2008;205:183-93.
- 24. Bry L, Falk P, Huttner K, Ouellette A, Midtvedt T, Gordon JI. Paneth cell differentiation in the developing intestine of normal and transgenic mice. Proc Natl Acad Sci USA 1994;91:10335-9.
- 25. Mukherjee S, Vaishnava S, Hooper LV. Multi-layered regulation of intestinal antimicrobial defense. Cell Mol Life Sci 2008;65:3019-27.
- 26. Putsep K, Axelsson LG, Boman A, et al. Germ-free and colonized mice generate the same products from enteric prodefensins. J Biol Chem 2000;275:40478-82.
- 27. DiGiulio DB, Romero R, Kusanovic JP, et al. Prevalence and diversity of microbes in the amniotic fluid, the fetal inflammatory response, and pregnancy outcome in women with preterm pre-labor rupture of membranes. Am J Reprod Immunol 2010;64:38-57.
- 28. Underwood MA, Gilbert WM, Sherman MP. Amniotic fluid: not just fetal urine anymore. J Perinatol 2005;25:341-8.
- 29. Schröder JM, Harder J. Human β-defensin-2. Int J Biochem Cell Biol 1999;31:645-51.
- 30. Ciccia F, Bombardieri M, Rizzo A, et al. Over-expression of paneth cell-derived anti-microbial peptides in the gut of patients with ankylosing spondylitis and subclinical intestinal inflammation. Rheumatology (Oxford) 2010;49:2076-83.
- Jäger S, Stange EF, Wehkamp J. Antimicrobial peptides in gastrointestinal 31. inflammation. Int J Inflam 2010;2010:910283.
- 32. Bevins CL, Stange EF, Wehkamp J. Decreased Paneth cell defensin expression in ileal Crohn's disease is independent of inflammation, but linked to the NOD2 1007fs genotype. Gut 2009;58:882-3;discussion 883.
- 33. Schlee M, Harder J, Köten B, Stange EF, Wehkamp J, Fellermann K. Probiotic lactobacilli and VSL#3 induce enterocyte β-defensin 2. Clin Exp Immunol 2008;151:528-35.
- 34. Amit-Romach E, Uni Z, Reifen R. Multistep mechanism of probiotic bacterium, the effect on innate immune system. Mol Nutr Food Res 2010;54:277-84.
- 35. Möndel M, Schroeder BO, Zimmermann K, et al. Probiotic E. coli treatment mediates antimicrobial human β-defensin synthesis and fecal excretion in humans. Mucosal Immunol 2009;2:166-72.
- 36. Chirgwin JM, Przybyla AE, MacDonald RJ, Rutter WJ. Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. Biochemistry 1979;18:5294-9.
- 37. Clark JA, Doelle SM, Halpern MD, et al. Intestinal barrier failure during experimental necrotizing enterocolitis: protective effect of EGF treatment. Am J Physiol Gastrointest Liver Physiol 2006;291:G938-49.
- 38. Bustin SA. Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. J Mol Endocrinol 2000:25:169-93.
- 39. Wehkamp J, Chu H, Shen B, et al. Paneth cell antimicrobial peptides: topographical distribution and quantification in human gastrointestinal tissues. FEBS Lett 2006;580:5344-50.
- 40. Dvorak B, Halpern MD, Holubec H, et al. Epidermal growth factor reduces the development of necrotizing enterocolitis in a neonatal rat model. Am J Physiol Gastrointest Liver Physiol 2002;282:G156-64.
- 41. Moore KS, Bevins CL, Tomassini N, et al. A novel peptide-producing cell in Xenopus: multinucleated gastric mucosal cell strikingly similar to the granular gland of the skin. J Histochem Cytochem 1992;40: 367-78.