

# *In utero* nicotine exposure promotes M2 activation in neonatal mouse alveolar macrophages

Cherry Wongtrakool<sup>1,2</sup>, Kora Grooms<sup>1</sup>, Xiao-Du Ping<sup>3</sup>, Hilda Rivera<sup>1</sup>, Janine Ward<sup>3</sup>, Susanne Roser-Page<sup>2</sup>, Jesse Roman<sup>4,5</sup>, Lou Ann S. Brown<sup>3</sup> and Theresa W. Gauthier<sup>3</sup>

**BACKGROUND:** Maternal smoking *in utero* has been associated with adverse health outcomes including lower respiratory tract infections in infants and children, but the mechanisms underlying these associations continue to be investigated. We hypothesized that nicotine plays a significant role in mediating the effects of maternal tobacco smoke on the function of the neonatal alveolar macrophage (AM), the resident immune cell in the neonatal lung.

**METHODS:** Primary AMs were isolated at postnatal day 7 from a murine model of *in utero* nicotine exposure. The murine AM cell line MH-S was used for additional *in vitro* studies.

**RESULTS:** *In utero* nicotine increased interleukin-13 and transforming growth factor- $\beta$ 1 (TGF $\beta$ 1) in the neonatal lung. Nicotine-exposed AMs demonstrated increased TGF $\beta$ 1 and increased markers of alternative activation with diminished phagocytic function. However, AMs from mice deficient in the  $\alpha$ 7 nicotinic acetylcholine receptor ( $\alpha$ 7 nAChR) had less TGF $\beta$ 1, reduced alternative activation, and improved phagocytic functioning despite similar *in utero* nicotine exposure.

**CONCLUSION:** *In utero* nicotine exposure, mediated in part via the  $\alpha$ 7 nAChR, may increase the risk of lower respiratory tract infections in neonates by changing the resting state of AM toward alternative activation. These findings have important implications for immune responses in the nicotine-exposed neonatal lung.

Despite the well-known health risks of smoking, the use of cigarettes and tobacco continues to contribute significantly to health-care problems in our society. Twenty-two percent of women of reproductive age smoke cigarettes, exposing nearly 20% of pregnancies to this toxin (1–3). It is clear that *in utero* and postnatal exposure to smoking increases the risk of serious lower respiratory tract infections in infants and children (4). Specifically for the newborn, smoking during pregnancy increases the risks of prematurity, alters immune defenses, and increases risk factors for neonatal sepsis (5,6). Despite advances in neonatal intensive care, infection increases mortality and morbidity, particularly for the premature newborn (7,8).

Nicotine, the fat soluble and addictive component of tobacco, readily crosses the placenta, is present in the amniotic fluid, and adversely affects lung development (9). Levels of nicotine and its metabolites in the fetus closely mirror those observed in maternal plasma (10). *In utero* nicotine exposure alters multiple developing organs including the lung and affects branching morphogenesis (11–13). The effects of nicotine on the developing lung are likely mediated through the  $\alpha$ 7 nicotinic acetylcholine receptor (nAChR), one of a family of cation channels predominantly found in the nervous system (12,14). However, the direct effects of nicotine or the role of the  $\alpha$ 7nAChR on inflammatory-mediated functioning of the resident immune cell of the lung, the alveolar macrophage (AM), are not fully known.

Several pathways govern the activation states of the normally resting AM (15). Classical activation of AM (TH1 response or M1 macrophage) is predominantly characterized by a robust cellular response to eliminate microbes with efficient phagocytosis, respiratory burst, and the generation of pro-inflammatory cytokines. In contrast, alternative activation of AM (TH2 response or M2 macrophage) is currently defined by signature gene expression profiles and response to pathogenic stimuli that have been observed mainly in murine models. Although the definition of alternative activation has been evolving, alternatively activated macrophages are presently characterized by a dampening of the immune response with decreased phagocytosis and respiratory burst. While the classical pathway involves microbial stimuli and efficient clearance of microbes, the alternative pathway can skew the AM response away from microbial clearance to leave the lung more susceptible to infections if the M2 phenotype persists. Another described feature of M2 activation is increased expression and activity of arginase 1, which shunts arginine away from the production of the antimicrobial nitric oxide and promotes remodeling and repair through enhanced collagen deposition (15).

In this study, we hypothesized that exposure to nicotine *in utero* places the neonate at risk for respiratory infections by influencing AM function through alternative activation (M2 activation). Using the mouse macrophage cell line MH-S

<sup>1</sup>Department of Medicine, Emory University School of Medicine, Atlanta, Georgia; <sup>2</sup>Department of Medicine, Atlanta Veterans Affairs Medical Center, Decatur, Georgia; <sup>3</sup>Department of Pediatrics, Emory University School of Medicine, Atlanta, Georgia; <sup>4</sup>Department of Medicine, University of Louisville, Louisville, Kentucky; <sup>5</sup>Department of Medicine, Louisville Veterans Affairs Medical Center, Louisville, Kentucky. Correspondence: Cherry Wongtrakool (cwongtr@emory.edu)

Received 30 September 2011; accepted 30 March 2012; advance online publication 20 June 2012. doi:10.1038/pr.2012.55

and a mouse model of *in utero* nicotine exposure, the goals of this study were (i) to determine whether chronic nicotine exposure polarized the AM toward an M2 phenotype and (ii) whether these changes were mediated via the  $\alpha 7$  nAChR.

## RESULTS

### Nicotine Exposure Increased $\alpha 7$ nAChR on the AM

By western blot techniques, chronic nicotine exposure to the MH-S cell line significantly increased the protein expression of  $\alpha 7$  nAChR 1.3-fold as compared with control cells (Figure 1a). Similarly, primary postnatal day 7 (P7) AMs exposed to *in utero* nicotine demonstrated significant increase in  $\alpha 7$  nAChR immunofluorescence as compared with controls (Figure 1b,c).

### *In Utero* Nicotine Skewed Cytokines in the Neonatal Lung and the AM Toward TH2 via the $\alpha 7$ nAChR

At baseline, neonatal lungs exposed to nicotine demonstrated significantly increased interleukin (IL)-13 and decreased IL1 $\beta$  at the mRNA level at P7 (Figure 2a,b), suggesting that the cytokine milieu of the nicotine-exposed lung was shifted toward TH2. This shift toward TH2 was supported by significantly increased active transforming growth factor- $\beta 1$  (TGF $\beta 1$ ) in the epithelial lining fluid (ELF) of P7 pups exposed to *in utero* nicotine (Figure 3). The  $\alpha 7$  nAChR modulated these nicotine-induced changes in active TGF $\beta 1$ , as nicotine did not induce active TGF $\beta 1$  in the ELF of the  $\alpha 7$  nAChR<sup>-/-</sup> neonatal mice.

The neonatal AM contributed to these nicotine-induced changes in IL-13 and TGF $\beta 1$ . Nicotine *in vitro* significantly increased IL-13 and TGF $\beta 1$  in MH-S cells at the mRNA level (Figure 4a,b). Blockage of the  $\alpha 7$  nAChR with the addition of  $\alpha$ -bungarotoxin ( $\alpha$ BGT) significantly blunted nicotine effects on both IL-13 and TGF $\beta 1$  mRNA. Furthermore, primary P7 AMs demonstrated a significant increase in TGF $\beta 1$  via immunohistochemistry (Figure 4c); this was significantly attenuated in the  $\alpha 7$ -knockout mice despite *in utero* nicotine exposure.


### Nicotine Exposure Induced Alternative Activation in the AM via the $\alpha 7$ nAChR

*In vitro* exposure to nicotine induced markers of alternative activation in the MH-S cell as demonstrated by significant increases in arginase 1, Ym1 (also known as ECF-L or T lymphocyte-derived eosinophil chemotactic factor) and fibronectin (FN) at the mRNA level (Figure 5a-c). Correspondingly, *in utero* nicotine also demonstrated significant increases in these markers of alternative activation (Figure 5d-f). Blockade of the  $\alpha 7$  nAChR with  $\alpha$ BGT blunted *in vitro* nicotine-induced increases in arginase 1, Ym1, and FN on the MH-S cell line. Similarly, nicotine-induced increases in these markers were blocked in the P7  $\alpha 7$  nAChR-deficient AM.


### Nicotine-Exposed AMs Promote FN Expression in a Paracrine Fashion

In addition to increased FN induced by nicotine exposure, AMs treated with nicotine *in vitro* can also promote FN transcription

in adjacent cells. MH-S cells treated with nicotine *in vitro* and cultured on permeable membrane supports separating them from NIH 3T3 fibroblasts permanently transfected with an FN-luciferase reporter were able to induce FN transcription in fibroblasts, as evidenced by increased luciferase production by the transfected NIH 3T3 cells (Figure 6a). These effects were dependent on  $\alpha 7$  nAChR signaling because concurrent incubation with  $\alpha$ BGT abrogated the nicotine effect. Increased FN transcription was absent in fibroblasts cultured without



**Figure 1.** Nicotine induced  $\alpha 7$  nicotinic acetylcholine receptor (nAChR) expression in alveolar macrophages. Cultured murine alveolar macrophages (MH-S cells) were serum-starved overnight and treated with 50  $\mu$ g/ml of nicotine for 24 h. By western blot ( $n = 4$ ), nicotine (N) treatment increases expression of  $\alpha 7$  nAChR in MH-S cells after 24 h as compared with the untreated control (C) (a). Female C57 BL/6J mice were administered 100  $\mu$ g/ml of nicotine in drinking water for 6–8 wk before the timed breeding. Alveolar macrophages (AMs) were harvested from the offspring at postnatal day 7. In AMs harvested from pups exposed to nicotine (bottom color panel, b), expression of  $\alpha 7$  nAChR in relative fluorescent units per cell (RFU/cell) was significantly increased compared to untreated control (top color panel, b) when analyzed with confocal microscopy (c). \* $P < 0.05$  vs. control, four separate litters in each condition were analyzed; image is representative image at 100 $\times$  original magnification. Scale bar represents 5  $\mu$ m.



**Figure 2.** Neonatal lungs exposed to nicotine *in utero* exhibited a shift towards a TH2 profile. Female C57 BL/6J mice were administered 100 µg/ml of nicotine in the drinking water for 6–8 wk before the timed breeding. Whole lung homogenates were collected from the offspring at postnatal day (P)7 for mRNA analysis. Using semi-quantitative bioluminescent reverse-transcriptase PCR, P7 lungs had increased mRNA expression of IL-13 (a) and decreased mRNA expression of interleukin-1β (b). \* $P < 0.05$  vs. control.  $n = 4$ . n-LU denotes luciferase units normalized to endogenous loading control.


MH-S cells in the membrane supports, demonstrating that the nicotine-exposed MH-S cells were capable of promoting FN transcription in fibroblasts. Nicotine-exposed MH-S cells also affected type I collagen expression in fibroblasts (Figure 6b).

#### In Utero Nicotine Impaired AM Phagocytic Function via the $\alpha 7$ nAChR

Last, we evaluated the effects of *in utero* nicotine exposure on the phagocytic function of the primary neonatal AM. *In utero* nicotine exposure significantly impaired the ability of the neonatal AM to phagocytose inactivated *Staphylococcus aureus* by approximately 50% (Figure 7). Despite nicotine exposure, AMs lacking the  $\alpha 7$  nAChR maintained phagocytosis at control levels, suggesting that nicotine-induced deranged phagocytic function was modulated via the  $\alpha 7$  nAChR.

## DISCUSSION


Despite the known risk of cigarette smoke exposure on childhood health, maternal use of cigarettes during pregnancy remains an issue with approximately 10–12% of pregnant women admitting to tobacco use (16). Use of cigarettes during



**Figure 3.**  $\alpha 7$  Nicotinic acetylcholine receptor (nAChR) signaling mediated increased active transforming growth factor- $\beta 1$  (TGF $\beta 1$ ) found in the alveolar epithelial lining fluid (ELF) of postnatal day (P)7 pups exposed to nicotine *in utero*. Alveolar ELF was collected from P7 litters and pooled for measurement of active TGF $\beta 1$  by enzyme-linked immunosorbent assay. *In utero* nicotine exposure induced increased active TGF $\beta 1$  expression in P7 pups. *In utero* nicotine exposure did not induce active TGF $\beta 1$  expression in  $\alpha 7$  nAChR<sup>-/-</sup> pups. \* $P < 0.05$  vs. control, † $P < 0.05$  vs. nicotine. Minimum of four separate litters in each experimental condition.

pregnancy exposes the developing neonatal lung to nicotine. There remains a gap of knowledge regarding the specific effects of nicotine on the characteristics and functioning of the developing AM, the resident immune cell in the lung. This study demonstrated that *in utero* nicotine exposure induced a TH2 milieu in the neonatal lung at baseline and skewed the resting state of the neonatal AM toward M2 activation, impairing phagocytic function. The  $\alpha 7$  nAChR played an important role in modulating the effect of nicotine as demonstrated by the effects of  $\alpha$ BGT *in vitro* and the attenuation of the effect of nicotine *in vivo* in the  $\alpha 7$  nAChR-deficient neonatal mouse.

The nAChRs are found in a number of cell types and non-neuronal organs, including the lung. Within the lung, nAChRs are present in epithelial cells, fibroblasts, smooth muscle cells, and AMs (12,14). We have previously demonstrated that nicotine increased branching morphogenesis of the developing lung in the pseudoglandular stage through  $\alpha 7$  nAChR-mediated signals (12). Others have shown that nicotine induced  $\alpha 7$  nAChR expression in fetal lung, particularly around the airways (14).  $\alpha 7$  nAChRs have been found in peripheral blood monocytes and macrophages, and they are capable of influencing immune responses (17). An anti-inflammatory role for the  $\alpha 7$  nAChR has been noted, as a specific  $\alpha 7$  nAChR agonist decreased tumor necrosis factor- $\alpha$  release in the lung, and  $\alpha 7$  nAChR agonists, including nicotine, also inhibited nuclear factor- $\kappa B$  activity and lipopolysaccharide-induced release of tumor necrosis factor- $\alpha$  in the lung (18). However, the literature also demonstrates that  $\alpha 7$  nAChR has been associated with pro-inflammatory effects. Prostaglandins and cyclo-oxygenase 2 expression were enhanced by activation of  $\alpha 7$  nAChR, whereas nicotine induced nuclear factor- $\kappa B$  and inducible nitric oxide synthase in peritoneal macrophages in a model of atherosclerosis (19,20). Expression of nAChRs was upregulated by



**Figure 4.** Nicotine-induced increases in mRNA expression of transforming growth factor- $\beta$ 1 (TGF $\beta$ 1) and interleukin (IL)-13 were dependent on  $\alpha$ 7 nicotinic acetylcholine receptor (nAChR)-mediated signals. MH-S cells were serum-starved overnight and treated with 50  $\mu$ g/ml of nicotine (N) after 1 h pretreatment with 5 nM of  $\alpha$ -bungarotoxin ( $\alpha$ BGT), a snake venom toxin that is an inhibitor of  $\alpha$ 7 nAChR. Cells were harvested after 24 h for mRNA analysis using quantitative real-time PCR. Nicotine induced mRNA expression of TGF $\beta$ 1 (a) and IL-13 (b) and pretreatment with  $\alpha$ BGT blocked induction of TGF $\beta$ 1 (a) and IL-13 (b) by nicotine. Primary alveolar macrophages (AMs) were harvested from the offspring at postnatal day 7. Immunofluorescent staining for TGF $\beta$ 1 was significantly increased after *in utero* nicotine exposure in primary AMs, whereas TGF $\beta$ 1 was not increased in  $\alpha$ 7 nAChR $^{-/-}$  pups despite nicotine exposure (c). \* $P < 0.05$  vs. control, † $P < 0.05$  vs. nicotine. Three to four separate litters in each experimental condition.

tumor necrosis factor- $\alpha$ , supporting a potential link between inflammation and nAChRs (21). For the developing AM, our study suggested that nAChR activation via *in utero* nicotine shifted the baseline cytokine milieu toward TH2, increased TGF $\beta$ 1, and skewed the resting AM state toward M2.

Currently available knowledge suggests the skewed responses of alternatively activated macrophages may leave the lung more susceptible to infections and remodeling. As a professional phagocyte within the lung, the AM patrols the lung, defending it against foreign particles and infection by

initiating immune responses, participating in phagocytosis and particle clearance, and orchestrating subsequent inflammatory processes (22). Studies have demonstrated that complete absence of macrophages led to dramatically increased mortality after bacterial infection (23). Alternatively activated macrophages have also been shown to promote fibroblast proliferation as well as production of collagen and FN, extracellular matrix proteins important in remodeling (24). Our study shows that nicotine-induced alternative activation of the AM leads to increased FN within the alveolar macrophage and induction of FN transcription in fibroblasts in the setting of increased TGF $\beta$ 1, a well-known stimulus for FN expression, potentially to promote airway remodeling (25). These effects on airway remodeling in the neonatal lung require additional investigation.

The nicotine-exposed neonatal lung and AM were hallmarked by increased TGF $\beta$ 1, a well described anti-inflammatory mediator in monocytic cells and AMs (26,27). Furthermore, TGF $\beta$ 1 contributes to the development of chronic lung disease (bronchopulmonary dysplasia) in the premature lung (28,29). In a neonatal hyperoxia model, neutralization of TGF $\beta$ 1 improved alveologenesis and microvascular development (30). Taken together, our data suggest that not only are the AMs functionally impaired with *in utero* nicotine exposure, but also the lung parenchyma and vasculature are at greater risk for injury in the setting of increased TGF $\beta$ 1.


Many studies focus on the effects of cigarette smoke exposure on the lung, but the complex nature of cigarette smoke can potentially impede full mechanistic understanding of pathophysiology. In other examinations of adult human AMs, adult human AMs exposed to cigarette smoke similarly exhibit signs of M2 activation (31,32). Taken together with previously published literature, our findings support the possible role of nicotine as a significant player in the pathophysiology behind the adverse effects of prenatal tobacco smoke exposure on the fetus because the effects from nicotine exposure alone mimic those seen with tobacco smoke exposure. Although cessation of cigarette smoking is advocated to all pregnant women, our results suggest that nicotine replacement therapy as a strategy for smoking cessation therapy may not be desirable in this population and alternative smoking cessation therapies may need to be considered.

## MATERIALS AND METHODS

All animal and experimental protocols were approved by the Emory University Institutional Animal Care and Use Committee and the Office of Biosafety.

### Model of *In Utero* Nicotine Exposure and Timed Breeding

Female C57BL6/J mice and *Chrna7* $^{-/-}$  mice ( $\alpha$ 7 nAChR $^{-/-}$ ,  $\alpha$ 7 knockout in C57BL6/J background; Jackson Laboratories, Bar Harbor, ME) were or were not administered nicotine (100  $\mu$ g/ml) in the drinking water ad libitum for 6–8 wk before timed breeding. Pregnant female mice continued to drink nicotine-treated water throughout the pregnancy. In this model, pregnant female mice are able to establish a steady state plasma level of nicotine similar to that seen in heavy smokers (12,33). P0 was determined by the day of birth. Pups were evaluated at P7. A total of 100% of the pups in the nicotine group remained exposed to nicotine through maternal breast milk until P7.



**Figure 5.** Nicotine promoted M2 activation of alveolar macrophages (AMs) *in vitro* and *in vivo*. Primary AMs were harvested at postnatal day (P)7. In addition, MH-S cells were serum-starved overnight and treated with 50 μg/ml of nicotine after 1 h pretreatment with 5 nM of α-bungarotoxin (αBGT). Cells were harvested after 24 h for mRNA analysis using quantitative real-time PCR. Nicotine induced mRNA expression of the following markers of M2 activation: arginase 1 (Arg1), Ym1, and fibronectin (a–c, respectively). Similarly, increased expressions of Arg1, Ym1, and fibronectin were seen by immunofluorescent staining in relative fluorescent units per cell (RFU/cell) using confocal microscopy in primary AMs at P7 (d–f, respectively). Pre-treatment with αBGT blocked in the nicotine-induced changes in mRNA expression. *In utero* nicotine exposure did not induce changes in mRNA expression of Arg1, Ym1, or fibronectin in primary AMs from α7 nAChR<sup>-/-</sup> pups. \*P < 0.05 vs. control, \*\*P < 0.01 vs. control, †P < 0.05 vs. nicotine, ‡P < 0.01 vs. nicotine. Figure is a composite of three to five separate litters in each experimental condition.

**In Vitro Nicotine Exposure**

The MH-S murine AM cell line (ATCC, Manassas, VA) was cultured in Roswell Park Memorial Institute media containing 10% fetal bovine serum, 100 U/ml penicillin, 100 μg/ml streptomycin, and 0.25 μg/ml amphotericin B. Cells were or were not exposed to ± nicotine (50 μg/ml) for 24 h. In some experiments, the cells were incubated with αBGT (5 nM), a selective inhibitor of α7 nAChR (34).

**Recovery of Primary AM and ELF**

AMs were isolated from P7 neonatal pups as previously described (27).

**Measurement of ELF Cytokine Levels**

Pooled pup ELF was evaluated for IL-13 and active TGFβ1 via commercial enzyme-linked immunosorbent assays per the manufacturers’ instructions (IL-13, R&D Systems, Minneapolis, MN; Active TGFβ1, Promega, Madison, WI). Values were normalized to protein as determined by the bicinchoninic acid protein assay (Bio-Rad,


Hercules, CA). Data are presented as the mean (IL-13 or TGF-β1 in pg/μg protein) ± SEM (27).

**Measurement of α7 nAChR**

Levels of AM α7 nAChR were measured by western blot analysis as previously described (35,36). Primary P7 mouse AMs obtained from pooled bronchoalveolar lavages were evaluated via immunofluorescence for α7 nAChR and markers of M2 activation using methods previously described (27).

**RNA Extraction, and Semi-Quantitative Bioluminescent and Quantitative Real-Time Reverse-Transcriptase PCR**

The determination of mRNA levels was done by a semi-quantitative bioluminescence-based reverse-transcriptase PCR assay and quantitative real-time reverse-transcriptase PCR as previously described (12). The primers used were synthesized on the basis of GenBank published sequences: IL-13: 5’ GGAGCTGAGCAACATCACACA



**Figure 6.** Nicotine promotes expression of extracellular matrix proteins associated with alternative activation. MH-S cells were serum-starved overnight, pretreated with 5 nM of bungarotoxin (BGT) to inhibit  $\alpha 7$  nicotinic acetylcholine receptor, then treated with 50  $\mu$ g/ml of nicotine and co-cultured with fibroblasts permanently transfected with a fibronectin-luciferase reporter for 24 h. MH-S cells treated with nicotine promoted transcription of fibronectin in co-cultured fibroblasts as measured by luciferase assay (a). Pretreatment with  $\alpha$ BGT blocked the nicotine-induced increase in fibronectin transcription. Collagen type I expression was also analyzed in the MH-S cells co-cultured with fibroblasts in (a). Nicotine-treated MH-S cells (N, black bars) also had increased collagen type I expression by western blot analysis after 24 h as compared with untreated MH-S cells (C, white bars) (b). Image is a representative western blot. \* $P < 0.05$  vs. control, † $P < 0.05$  vs. nicotine.  $n = 3$ .


3' and 5' GGTCCTGTAGATGGCATTGCA 3'; TGF $\beta$ 1: 5' CCCACT CCCGTGGCTTC 3' and 5' TAGTAGTCCGCTTCGGGCT 3'; IL1- $\beta$ : 5' GAGCACCTTCTTTTCC 3' and 5' GGAAAAAGAAGGTGGTC 3'; Ym1/2: 5' TTATCCTGAGTGACCCTTCTAAG 3' and 5' TCATTA CCCAGATAGGCATAGG 3'; arginase 1: 5' TGGACCTGGCCTTTG TTGA 3' and 5' GGTTGTGACGGGAGTGTT 3'; FN: 5' CTGTG ACAACTGCCGTAG 3' and 5'ACCAAGGTCAATCCACAC 3'.

**FN Luciferase Assay**

MH-S cells were cultured on a permeable membrane Transwell (Corning, Tewksbury, MA) support with NIH 3T3 fibroblasts permanently transfected with the FN promoter attached to a luciferase reporter (FN-luciferase) in the bottom well. After overnight serum starvation, MH-S cells on the permeable membrane were then treated with 5 nM  $\alpha$ BGT for 1 h before the addition of 50  $\mu$ g/ml of nicotine. After 24 h, the MH-S cells and membrane supports were removed and FN transcription in the transfected 3T3 fibroblasts was measured by luciferase activity. Relative luciferase units were recorded and are presented as mean  $\pm$  SD (37).

**Measurement of Phagocytosis**

Phagocytosis was measured in freshly isolated P7 AMs as previously described (27,38).



**Figure 7.** *In utero* nicotine exposure impaired the phagocytic response of alveolar macrophages (AMs), and nicotine's effect was dependent on  $\alpha 7$  nicotinic acetylcholine receptor (nAChR)-mediated signals. Primary AMs were harvested at postnatal day 7 and were administered pH rhodo-labeled *Staphylococcus aureus*, which fluoresces when incorporated into an acidic phagolysosome. AMs from *in utero* nicotine-exposed pups demonstrated diminished phagocytosis, but AMs from  $\alpha 7$  nAChR $^{-/-}$  pups maintained their phagocytic ability despite *in utero* nicotine exposure. \* $P < 0.05$  vs. control, † $P < 0.05$  vs. nicotine. Minimum of four separate litters in each experimental condition.

**Statistical Analyses**

Sigma Stat for Windows (Systat Software, San Jose, CA) was used for all statistical analyses. ANOVA or ANOVA on Ranks was used as appropriate. Student Newman Keul's or Dunn's test was used for multiple comparisons. A  $P \leq 0.05$  was considered significant. Each  $n$  represents a separate mouse litter or a separate experimental condition.

**STATEMENT OF FINANCIAL SUPPORT**

This work was funded by National Institutes of Health grant K08 HL080293 (to C.W.) and a grant from the Children's Center for Developmental Lung Biology (to T.W.G. and L.A.S.B.).

**REFERENCES**

- Centers for Disease Control and Prevention. Cigarette smoking among adults—United States, 2007. *MMWR Morb Mortal Wkly Rep* 2008;57:1221–6.
- D'Angelo D, Williams L, Morrow B, et al. Preconception and interconception health status of women who recently gave birth to a live-born infant—Pregnancy Risk Assessment Monitoring System (PRAMS), United States, 26 reporting areas, 2004. *MMWR Surveill Summ* 2007;56:1–35.
- Tong VT, Jones JR, Dietz PM, D'Angelo D, Bombard JM. Trends in smoking before, during, and after pregnancy - Pregnancy Risk Assessment Monitoring System (PRAMS), United States, 31 sites, 2000-2005. *MMWR Surveill Summ* 2009;58:1–29.
- DiFranza JR, Aligne CA, Weitzman M. Prenatal and postnatal environmental tobacco smoke exposure and children's health. *Pediatrics* 2004;113(4 Suppl):1007–15.
- Mercelina-Roumans PE, Breukers RB, Ubachs JM, van Wersch JW. Hematological variables in cord blood of neonates of smoking and nonsmoking mothers. *J Clin Epidemiol* 1996;49:449–54.
- Suzuki K, Tanaka T, Kondo N, Minai J, Sato M, Yamagata Z. Is maternal smoking during early pregnancy a risk factor for all low birth weight infants? *J Epidemiol* 2008;18:89–96.
- Alarcon A, Peña P, Salas S, Sancha M, Omeñaca F. Neonatal early onset *Escherichia coli* sepsis: trends in incidence and antimicrobial resistance in the era of intrapartum antimicrobial prophylaxis. *Pediatr Infect Dis J* 2004;23:295–9.

8. Cordero L, Rau R, Taylor D, Ayers LW. Enteric gram-negative bacilli bloodstream infections: 17 years' experience in a neonatal intensive care unit. *Am J Infect Control* 2004;32:189–95.
9. Rehan VK, Wang Y, Sugano S, et al. In utero nicotine exposure alters fetal rat lung alveolar type II cell proliferation, differentiation, and metabolism. *Am J Physiol Lung Cell Mol Physiol* 2007;292:L323–33.
10. Luck W, Nau H, Hansen R, Steldinger R. Extent of nicotine and cotinine transfer to the human fetus, placenta and amniotic fluid of smoking mothers. *Dev Pharmacol Ther* 1985;8:384–95.
11. Rehan VK, Asotra K, Torday JS. The effects of smoking on the developing lung: insights from a biologic model for lung development, homeostasis, and repair. *Lung* 2009;187:281–9.
12. Wongtrakool C, Roser-Page S, Rivera HN, Roman J. Nicotine alters lung branching morphogenesis through the alpha7 nicotinic acetylcholine receptor. *Am J Physiol Lung Cell Mol Physiol* 2007;293:L611–8.
13. Zhao Z, Reece EA. Nicotine-induced embryonic malformations mediated by apoptosis from increasing intracellular calcium and oxidative stress. *Birth Defects Res B Dev Reprod Toxicol* 2005;74:383–91.
14. Sekhon HS, Jia Y, Raab R, et al. Prenatal nicotine increases pulmonary alpha7 nicotinic receptor expression and alters fetal lung development in monkeys. *J Clin Invest* 1999;103:637–47.
15. Martinez FO, Helming L, Gordon S. Alternative activation of macrophages: an immunologic functional perspective. *Annu Rev Immunol* 2009;27:451–83.
16. Tong VT, Jones JR, Dietz PM, D'Angelo D, Bombard JM. Trends in smoking before, during, and after pregnancy - Pregnancy Risk Assessment Monitoring System (PRAMS), United States, 31 sites, 2000–2005. *MMWR Surveill Summ* 2009;58:1–29.
17. Pavlov VA, Wang H, Czura CJ, Friedman SG, Tracey KJ. The cholinergic anti-inflammatory pathway: a missing link in neuroimmunomodulation. *Mol Med* 2003;9:125–34.
18. Su X, Lee JW, Matthay ZA, et al. Activation of the alpha7 nAChR reduces acid-induced acute lung injury in mice and rats. *Am J Respir Cell Mol Biol* 2007;37:186–92.
19. De Simone R, Ajmone-Cat MA, Carnevale D, Minghetti L. Activation of alpha7 nicotinic acetylcholine receptor by nicotine selectively up-regulates cyclooxygenase-2 and prostaglandin E2 in rat microglial cultures. *J Neuroinflammation* 2005;2:4.
20. Lau PP, Li L, Merched AJ, Zhang AL, Ko KW, Chan L. Nicotine induces proinflammatory responses in macrophages and the aorta leading to acceleration of atherosclerosis in low-density lipoprotein receptor(-/-) mice. *Arterioscler Thromb Vasc Biol* 2006;26:143–9.
21. Gahring LC, Osborne-Hereford AV, Vasquez-Opazo GA, Rogers SW. Tumor necrosis factor alpha enhances nicotinic receptor up-regulation via a p38MAPK-dependent pathway. *J Biol Chem* 2008;283:693–9.
22. Gordon SB, Read RC. Macrophage defences against respiratory tract infections. *Br Med Bull* 2002;61:45–61.
23. Broug-Holub E, Toews GB, van Iwaarden JF, et al. Alveolar macrophages are required for protective pulmonary defenses in murine *Klebsiella pneumoniae*: elimination of alveolar macrophages increases neutrophil recruitment but decreases bacterial clearance and survival. *Infect Immun* 1997;65:1139–46.
24. Gratchev A, Guillot P, Hakiy N, et al. Alternatively activated macrophages differentially express fibronectin and its splice variants and the extracellular matrix protein betaG-H3. *Scand J Immunol* 2001;53:386–92.
25. Ramirez AM, Wongtrakool C, Welch T, Steinmeyer A, Zügel U, Roman J. Vitamin D inhibition of pro-fibrotic effects of transforming growth factor beta1 in lung fibroblasts and epithelial cells. *J Steroid Biochem Mol Biol* 2010;118:142–50.
26. Fadok VA, Bratton DL, Konowal A, Freed PW, Westcott JY, Henson PM. Macrophages that have ingested apoptotic cells *in vitro* inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF-beta, PGE2, and PAF. *J Clin Invest* 1998;101:890–8.
27. Gauthier TW, Ping XD, Gabelaia L, Brown LA. Delayed neonatal lung macrophage differentiation in a mouse model of in utero ethanol exposure. *Am J Physiol Lung Cell Mol Physiol* 2010;299:L8–16.
28. Gauldie J, Galt T, Bonniaud P, Robbins C, Kelly M, Warburton D. Transfer of the active form of transforming growth factor-beta 1 gene to newborn rat lung induces changes consistent with bronchopulmonary dysplasia. *Am J Pathol* 2003;163:2575–84.
29. Vicencio AG, Lee CG, Cho SJ, et al. Conditional overexpression of bioactive transforming growth factor-beta1 in neonatal mouse lung: a new model for bronchopulmonary dysplasia? *Am J Respir Cell Mol Biol* 2004;31:650–6.
30. Nakanishi H, Sugiura T, Streisand JB, Lonning SM, Roberts JD Jr. TGF-beta-neutralizing antibodies improve pulmonary alveologenesis and vasculogenesis in the injured newborn lung. *Am J Physiol Lung Cell Mol Physiol* 2007;293:L151–61.
31. Hodge S, Matthews G, Mukaro V, et al. Cigarette smoke-induced changes to alveolar macrophage phenotype and function are improved by treatment with procysteine. *Am J Respir Cell Mol Biol* 2011;44:673–81.
32. Shaykhiiev R, Krause A, Salit J, et al. Smoking-dependent reprogramming of alveolar macrophage polarization: implication for pathogenesis of chronic obstructive pulmonary disease. *J Immunol* 2009;183:2867–83.
33. Rowell PP, Hurst HE, Marlowe C, Bennett BD. Oral administration of nicotine: its uptake and distribution after chronic administration to mice. *J Pharmacol Methods* 1983;9:249–61.
34. Orr-Urtreger A, Göldner FM, Saeki M, et al. Mice deficient in the alpha7 neuronal nicotinic acetylcholine receptor lack alpha-bungarotoxin binding sites and hippocampal fast nicotinic currents. *J Neurosci* 1997;17:9165–71.
35. Roman J, Ritzenthaler JD, Gil-Acosta A, Rivera HN, Roser-Page S. Nicotine and fibronectin expression in lung fibroblasts: implications for tobacco-related lung tissue remodeling. *FASEB J* 2004;18:1436–8.
36. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248–54.
37. Tomic R, Lassiter CC, Ritzenthaler JD, Rivera HN, Roman J. Anti-tissue remodeling effects of corticosteroids: fluticasone propionate inhibits fibronectin expression in fibroblasts. *Chest* 2005;127:257–65.
38. Gauthier TW, Ping XD, Harris FL, Wong M, Elbahesh H, Brown LA. Fetal alcohol exposure impairs alveolar macrophage function via decreased glutathione availability. *Pediatr Res* 2005;57:76–81.