doi:10.1038/pr.2014.210

Articular cartilage and joint disease

A better understanding of the molecular mechanisms that regulate the growth of articular cartilage may give insight into the antecedents of joint disease. Using laser-capture microdissection, Lui and colleagues isolated chondrocytes from growing tibial cartilage in one-week-old mice. Expression patterns obtained by microarray identified novel signaling pathways that are spatially regulated. [See page 406](#)

Intermittent hypoxia and adult cardiovascular dysfunction

Eli Revzin/Henara/Thinkstock

Chu and coinvestigators hypothesized that postnatal exposure to intermittent hypoxia (IH) leads to cardiovascular dysfunction in adulthood.

Newborn mice exposed to IH for four weeks exhibited higher systolic blood pressure, impaired baroreflex responses, and decreased heart rate variability at three months. Exposure to postnatal IH might alter normal development of the renin-angiotensin system and lead to cardiovascular dysfunction in adult mice. [See page 425](#)

Adipose tissue and obesity comorbidities

It is unknown how excess adipose tissue (AT) contributes to comorbidities of obesity. Ferrante and coauthors quantified and characterized exosomes shed by adipocytes from obese and lean preadolescents and teenagers. Their findings suggest

that lean visceral AT sheds a higher concentration of exosomes than obese visceral AT. [See page 447](#)

Diagnosing venous thromboembolism

Emw

Venous thromboembolism (VTE) is difficult to diagnose in children. Kerlin *et al.* hypothesized that a child-specific clinical probability tool (CPT) to estimate the likelihood of VTE is feasible. Radiology reports were used to identify 389 children imaged for suspected VTE. Relevant signs, symptoms, and comorbidity variables were extracted from the corresponding medical records. The results suggest that the CPT is feasible in children and could facilitate early diagnosis of VTE. [See page 463](#)

Genes and bronchopulmonary dysplasia

Sampath and colleagues hypothesized that genetic variants of antioxidant

Krishna Kumar/Henara/Thinkstock

response elements (AREs) modulate susceptibility to or severity of bronchopulmonary dysplasia (BPD) in very-low-birth-weight (VLBW) infants. Blood samples from 659 VLBW infants were genotyped for ARE variants. The presence of the hypomorphic *NQO1* single-nucleotide polymorphism (SNP) in a homozygous state was associated with an increased risk of severe BPD whereas the presence of the *NFE2L2* SNP was associated with a decreased risk. [See page 477](#)

Prenatal LC-PUFA

annie-claude/© iStockphoto/Thinkstock

It is unclear whether maternal LC-PUFA status during pregnancy influences behavioral characteristics in later childhood. Steenweg-de Graaff and colleagues measured maternal plasma concentrations of docosahexaenoic acid (DHA), eicosapentaenoic acid, and arachidonic acid (AA) in mid-pregnancy. Higher maternal DHA and n-3:n-6 ratio seem to be associated with fewer child behavioral difficulties at six years of age, and higher AA appears to be associated with more of such problems. [See page 489](#)