Open

Does prenatal stress alter the developing connectome?

Dustin Scheinost¹, Rajita Sinha^{2,3,4}, Sarah N. Cross⁵, Soo Hyun Kwon⁶, Gordon Sze¹, R. Todd Constable^{7,8} and Laura R. Ment^{9,10}

Human neurodevelopment requires the organization of neural elements into complex structural and functional networks called the connectome. Emerging data suggest that prenatal exposure to maternal stress plays a role in the wiring, or miswiring, of the developing connectome. Stress-related symptoms are common in women during pregnancy and are risk factors for neurobehavioral disorders ranging from autism spectrum disorder, attention deficit hyperactivity disorder, and addiction, to major depression and schizophrenia. This review focuses on structural and functional connectivity imaging to assess the impact of changes in women's stress-based physiology on the dynamic development of the human connectome in the fetal brain.

Human neurodevelopment requires the organization of neural elements into complex structural and functional networks called the connectome (1–3). While development of the connectome is contingent on many factors, emerging data suggest that prenatal exposure to maternal stress may also play a role (4–7). Stress is a signal in response to challenging and uncontrollable adverse events and perceived threat (8,9), and exposure to early life stress is a risk factor for neurobehavioral disorders ranging from autism spectrum disorder (ASD), attention deficit hyperactivity disorder and addiction, to depression and schizophrenia (10–21). Both high stress and stress-related conditions, including depression and anxiety, potentially stimulate biological stress pathways (7), alter synaptogenesis (22,23), and change brain development (5,24–28). The prenatal period is critical for brain development, and prenatal stressors exhibit long-lasting influence on adult disorders, making stressor type and timing important factors to explore (27,29,30). Fetal sex and genetic variants may also mediate stress responsiveness (6,7,31–34).

Recent reports suggest that prenatal stress exposure (PNSE) is a global public health problem (13,29,35,36). PNSE has been reported in 10–35% of children worldwide (37). Nearly 8–23% of infants in the United States, or almost 800,000 neonates/year, experience prenatal exposure to depression (38,39), and reports from developing countries support similar numbers

(40–42). Likewise, 1 in 7 to 1 in 13 pregnant women in the United States affirm symptoms of anxiety, while 5.6–14.8% in developing countries suffer a similar diagnosis (40–44). Since a nationally representative study found that more than half of the pregnant women (65.9%) experiencing depression in the United States went undiagnosed (45), these data may underrepresent the problem.

PNSE is believed to both activate the hypothalamic-pituitary-adrenal (HPA) axis and result in epigenetic changes in the developing brain. This review will focus on converging preclinical and clinical imaging data to assess the impact of these changes in women's stress-based physiology on the functional development of the human fetal brain. Prior to reviewing published data, we review common causes of PNSE and methods for measuring the structural and functional connectome. We also provide preliminary human data demonstrating increasing connectivity in limbic system structures across the third trimester of gestation.

STRESS MODELS IN CLINICAL AND TRANSLATIONAL STUDIES

While the relationship between maternal psychosocial stress and adverse pregnancy outcomes has been shown in many studies, it is important to define the nature of the stressor and the subject population (46). Stressors range from depression and anxiety to natural disasters, bereavement and steroid administration. As stressors vary, so may the outcomes. For a listing of outcomes, putative prenatal stressors and representative publications, please see [Table 1](#).

Depression and Anxiety

Although some older studies relied on retrospective recall measures and few evaluated the effects of increasing duration and strength of psychosocial stressors, more recent investigations have employed depression and anxiety as markers of maternal stress. Estimates suggest that 8–23% of women have symptoms of depression during their pregnancy (47). Likewise, 7.7–14% report anxiety, and there are numerous reports of coexisting depression and anxiety in the same pregnant woman at any given time. While depression and anxiety are common proxies

¹Department of Radiology and Biomedical Imaging, Yale School of Medicine, New Haven, Connecticut; ²Department of Psychiatry, Yale School of Medicine, New Haven, Connecticut; ³Department of Child Study, Yale School of Medicine, New Haven, Connecticut; ⁴Department of Neuroscience, Yale School of Medicine, New Haven, Connecticut; ⁵Department of Obstetrics, Gynecology, and Reproductive Sciences, Yale School of Medicine, New Haven, Connecticut; ⁶Department of Pediatrics, Yale School of Medicine, New Haven, Connecticut; ⁷Department of Radiology and Biomedical Imaging, Yale School of Medicine, New Haven, Connecticut; ⁸Department of Neurosurgery, Yale School of Medicine, New Haven, Connecticut; ⁹Department of Pediatrics, Yale School of Medicine, New Haven, Connecticut; ¹⁰Department of Neurology, Yale School of Medicine, New Haven, Connecticut. Correspondence: Laura R. Ment (laura.ment@yale.edu)

Received 18 May 2016; accepted 30 August 2016; advance online publication 2 November 2016. doi:10.1038/pr.2016.197

Table 1. Disorders and putative prenatal stressors

Disorder	Prenatal stress and representative references
Autism spectrum disorder	Anxiety (13,103) Conjugal conflict (104) Depression (105,106) Maternal bereavement (55) Natural disasters (13,52)
Attention deficit hyperactivity disorder	Anxiety (61,103,107,108) Maternal bereavement (15,55)
Bipolar affective disorder	Stress (109)
Cognition	Anxiety (107,110) Depression (48) Natural disaster (111) Psychosocial stress (60,112)
Depression	Depression (16,48,113) PTSD (113)
Internalizing problems	Depression (114)
Neonatal behavioral changes	Anxiety (115,116) Depression (48,115,117) Natural disasters (49,51) Perceived stress scale (118,119) PTSD (51)
Pervasive developmental disorder	Depression (53)
Psychosis	Cumulative life experiences (120,121) Depression (121)
Schizophrenia	PTSD (122,123)

PTSD, post-traumatic stress disorder.

for stress, stress in pregnant women does not always coincide with elevated depression or anxiety scales. As such, cases of PNSE may be missed in such analysis. Finally, depression and anxiety may have independent or additive effects in regards to PNSE, making it difficult to fully disentangle these effects with this model. For a more complete review of this topic, please see Suri *et al.* (48).

Natural Disasters

Another approach to test the hypothesis that PNSE results in neurobehavioral disorders is the use of natural disasters as “experiments of nature.” Unlike depression and anxiety, natural disasters are independent of the subject’s genetic background, personality or other confounding characteristics. Disasters strike in a random manner, similar to a randomized controlled experiment, and thus can provide data on prenatal stressors to which a given cohort of pregnant women were exposed (13). Using this strategy, the impact of disasters ranging from hurricanes to terrorist attacks on neurobehavioral outcomes of the offspring have been assessed (13,49–52).

Preconception Stress

In contrast, the influence of preconception adversity and the impact of high cumulative stress on maternal perception of prenatal stress on the developing connectome are just beginning to be explored (53–55). Consistent with preclinical studies showing effects of repeated stress on neural atrophy and neurobehavioral effects (56), human studies link altered structure and function of limbic, subcortical, and frontal regions to higher levels of cumulative stress (28,57,58). These data suggest that preconception adversity may shape perception and control of prenatal stress levels and should be considered in investigations of PNSE on neurobehavioral and MRI outcomes.

Prenatal Maternal Stress

PNSE has been widely associated with preterm birth, intra-uterine growth restriction, and reduced fetal head growth (50,51,59–61). In addition, several studies have reported that increased acute maternal stress is associated with changes in fetal heart rate, activity level, sleep patterns, and higher pulsatility indices in the middle cerebral artery (21,60). PNSE can also be directly measured using prospective data collections in samples of pregnant women with questionnaires, clinical interviews, and biological samples such as cortisol from maternal saliva, blood, or amniotic fluid.

METHODS TO ASSESS CONNECTIVITY USING MRI

Advances in neuroimaging provide important information about microstructural and functional connectivity (62), and offer opportunities to understand the impact of PNSE on the developing connectome (1–3). In the following section, we define measures commonly used in connectomics with examples shown in [Figure 1](#).

Functional connectivity provides information about neural regions that are physiologically functionally coupled, independent of structural connectivity (63). Based on the blood oxygen level dependent signal and derived from time series observations, it assesses “temporal correlations between spatially remote neurophysiological events.” (63) High correlation between time courses of two regions or voxels implies high functional connectivity. For the references included in this review, functional MRI (fMRI) data are largely collected in the resting state, or resting state-fMRI (rs-fMRI).

Methods to assess rs-fMRI data (64,65) include seed, independent components analysis, and voxel-wise connectivity. Seed-based connectivity is most frequently used in human studies and involves (i) selecting a predefined region of interest (ROI), (ii) extracting the average time course from this ROI, and (iii) correlating this average time course with the time courses of every other voxel in the gray matter. Independent components analysis is mathematical modeling technique that parcellates the brain into independent spatial components or networks. These networks can be compared across subject groups or used for later analysis. Voxel-wise connectivity methods are generalizations of seed-based connectivity where many seed connectivity analyses are performed treating each voxel in the gray matter as a unique ROI. As these methods

Figure 1. Examples of functional and structural connectivity. (a) Functional connectivity. Functional connectivity measures the synchrony or correlation of brain activity between two or more regions of the brain. Common methods include seed connectivity, independent components analysis, and voxel-wise connectivity. Seed-based connectivity measures functional connectivity from a predefined region of interest (ROI, or seed, shown in green) and the rest of the gray matter. Regions of positive or negative functional connectivity are shown as red and blue regions. Independent components analysis is mathematical modeling technique that parcellates the brain into independent spatial components or networks. Example components shown are the motor network and the default mode network. Voxel-wise connectivity methods involve correlating the time course of every voxel in the gray matter with the time course of every other voxel in the gray matter. Connectivity for each voxel are often summarized to a single number using network theory to highlight so-called hub regions in the brain. (b) Structural connectivity. Structural connectivity measures anatomical white matter connections linking different cortical and sub cortical regions. Common methods include ROI quantification, tract-based and tractography, and voxel-based morphometry. For ROI quantification, average FA across all voxels in *a priori* ROIs (shown in white outline and overlay) is compared across study groups. Tractography is modeling technique used to identifying white matter tracts used in further analyses. In VBM analysis, FA data from all subjects is transformed into a common space and comparison across each voxel of the white matter is performed. (Figure modified with permission, John Wiley & Sons, Hoboken, NJ)

produce a large amount of data (approximately 20,000 seed-connectivity results), seed-connectivity results for each voxel are often summarized to a single number using network theory.

Anatomical connections in the developing brain represent *microstructural connectivity* (66). Diffusion-weighted imaging (dMRI) assesses the diffusion of water along axons and permits visualization of axonal pathways. By modeling the directional diffusion of water as an ellipsoidal shape, or “tensor”, at each voxel in the brain, dMRI permits assessment of white matter tracts. The first eigenvector, λ_1 , describes the direction

of maximal diffusion, while the second and third define diffusivity perpendicular to this principle axis. Radial diffusivity represents the average of λ_2 and λ_3 and is affected by changes in axon caliber and myelination. Fractional anisotropy (FA) measures the degree to which water diffuses in one direction (along the axon) by computing the ratio of λ_1 to λ_2 and λ_3 and is the most common measure used to assess axonal integrity. High values of FA suggest more highly organized, strongly myelinated tracts.

The three main approaches to analyzing dMRI data include region ROI quantification, tract-based analysis and tractography, and voxel-based morphometry (VBM). ROI quantification is frequently used in human studies investigating the impact of PNSE on the developing connectome. In this method, one or more ROIs are selected *a priori* and the average FA across all voxels in the ROI calculated. Typically, ROIs are major white matter tracts. Tractography is modeling technique used to identify these tracts. Once identified, they can be analyzed using graph theory or ROI analyses. In VBM analysis, FA data from all subjects are transformed into a common space and compared across each voxel of the white matter.

Finally, although not direct measures of the connectome, we include studies assessing *brain morphometry*, including cortical volumes and thickness. Morphological features of different brain regions are not independent of those of other areas, and the brain shows a high level of coordination between different structures (67). This coordination of morphological features is often referred to as anatomical covariance (67–69) and resembles functional and structural connectivity.

PRECLINICAL DATA SUPPORT THE IMPACT OF PNSE ON DEVELOPING CONNECTOME

Across multiple species and numerous time points, converging data suggest that gestational stress influences brain development. Similar to the human subjects, the offspring of numerous species exposed to PNSE demonstrate increases in anxiety and depression, impaired spatial memory and alterations in cognition (70,71). Systematic experimental investigations using standardized animal models and outcome measures (6,72,73) address not only the impact of PNSE on maternal endocrine functions and the “re-programming” of the fetal HPA axis (74–79), but also suggest that changes in corticogenesis contribute to the long-lasting effects on brain and behavior (Table 2) (74,80,81).

MRI STUDIES OF PNSE AND THE DEVELOPING BRAIN

While the neural correlates of acute and cumulative post-natal stress in human subjects are active fields of study, MRI research investigating PNSE in human subjects is just starting to be explored. As described below and shown in Table 3, many investigators have interrogated the impact of PNSE on the limbic system and connected regions in the developing brain.

Studies During Infancy

Recent studies suggest a significant relationship between antenatal maternal depression and/or anxiety and structure and

Table 2. Prenatal stress and the connectome: preclinical data

Author/year	Model	Protocol—fetuses	Outcome
HPA axis function Jutapakdeegul 2010 (124)	Pregnant rats Restraint stress Or Corticosterone 40mg/kg/d E14 – E21	Quantitative immuno-histochemistry	GAP-43 increased in prefrontal cortex (PFC) at PND 7 and 14
Afadlal 2010 (125)	Pregnant rats Corticosterone 40 mgkg/d	Levels of GAP-43PND 7, 14 and 60 Quantitative immuno-histochemistry	GAP-43 decreased in PFC at PND 60 Results identical for prenatal stress and corticosterone injection studies Corticosterone resulted in significant increase of GAP-43 and pGAP-43 in the hippocampus at PND 7 and PND 14
Antonow-Schlörke 2009 (126)	E14 – E21 Pregnant sheep	Levels of GAP-43, pGAP-43 and synaptophysin in the hippocampus PND 7 and 14 Outcomes at 20 d after treatment Immunohistochemistry Cell counting – mature oligodendroglia Immunohistochemistry – myelin basic protein (MBP)	Significant decrease in synaptophysin in hippocampus at same time points Within 24 h after treatment, betamethasone reduced number of mature oligodendrocytes and MBP immunoreactivity Maternal and fetal treatment had similar outcomes Loss of MBP immunoreactivity was not reversed 20 d after two treatment courses
Raschke 2008 (127)	Third trimester Betamethasone at 0.63, 0.75 and 0.87 gestation intramuscularly to mother or 48 h continuous infusion to fetus at 0.75 and 0.87 gestation Pregnant sheep Betamethasone administered at E 106/147 and E116/147 Lambs delivered preterm at 129/147 d gestation	Electron microscopy	Betamethasone resulted in a change in the formation of the myelin sheath in the commissural fibers of the corpus callosum but not in the association fibers of the subcortical white matter
Brain structure Zucchi 2013 (128)	Prenatal stress	Offspring: brain miRNA	Stress upregulated miR-219, a suppressor of neural stem cells Stress also upregulated miR-98, an miRNA shown to decrease the cerebral inflammatory response and increase blood brain barrier tightness Transcriptomic changes included genes related to development, axonal guidance and neuropathology Delay in embryonic neurogenesis of GABAergic progenitors
Uchida 2014 (129)	GAD67-green fluorescent protein (GFP) knock-in pregnant mice Restraint and light stress from E 15.0 – E17.5 Parvalbumin-(PV)positive interneurons	Offspring transcription profiles Outcomes at PND 21 Immunohistochemistry	PV-positive GABAergic neurons decreased in medial PFC, hippocampus (HPC) and somatosensory cortex of GAD67 ⁺ /GFP but not wild-type offspring Migration of GABAergic progenitors to cortex delayed in PNSE
Stevens 2013 (130)	BrdU at E15 or E12 Pregnant CAD67GFP transgenic mice Restraint stress E12 onwards	Unbiased cell counting Immunohistochemistry <i>In situ</i> hybridization PCR – gene expression	Significant changes in dlx2 and nkx2.1, transcription factors which regulate interneuron migration in PNSE forebrain No change mash 1 (determinant of interneuron fate), bdnf (maturation factor for GABAergic cells) or fgf2 (growth/differentiation factor) Total GABAergic cell number showed altered trajectories in medial PFC and HPC in PNSE mice
Lussier 2016 (80)	BrdU E13 Pregnant mice Restraint stress E12-birth Male offspring only	Cell counting Immunohistochemistry Quantitative PCR Behavior at 2–3 mo	Parvalbumin neuron proportion in juvenile brain was altered by PNSE but parvalbumin gene expression showed no changes PNSE exhibit behavior changes which correlate with GABAergic populations in medial PFC and HPC

Table 2. Continued on next page

Table 2. Continued

Author/year	Model	Protocol—fetuses	Outcome
Ehrlich 2015 (131)	Pregnant rats	Amygdala GABA interneuron expression Cl- trans-porters KCC2 and NKCC1	PNSE decreased KCC2 but increased NKCC1 expression
Petit 2015 (132)	Stress model E9-E20 Pregnant ewes	Behavior PND 28 Offspring expression levels and dendritic morphology PND 1; Dendritic morphology: Dlg4, Rac1, RhoA, Doc2B	PNSE increased anxiety in female pups Decreased expression Rac1 and NR1 in PFC
Bennett 2015 (133)	Last third gestation Unpredictable management challenges Pregnant guinea pigs	Synaptic transmission: Nr1, Grin2A, Grin2B HPA axis: Nr3C1 Immunohistochemistry, RT-PCR PND 21 Myelin basic protein (MBP), glial fibrillary acidic protein (GFAP)	Overexpression Dlg4 in amygdale Alterations in spine density and structure in PFC and CA1 region HPC Reduction in MBP and GFAP in CA1 region HPC
Brain function and connectivity	Last third gestation Strobe light exposure	GABAA receptor subunit Behavior testing PND 18	No change GABAA receptor subunit PNSE pups showed higher anxiety
	Ladostigil (L, a mono-amine oxidase inhibitor) ± stress	Monoamine oxidase levels Rs fMRI	Ladostigil inhibited MAO A & B by 45–50%
	Pregnant rats L or control - E7–PND21		PNSE increased limbic connectivity in the R hemisphere All connections associated with dopaminergic system (R nucleus accumbens with frontal, cingulate, septum and motor/sensory cortices; R globus pallidus with infra-limbic and dentate gyrus) L prevented stress-induced changes
Skelin 2015 (135)	Stress exposure E13–214 groups: PNSE L; PNSE no L; No PNSE – L; no PNSE no L Pregnant rats PNSE during last third gestation over four generations (multigenerational prenatally stressed, MGPNs) Pregnant rats	Unilateral field potential recording power (FP)	FP power lower in medial PFC, amygdala, HPC and striatum in MGPNs pups Coherence of FPs between brain regions high in MGPNs animals among all structures
Ehrlich 2015 (136)		Electrophysiology – PND 10, 14, 17, 20, 28 and 60 Behavior – PND 60 In vivo electrophysiology	PNSE reduced amygdala neuron excitability across all days PNSE neurons had more hyperpolarized resting membrane potential and produced fewer action potentials PNSE animals decreased socio-emotional behavior Decreased basic synaptic activity in PNSE rats Long-term potentiation induction was decreased in CA3-CA1 pathway of hippocampus
Barzegar 2015 (137)	PNSE E17–20 Pregnant rats	Behavior at 1–2 mo Electrophysiology	Lower behavioral performance in PNSE rats PNSE showed decreased functional connectivity between neuronal discharge in medial PFC and hippocampal sharp-wave ripples
Negron-Oyarzo 2015 (138)	Noise stress Pregnant mice Prenatal stress model E14–21		
Inflammation Holloway 2013 (139)	Preclinical prenatal immune activation model	Expression serotonin 5-HT(2A) and metabotropic glutamate 2 (mGlu2) receptors Behavior	Prenatal immune activation increased 5-HT(2A) and decreased mGlu2 expression in frontal cortex Pattern of expression associated with behavioral changes Similar changes in prenatal stress model

Table 3. Prenatal stress and the connectome: clinical data

Author/year	Number	Risk factors/time	Age at scan	Outcome	Results
Neonates					
Rifkin-Graboi 2013 (24)	157	EPDS at 26 wk GA	6–14 d	dMRI – amygdala	Lower FA ($P = 0.009$) but not volume in R amygdala in infants with high prenatal stress compared to low-normal stress
Qiu 2013 (27)	175	STAI at 26 wk GA	All at birth 35 wk/repeat scans at 6 mo	Hippocampal volume and growth trajectory	No influence of maternal anxiety on hippocampal volumes at birth Neonates with maternal anxiety showed slower growth of bilateral hippocampal volumes over first 6 mo compared to neonates with no antenatal maternal anxiety
Rifkin-Graboi 2015 (25)	54	STAI at 26 wk GA 21 with high scores 33 with low scores	5–17 d	dMRI	High anxiety group – reduced FA in R insula, middle occipital, inf temporal, angular gyrus, uncinate, posterior cingulate, dorsolateral prefrontal, inferior frontal region, cerebellum and inferior fronto-occipital fasciculus; also bilat superior and L postcentral regions
Qiu 2015 (26)	24	EPDS at 26 wk GA	6 mo	fMRI – amygdala connectivity	Neonates with high prenatal stress had greater connectivity of amygdala with L temporal cortex, insula, bilat ACC, medial orbitofrontal ventromedial prefrontal cortices
Scheinost 2016 (82)	26	Retrospective review of maternal chart for prenatal stress as documented by diagnosis of maternal depression and/or anxiety	Term equivalent age (35–40 wk PMA)	fMRI – amygdala connectivity	Preterm neonates with high prenatal stress had reduced connectivity of amygdala to subcortical regions, including the thalamus.
Childhood					
Buss 2010 (83)	35	Pregnancy anxiety scale administered at 19, 25, and 31 wk GA	6–10 y	Voxel-based morphometry	Anxiety at 19 wk GA associated with decreased gray matter in prefrontal cortex, premotor cortex, medial temporal lobe, lateral temporal cortex, postcentral gyrus and cerebellum extending to the middle occipital gyrus and fusiform gyrus
Sarkar 2014 (84)	22	Prenatal stress	7 y	dMRI – FA and perpendicular diffusivity ($D_{\perp}$) – Uncinate and a control tract	Prenatal stress correlated positively with R uncinate FA and negatively with R uncinate $D_{\perp}$ No correlations with control tract
Sandman 2015 (47)	81	Centers for Epidemiologic Studies Depression scale (CESDS) at 19, 25, and 31 wk GA	6–9 y	MRI cortical thickness	Antenatal exposure to maternal depression at 25 wk GA was associated with cortical thinning in 24% of the frontal lobes, primarily R superior, medial orbital and frontal pole regions of the prefrontal cortex
Young adults					
Favaro 2015 (90)	35 females	Prenatal stress interview (140)	14–40 y	CBCL Voxel-based morphometry fMRI Depression survey	Association between cortical thinning in R prefrontal region and child externalizing behavior High prenatal stress associated with decreased gray matter in L medial temporal lobe and both amygdalae Prenatal stress showed positive linear relationship with connectivity between L medial temporal lobe and pregenual cortex Connectivity between L medial temporal lobe and L medial orbitofrontal cortex partially explained variance in depressive symptoms of offspring

EPDS, Edinburgh Postnatal Depression Scale.

function in the developing brain. Rifkin-Graboi performed structural MRI and dMRI on 157 nonsedated 6–14-day-old newborns whose mothers participated in the GUSTO study (Growing Up in Singapore Towards Healthy Outcomes), a cohort of Asian women enrolled during the first trimester of pregnancy. Socioeconomic status, prenatal exposures, pregnancy measures, and birth outcomes were recorded, and imaging data were analyzed only for those infants who met the following criteria: (i) gestational age (GA) ≥ 37 wk, (ii) birth weight (BW) $> 2,500$ g, and (iii) $Apgar_{5\min} > 7$. The Edinburgh Postnatal Depression Scale and the State Trait Anxiety Inventory (STAI) were administered to all women at 26 wk of pregnancy. Adjusting for household income, maternal age and smoking exposure, postmenstrual age (PMA) at MRI, and BW, Rifkin-Graboi (24) found significantly lower FA but not volume in the right amygdala in infants of mothers with high EDPS scores. This suggests a significant relationship between PNSE and microstructure of the right amygdala, a region associated with stress reactivity and vulnerability for mood disorders.

Similarly, Qiu interrogated the GUSTO cohort to examine the consequences of PNSE to maternal anxiety on neonatal development of the hippocampus, a structure critical for stress regulation (27). Entry criteria for this analysis differed from those of Rifkin-Graboi's 2013 study, and included both term and late preterm infants who met the following criteria: (i) GA ≥ 35 wk; (ii) BW $> 2,000$ g; and (iii) $Apgar_{5\min} > 9$. There were 175 GUSTO infants available for this analysis; 42 underwent repeat scans at age 6 mo, and 35 (83%) had usable data. In Qiu's analysis, antenatal maternal anxiety did not influence bilateral hippocampal volume at birth, but children of women with increased anxiety during pregnancy showed slower growth of both the left and right hippocampus between birth and age 6 mo. Subsequently, evaluating 21 GUSTO infants with high PNSE (i.e., maternal STAI > 90) and 34 with low PNSE (i.e., maternal STAI < 70), Rifkin-Graboi showed that antenatal anxiety predicted decreases in FA of regions important for cognitive-emotional responses to stress (i.e., right insula and dorsolateral prefrontal cortices (PFC)), sensory processing (right middle occipital cortex), and socio-emotional function (i.e., right angular gyrus, uncinate fasciculus, posterior cingulate, and parahippocampus) at age 5–17 d (25). Of note, infants were eligible for this analysis if met the following criteria: (i) GA ≥ 36 wk; (ii) BW $> 2,000$ g; and (iii) $Apgar_{5\min} > 7$.

Finally, Scheinost (82) and Qiu (26) investigated prenatal depression/anxiety exposure and amygdala connectivity using rs-fMRI in preterm neonates at term equivalent age and infants at age 6 mo, respectively. These data showed that, in the neonatal period, the amygdala is functionally connected to subcortical and posterior cortical regions, and, by age 6 mo, is connected to widespread networks subserving emotional regulation, memory, and social cognition. In preterm neonates, Scheinost showed that PNSE reduces amygdalar-thalamic connectivity and is additive to effects of preterm birth. Using 24 GUSTO infants, Qui showed that infants born to mothers with higher prenatal depressive symptoms had greater

Figure 2. Development of left amygdala functional connectivity during the perinatal period. During the perinatal period, (a) left amygdala connectivity is first characterized by a largely local circuitry, (b) then begins to connect to ipsilateral regions in the frontal and temporal lobes, and (c) finally develops connections to the contralateral amygdala.

rs-fMRI of the amygdala with the left temporal cortex, insula, anterior cingulate (ACC), medial orbitofrontal, and ventromedial PFC. These networks are reported in children and adults with depression, suggesting that rs-fMRI data may foreshadow future neuropsychiatric disease.

Studies During Childhood

Studies of older children also suggest that maternal anxiety is associated with specific changes in brain morphology. Buss evaluated children ages 6–10 y whose mothers had been enrolled in a prospective study of pregnancy at the University of California, Irvine or Cedars Sinai Hospital in Los Angeles, CA, between 1998 and 2002 (83). Families were contacted again in 2007 and invited to participate in a follow-up study of their children to assess the influence of PNSE on brain development. At the time of this report, 35 mother–child dyads had both usable MRI data and complete maternal data. VBM on these children demonstrated that exposure to high maternal stress at 19 wk of gestation correlated with gray matter reductions in the PFC, premotor cortex, medial temporal lobe, lateral temporal cortex, post-central gyrus, and cerebellum extending to the middle occipital and fusiform gyri. Although the numbers are small and assessments of postnatal stress exposure were not included in the authors' analyses, high pregnancy stress at 25 and 31 wk of gestation was not associated with local reductions in gray matter volume, suggesting the importance of earlier exposure to gestational psychological stress. Similarly, Sarkar performed dMRI studies to assess both FA and perpendicular diffusivity (D_{perp}) on 22 children ages 6–9 y whose mothers were retrospectively assessed for PNSE when the children were age 17 mo (84). For these children, PNSE was positively correlated with right uncinate FA and negatively with right uncinate D_{perp} , while PNSE was not associated with control tract properties.

In addition, since reduced cortical volume and thickness have both been associated with a history of depression in adult populations (85,86), Sandman measured cortical thickness in

81 school aged children whose mothers had participated in the longitudinal study described above; (83) all were prospectively evaluated for depression at 19, 25, and 31 wk of gestation (47). Prenatal maternal depression exposure was associated with thinning in the right frontal lobe, and the strongest association was with exposure at 25 wk gestation. Morphological changes were primarily found in the superior, medial orbital, and frontal pole regions of the right PFC, consistent with data in adults with depressive symptomatology (85,86). Further, the significant association between prenatal depression exposure and child externalizing behavior in this cohort of children was mediated by these changes.

Studies During Adulthood

Finally, although MR studies of young adults with early life stress exposures are just beginning to emerge (87–89), Favaro explored the relationship between PNSE, cortical volumes and rs-fMRI in a sample of 35 healthy women aged 14–40 y (90). The sample was composed of volunteers to whose mothers a semi-structured interview assessing stress related events during pregnancy was administered. Subject scores were assigned based on interview data and used for MRI analyses. For these women, greater PNSE was associated with decreased gray matter volume in the left medial temporal lobe and both amygdalae. Strength of PNSE was positively correlated with rs-fMRI between the left medial temporal lobe and pre-genual cortex, and connectivity between the left medial temporal lobe and

left medial-orbitofrontal cortex partially explained variance in depressive symptoms in this cohort.

EMERGING FACTORS

As studies begin to investigate the impact of PNSE on the connectome, several factors from both preclinical and clinical data have emerged as key considerations for future studies. These factors include defining the normal developmental trajectories of the fetal connectome, the type, timing and duration of PNSE, and the fetal sex. Other factors not addressed within this review include assessing the role of paternal preconception stress and identifying the molecular signatures of PNSE.

Fetal Networks

“Trajectory analysis is central to the assessment of the impact of PNSE on the developing brain.” (2) Fetal rs-fMRI is an emerging technology obtaining information about neural network development *in utero* by directly measuring the fetal brain (91). These methods are needed to investigate the prenatal connectome as it develops and pinpoint how and when PNSE alters its development. Using cross-sectional functional connectivity data between 21–38 wk of gestation, fetuses show evidence of both long-range functional connectivity and the emergence of neural networks across the third trimester, mimicking those in older children and adults (92,93). However, both longitudinal and cross-sectional data are needed to more fully characterize the developmental trajectories of PNSE. To

Table 4. Prenatal stress and the connectome: Endocrine and genetic mechanisms

Author/year	Number	Risk factors/time	Age at scan	Outcome	Results
Hypothalamic-pituitary-adrenal axis					
Buss 2012 (33)	65	Maternal cortisol at 15, 19, 25, 31, and 37 wk GA	7 y	Child amygdala and hippocampal volumes Child affective problems	Higher cortisol levels at 15 wk GA associated with larger R amygdala volume in girls but not in boys Higher cortisol levels at 15 wk GA Associated with more affective problems in girls but not in boys
Davis 2013 (141)	54	Subjects with and without exposure to antenatal steroid exposure (ANS)	6–10 y	MRI – cortical thickness Child Behavior Check List (CBCL)	ANS children had bilateral cortical thinning; most significant region was rostral ACC Children with more affective problems had a thinner left rostral ACC
Candidate genes					
Qiu 2015 (72)	146	Pregnancy anxiety scale administered at 19, 25, and 31 wk GA	Newborn	Voxel-based morphometry Catechol-O-methyltransferase (COMT) genotypes	Individual COMT SNPs modulated association between antenatal maternal anxiety and prefrontal and parietal cortical thickness Among rs737865-val158met-rs165599 haplotypes, the A-val-G haplotype modulated positive associations of maternal anxiety with cortical thickness in right PFC and right parietal cortex The G-met-A mediates negative associations of anxiety with thickness in bilateral precentral gyrus and prefrontal cortex
Epigenetic mechanisms					
Chen 2015 (142)	247	Maternal anxiety (STAI) at 26 wk of gestation	Newborn	Regional brain volumes BDNF genotype and methylation status	Infant brain-derived neurotrophic factor (BDNF) genotype influenced association of prenatal anxiety on both epigenome as well as that between epigenome and right amygdala and left hippocampus volumes

STAI, State Trait Anxiety Inventory.

begin to address this problem, we performed longitudinal rs-fMRI on 10 typically developing fetuses at 30–32, 34–36 wk PMA and following term delivery. This study was approved by the Yale University Human Investigation Committee, and pregnant women signed consent for the protocol. Because of its documented role in neurobehavioral disorders and alterations in studies of PNSE described above, we interrogated the emergence of amygdala networks during the prenatal period. During the 3rd trimester, left amygdala connectivity is first characterized by local circuitry, then begins to connect to ipsilateral regions in the frontal and temporal lobes, and finally develops connections to the contralateral amygdala ([Figure 2](#)). The development of these important cross-hemispheric connections between the right and left amygdala develop during the end of the third trimester and likely increases the vulnerability of this circuitry to PNSE (55).

Timing of Stress Exposure

There is increasing recognition that fetal stress exposure has a particularly pronounced impact during early periods of corticogenesis, commonly known as critical periods in the developing brain. Critical periods refer to epochs characterized by both increasing plasticity and greater vulnerability; thus, these are times when the developing brain may be most easily modified in either favorable or unfavorable directions. Critical periods are thought to be environmentally sensitive, and many authors believe they underlie the developmental origins of neurobehavioral disorders such as ASD.

Typically developing fetuses with PNSE during the middle second and third trimesters of gestation are reported to be at the greatest risk for neurobehavioral disorders (13,52). Reviewing Swedish birth registries, Class examined associations between PNSE in 738,144 offspring born in 1992–2000 for childhood outcomes and 2,155,221 offspring born in 1973–1997 for adult outcomes. Although data for GA are not available, third trimester bereavement stress significantly increased risk of both ASD and attention deficit hyperactivity disorder (55). Similarly, children who had been exposed to tropical storms during gestation months 5–6 or 9–10 had 3.8 times greater risk of developing ASD than children who had been exposed to the same storms, in the same place, but during other months of gestation (52). Duration of maternal stress may also play a role. Analyzing data from 4,682 live births, Latendresse reported that children of mothers with the longest periods of prenatal depression exposure experienced more than seven times increased risk for pervasive developmental disorder when compared to children with no PNSE (53).

In contrast, in the GUSTO study, mothers were assessed for gestational depression and/or anxiety at 26 wk, and MRI measures were correlated with these data (24–27). In addition, Sandman performed depression screening on 82 pregnant mothers at 19, 25, and 31 wk gestation and found that antenatal exposure to maternal depression at 25 wk gestation was significantly correlated with cortical thinning in 24% of the frontal lobes in the offspring (47). Finally, although cortisol levels are not available for subjects in the prior MRI studies,

high levels of maternal cortisol at 15 wk (but not 19, 25, 31, or 37 wk) of gestation were associated with amygdala volumetric changes in girls but not in boys (33). Since high levels of cortisol are believed to reprogram the fetal HPA axis and maternal stress has been reported to downregulate 11 β -hydroxysteroid dehydrogenase (75), the placental enzyme which metabolizes cortisol (75), future studies of maternal psychological stress during gestation should consider longitudinal assessments of maternal cortisol in tandem with fetal neuroimaging.

Sex Differences in Prenatal Stress Outcomes

The link between PNSE and outcomes may be moderated by fetal sex. The source of sex differences upon early development is unclear but may include placental functioning, exposure to adrenal hormones and testosterone and an assortment of epigenetic mechanisms (94–97). Recent fetal pathways also proposed include sex-dependent responses of the transcriptome (6,98–100), naturally occurring sexually-dimorphic processes mediating neuron-glia interactions (101), and differential responses of target regions in the developing brain (102). Thus, while PNSE may have consequences for both males and females, the specificity of effects may differ. To the best of our knowledge, however, only a single study has reported sex differences in MRI outcome measures. These data suggest that higher cortisol levels at 15 wk of gestation were associated with larger right amygdala volumes and more affective problems in female but not male offspring (33).

MECHANISMS OF PRENATAL STRESS AND THE CONNECTOME

Taken together, published studies of PNSE suggest both proximate and long-lasting influence on the connectome. However, mechanisms of how PNSE alters the developing connectome must be explored. Mechanistic studies have focused on the HPA axis, candidate genes, and epigenetic pathways (see [Table 4](#)).

MOVING FORWARD: INVESTIGATION OF THE CLINICAL PROBLEM, CHANGES IN CARE

Converging data suggest that PNSE alters the developing connectome. As noted by Sporns, “The placement of brain connectivity as an intermediate phenotype between environmental exposures and behavior makes it an important target for studies that link networks across levels from behavior to molecules, neurons and emerging networks in the developing brain” (62). To better address the impact of PNSE on the connectome, longitudinal studies of maternal/fetal dyads with and without stress exposure are needed. Such investigations would benefit from repeated assessments of maternal stress in order to identify type, time of onset, and duration of PNSE and correlate these data with sequential imaging. In addition, preconceptional stress may influence offspring outcome, and pregnant women should be surveyed for cumulative stress at the time of study enrollment. Likewise, both genetic variants and epigenetic changes may contribute to outcome in the offspring, and consideration should be made to include these data in PNSE-offspring outcome analyses. Finally, longitudinal

fetal imaging will provide important information about target regions, and developmental trajectory analyses are well suited for interrogation of the developing connectome.

These strategies can be used to detect developmental disturbances of the connectome that may underlie the development of neurobehavioral disorders.

ACKNOWLEDGMENTS

We are indebted to our medical, nursing and research colleagues and the infants and their mothers who agreed to take part in this study. We are also grateful to Michael Labrec, R.T.R., for technical expertise.

STATEMENT OF FINANCIAL SUPPORT

This work was supported by Gates Foundation OPP1119263, Seattle, WA, and National Institutes of Health T32 HD07094, Bethesda, MD.

Disclosures: None of the authors have disclosures.

REFERENCES

- Dennis EL, Thompson PM. Typical and atypical brain development: a review of neuroimaging studies. *Dialogues Clin Neurosci* 2013;15:359–84.
- Di Martino A, Fair DA, Kelly C, et al. Unraveling the miswired connectome: a developmental perspective. *Neuron* 2014;83:1335–53.
- Graham AM, Pfeifer JH, Fisher PA, Lin W, Gao W, Fair DA. The potential of infant fMRI research and the study of early life stress as a promising exemplar. *Dev Cogn Neurosci* 2015;12:12–39.
- Provençal N, Binder EB. The effects of early life stress on the epigenome: From the womb to adulthood and even before. *Exp Neurol* 2015;268:10–20.
- Chen Y, Baram TZ. Toward understanding how early-life stress reprograms cognitive and emotional brain networks. *Neuropsychopharmacology* 2016;41:197–206.
- Bronson SL, Bale TL. Prenatal stress-induced increases in placental inflammation and offspring hyperactivity are male-specific and ameliorated by maternal antiinflammatory treatment. *Endocrinology* 2014;155:2635–46.
- Constantinof A, Moisiadis VG, Matthews SG. Programming of stress pathways: A transgenerational perspective. *J Steroid Biochem Mol Biol* 2016;160:175–80.
- Sinha R. Chronic stress, drug use, and vulnerability to addiction. *Ann N Y Acad Sci* 2008;1141:105–30.
- McEwen BS. The neurobiology and neuroendocrinology of stress. Implications for post-traumatic stress disorder from a basic science perspective. *Psychiatr Clin North Am* 2002;25:469–94, ix.
- Brown AS. Epidemiologic studies of exposure to prenatal infection and risk of schizophrenia and autism. *Dev Neurobiol* 2012;72:1272–6.
- Howerton CL, Bale TL. Prenatal programming: at the intersection of maternal stress and immune activation. *Horm Behav* 2012;62:237–42.
- Khashan AS, Abel KM, McNamee R, et al. Higher risk of offspring schizophrenia following antenatal maternal exposure to severe adverse life events. *Arch Gen Psychiatry* 2008;65:146–52.
- Kinney DK, Munir KM, Crowley DJ, Miller AM. Prenatal stress and risk for autism. *Neurosci Biobehav Rev* 2008;32:1519–32.
- Koenig JI, Kirkpatrick B, Lee P. Glucocorticoid hormones and early brain development in schizophrenia. *Neuropsychopharmacology* 2002;27:309–18.
- Li J, Olsen J, Vestergaard M, Obel C. Attention-deficit/hyperactivity disorder in the offspring following prenatal maternal bereavement: a nationwide follow-up study in Denmark. *Eur Child Adolesc Psychiatry* 2010;19:747–53.
- Pearson RM, Evans J, Kounali D, et al. Maternal depression during pregnancy and the postnatal period: risks and possible mechanisms for offspring depression at age 18 years. *JAMA Psychiatry* 2013;70:1312–9.
- Talge NM, Neal C, Glover V; Early Stress, Translational Research and Prevention Science Network: Fetal and Neonatal Experience on Child and Adolescent Mental Health. Antenatal maternal stress and long-term effects on child neurodevelopment: how and why? *J Child Psychol Psychiatry* 2007;48:245–61.
- Entringer S, Buss C, Wadhwa PD. Prenatal stress, development, health and disease risk: a psychobiological perspective-2015 Curt Richter Award Paper. *Psychoneuroendocrinology* 2015;62:366–75.
- Montagna A, Nosarti C. Socio-emotional development following very pre-term birth: pathways to psychopathology. *Front Psychol* 2016;7:80.
- Ramchandani PG, Richter LM, Norris SA, Stein A. Maternal prenatal stress and later child behavioral problems in an urban South African setting. *J Am Acad Child Adolesc Psychiatry* 2010;49:239–47.
- Kinsella MT, Monk C. Impact of maternal stress, depression and anxiety on fetal neurobehavioral development. *Clin Obstet Gynecol* 2009;52:425–40.
- Bale TL, Baram TZ, Brown AS, et al. Early life programming and neurodevelopmental disorders. *Biol Psychiatry* 2010;68:314–9.
- Singh-Taylor A, Korosi A, Molet J, Gunn BG, Baram TZ. Synaptic rewiring of stress-sensitive neurons by early-life experience: a mechanism for resilience? *Neurobiol Stress* 2015;1:109–15.
- Rifkin-Graboi A, Bai J, Chen H, et al. Prenatal maternal depression associates with microstructure of right amygdala in neonates at birth. *Biol Psychiatry* 2013;74:837–44.
- Rifkin-Graboi A, Meaney MJ, Chen H, et al. Antenatal maternal anxiety predicts variations in neural structures implicated in anxiety disorders in newborns. *J Am Acad Child Adolesc Psychiatry* 2015;54:313–21.e2.
- Qiu A, Anh TT, Li Y, et al. Prenatal maternal depression alters amygdala functional connectivity in 6-month-old infants. *Transl Psychiatry* 2015;5:e508.
- Qiu A, Rifkin-Graboi A, Chen H, et al. Maternal anxiety and infants' hippocampal development: timing matters. *Transl Psychiatry* 2013;3:e306.
- Van Dam NT, Rando K, Potenza MN, Tuit K, Sinha R. Childhood maltreatment, altered limbic neurobiology, and substance use relapse severity via trauma-specific reductions in limbic gray matter volume. *JAMA Psychiatry* 2014;71:917–25.
- Rubin LP. Maternal and pediatric health and disease: integrating biosychosocial models and epigenetics. *Pediatr Res* 2016;79:127–35.
- Grandjean P, Barouki R, Bellinger DC, et al. Life-long implications of developmental exposure to environmental stressors: new perspectives. *Endocrinology* 2015;156:3408–15.
- Bock J, Wainstock T, Braun K, Segal M. Stress in utero: prenatal programming of brain plasticity and cognition. *Biol Psychiatry* 2015;78:315–26.
- Howerton CL, Morgan CP, Fischer DB, Bale TL. O-GlcNAc transferase (OGT) as a placental biomarker of maternal stress and reprogramming of CNS gene transcription in development. *Proc Natl Acad Sci USA* 2013;110:5169–74.
- Buss C, Davis EP, Shahbaba B, Pruessner JC, Head K, Sandman CA. Maternal cortisol over the course of pregnancy and subsequent child amygdala and hippocampus volumes and affective problems. *Proc Natl Acad Sci USA* 2012;109:E1312–9.
- Buss C, Entringer S, Davis EP, et al. Impaired executive function mediates the association between maternal pre-pregnancy body mass index and child ADHD symptoms. *PLoS One* 2012;7:e37758.
- Kertes DA, Kamin HS, Hughes DA, Rodney NC, Bhatt S, Mulligan CJ. Prenatal maternal stress predicts methylation of genes regulating the hypothalamic-pituitary-adrenocortical system in mothers and newborns in the Democratic Republic of Congo. *Child Dev* 2016;87:61–72.
- Janssen AB, Kertes DA, McNamara GI, et al. A role for the placenta in programming maternal mood and childhood behavioural disorders. *J Neuroendocrinol* 2016;28.
- Maselko J, Sikander S, Bhalotra S, et al. Effect of an early perinatal depression intervention on long-term child development outcomes: follow-up of the Thinking Healthy Programme randomised controlled trial. *Lancet Psychiatry* 2015;2:609–17.
- Yonkers KA, Wisner KL, Stewart DE, et al. The management of depression during pregnancy: a report from the American Psychiatric Association and the American College of Obstetricians and Gynecologists. *Gen Hosp Psychiatry* 2009;31:403–13.
- American College of Obstetricians and Gynecologists. Perinatal Depression Toolkit. 2016 <http://mail.ny.acog.org/website/DepressionToolkit.pdf>.

40. Mapayi B, Makanjuola RO, Mosaku SK, et al. Impact of intimate partner violence on anxiety and depression amongst women in Ile-Ife, Nigeria. *Arch Womens Ment Health* 2013;16:11–8.
41. Fisher J, Cabral de Mello M, Patel V, et al. Prevalence and determinants of common perinatal mental disorders in women in low- and lower-middle-income countries: a systematic review. *Bull World Health Organ* 2012;90:139G–49G.
42. Sawyer A, Ayers S, Smith H. Pre- and postnatal psychological wellbeing in Africa: a systematic review. *J Affect Disord* 2010;123:17–29.
43. Barthel D, Kriston L, Barkmann C, et al.; International CDS Study Group. Longitudinal course of ante- and postpartum generalized anxiety symptoms and associated factors in West-African women from Ghana and Côte d'Ivoire. *J Affect Disord* 2016;197:125–33.
44. ACOG Practice Bulletin: Clinical management guidelines for obstetrician-gynecologists number 92, April 2008 (replaces practice bulletin number 87, November 2007). Use of psychiatric medications during pregnancy and lactation. *Obstet Gynecol* 2008 111:1001–1020.
45. Ko JY, Farr SL, Dietz PM, Robbins CL. Depression and treatment among U.S. pregnant and nonpregnant women of reproductive age, 2005–2009. *J Womens Health (Larchmt)* 2012;21:830–6.
46. Lee AM, Lam SK, Sze Mun Lau SM, Chong CS, Chui HW, Fong DY. Prevalence, course, and risk factors for antenatal anxiety and depression. *Obstet Gynecol* 2007;110:1102–12.
47. Sandman CA, Buss C, Head K, Davis EP. Fetal exposure to maternal depressive symptoms is associated with cortical thickness in late childhood. *Biol Psychiatry* 2015;77:324–34.
48. Suri R, Lin AS, Cohen LS, Altshuler LL. Acute and long-term behavioral outcome of infants and children exposed in utero to either maternal depression or antidepressants: a review of the literature. *J Clin Psychiatry* 2014;75:e1142–52.
49. Harville E, Xiong X, Buekens P. Disasters and perinatal health: a systematic review. *Obstet Gynecol Surv* 2010;65:713–28.
50. Engel SM, Berkowitz GS, Wolff MS, Yehuda R. Psychological trauma associated with the World Trade Center attacks and its effect on pregnancy outcome. *Paediatr Perinat Epidemiol* 2005;19:334–41.
51. Brand SR, Engel SM, Canfield RL, Yehuda R. The effect of maternal PTSD following in utero trauma exposure on behavior and temperament in the 9-month-old infant. *Ann N Y Acad Sci* 2006;1071:454–8.
52. Kinney DK, Miller AM, Crowley DJ, Huang E, Gerber E. Autism prevalence following prenatal exposure to hurricanes and tropical storms in Louisiana. *J Autism Dev Disord* 2008;38:481–8.
53. Latendresse G, Wong B, Dyer J, Wilson B, Baksh L, Hogue C. Duration of Maternal Stress and Depression: Predictors of Newborn Admission to Neonatal Intensive Care Unit and Postpartum Depression. *Nurs Res* 2015;64:331–41.
54. Bock J, Poeschel J, Schindler J, et al. Transgenerational sex-specific impact of preconception stress on the development of dendritic spines and dendritic length in the medial prefrontal cortex. *Brain Struct Funct* 2016;221:855–63.
55. Class QA, Abel KM, Khashan AS, et al. Offspring psychopathology following preconception, prenatal and postnatal maternal bereavement stress. *Psychol Med* 2014;44:71–84.
56. Hara Y, Waters EM, McEwen BS, Morrison JH. Estrogen Effects on Cognitive and Synaptic Health Over the Lifecourse. *Physiol Rev* 2015;95:785–807.
57. Ansell EB, Rando K, Tuit K, Guarnaccia J, Sinha R. Cumulative adversity and smaller gray matter volume in medial prefrontal, anterior cingulate, and insula regions. *Biol Psychiatry* 2012;72:57–64.
58. Seo D, Tsou KA, Ansell EB, Potenza MN, Sinha R. Cumulative adversity sensitizes neural response to acute stress: association with health symptoms. *Neuropsychopharmacology* 2014;39:670–80.
59. Lewis AJ, Austin E, Galbally M. Prenatal maternal mental health and fetal growth restriction: a systematic review. *J Dev Orig Health Dis* 2016;7:416–28.
60. Sandman CA, Davis EP, Buss C, Glynn LM. Exposure to prenatal psychological stress exerts programming influences on the mother and her fetus. *Neuroendocrinology* 2012;95:7–21.
61. Rice F, Harold GT, Boivin J, van den Bree M, Hay DF, Thapar A. The links between prenatal stress and offspring development and psychopathology: disentangling environmental and inherited influences. *Psychol Med* 2010;40:335–45.
62. Sporns O. Structure and function of complex brain networks. *Dialogues Clin Neurosci* 2013;15:247–62.
63. Friston KJ. Functional and effective connectivity: a review. *Brain Connect* 2011;1:13–36.
64. Fox MD, Raichle ME. Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. *Nat Rev Neurosci* 2007;8:700–11.
65. Scheinost D, Finn ES, Tokoglu F, et al. Sex differences in normal age trajectories of functional brain networks. *Hum Brain Mapp* 2015;36:1524–35.
66. Mori S, Zhang J. Principles of diffusion tensor imaging and its applications to basic neuroscience research. *Neuron* 2006;51:527–39.
67. Mechelli A, Friston KJ, Frackowiak RS, Price CJ. Structural covariance in the human cortex. *J Neurosci* 2005;25:8303–10.
68. Evans AC. Networks of anatomical covariance. *Neuroimage* 2013;80:489–504.
69. Scheinost D, Kwon SH, Lacadie C, Vohr BR, Schneider KC, Papademetris X, Constable RT, Ment LR. Alterations in anatomical covariance in the prematurely born. *Cereb Cortex* 2016 doi:10.1093/cercor/bhv248.
70. Kolb B, Mychasiuk R, Muhammad A, Li Y, Frost DO, Gibb R. Experience and the developing prefrontal cortex. *Proc Natl Acad Sci USA* 2012;109 Suppl 2:17186–93.
71. Maniam J, Antoniadis C, Morris MJ. Early-life stress, HPA axis adaptation, and mechanisms contributing to later health outcomes. *Front Endocrinol (Lausanne)* 2014;5:73.
72. Qiu A, Tuan TA, Ong ML, et al. COMT haplotypes modulate associations of antenatal maternal anxiety and neonatal cortical morphology. *Am J Psychiatry* 2015;172:163–72.
73. Tsotsi S, Abdulla N, Li C, et al. Fetal DNA methylation of cortisol-related genes and infant neurobehavior at 6 months: The role of antenatal maternal anxiety. *Psychoneuroendocrinology* 2015;61:35.
74. Bock J, Rether K, Gröger N, Xie L, Braun K. Perinatal programming of emotional brain circuits: an integrative view from systems to molecules. *Front Neurosci* 2014;8:11.
75. O'Donnell KJ, Bugge Jensen A, Freeman L, Khalife N, O'Connor TG, Glover V. Maternal prenatal anxiety and downregulation of placental 11 β -HSD2. *Psychoneuroendocrinology* 2012;37:818–26.
76. Mairesse J, Lesage J, Breton C, et al. Maternal stress alters endocrine function of the feto-placental unit in rats. *Am J Physiol Endocrinol Metab* 2007;292:E1526–33.
77. Jensen Peña C, Monk C, Champagne FA. Epigenetic effects of prenatal stress on 11 β -hydroxysteroid dehydrogenase-2 in the placenta and fetal brain. *PLoS One* 2012;7:e39791.
78. Lester BM, Padbury JF. Third pathophysiology of prenatal cocaine exposure. *Dev Neurosci* 2009;31:23–35.
79. Li Y, Gonzalez P, Zhang L. Fetal stress and programming of hypoxic/ischemic-sensitive phenotype in the neonatal brain: mechanisms and possible interventions. *Prog Neurobiol* 2012;98:145–65.
80. Lussier SJ, Stevens HE. Delays in GABAergic interneuron development and behavioral inhibition after prenatal stress. *Dev Neurobiol* 2016;76:1078–91.
81. Turkheimer FE, Leech R, Expert P, Lord LD, Vernon AC. The brain's code and its canonical computational motifs. From sensory cortex to the default mode network: A multi-scale model of brain function in health and disease. *Neurosci Biobehav Rev* 2015;55:211–22.
82. Scheinost D, Kwon SH, Lacadie C, et al. Prenatal stress alters amygdala functional connectivity in preterm neonates. *Neuroimage Clin* 2016;12:381–8.
83. Buss C, Davis EP, Muftuler LT, Head K, Sandman CA. High pregnancy anxiety during mid-gestation is associated with decreased gray matter density in 6–9-year-old children. *Psychoneuroendocrinology* 2010;35:141–53.
84. Sarkar S, Craig MC, Dell'Acqua F, et al. Prenatal stress and limbic-prefrontal white matter microstructure in children aged 6–9 years: a preliminary diffusion tensor imaging study. *World J Biol Psychiatry* 2014;15:346–52.

85. Peterson BS, Warner V, Bansal R, et al. Cortical thinning in persons at increased familial risk for major depression. *Proc Natl Acad Sci USA* 2009;106:6273–8.
86. Peterson BS, Weissman MM. A brain-based endophenotype for major depressive disorder. *Annu Rev Med* 2011;62:461–74.
87. Grant MM, Wood K, Sreenivasan K, et al. Influence of early life stress on intra- and extra-amygdaloid causal connectivity. *Neuropsychopharmacology* 2015;40:1782–93.
88. Holz NE, Boecker R, Hohm E, et al. The long-term impact of early life poverty on orbitofrontal cortex volume in adulthood: results from a prospective study over 25 years. *Neuropsychopharmacology* 2015;40:996–1004.
89. Fonzo GA, Ramsawh HJ, Flagan TM, et al. Early life stress and the anxious brain: evidence for a neural mechanism linking childhood emotional maltreatment to anxiety in adulthood. *Psychol Med* 2016;46:1037–54.
90. Favaro A, Tenconi E, Degortes D, Manara R, Santonastaso P. Neural correlates of prenatal stress in young women. *Psychol Med* 2015;45:2533–43.
91. Schöpf V, Kasprian G, Brugger PC, Prayer D. Watching the fetal brain at 'rest'. *Int J Dev Neurosci* 2012;30:11–7.
92. Anderson AL, Thomason ME. Functional plasticity before the cradle: a review of neural functional imaging in the human fetus. *Neurosci Biobehav Rev* 2013;37(9 Pt B):2220–32.
93. Thomason ME, Grove LE, Lozon TA Jr, et al. Age-related increases in long-range connectivity in fetal functional neural connectivity networks in utero. *Dev Cogn Neurosci* 2015;11:96–104.
94. Weinstock M. Sex-dependent changes induced by prenatal stress in cortical and hippocampal morphology and behaviour in rats: an update. *Stress* 2011;14:604–13.
95. Schaafsma SM, Pfaff DW. Etiologies underlying sex differences in Autism Spectrum Disorders. *Front Neuroendocrinol* 2014;35:255–71.
96. Lai MC, Lombardo MV, Auyeung B, Chakrabarti B, Baron-Cohen S. Sex/gender differences and autism: setting the scene for future research. *J Am Acad Child Adolesc Psychiatry* 2015;54:11–24.
97. Romano E, Cosentino L, Laviola G, De Filippis B. Genes and sex hormones interaction in neurodevelopmental disorders. *Neurosci Biobehav Rev* 2016;67:9–24.
98. Grundwald NJ, Benitez DP, Brunton PJ. Sex-dependent effects of prenatal stress on social memory in rats: a role for differential expression of central vasopressin-1a receptors. *J Neuroendocrinol* 2015;28 doi:10.1111/jne.12343
99. Grundwald NJ, Brunton PJ. Prenatal stress programs neuroendocrine stress responses and affective behaviors in second generation rats in a sex-dependent manner. *Psychoneuroendocrinology* 2015;62:204–16.
100. Mychasiuk R, Gibb R, Kolb B. Prenatal stress produces sexually dimorphic and regionally specific changes in gene expression in hippocampus and frontal cortex of developing rat offspring. *Dev Neurosci* 2011;33:531–8.
101. Werling DM, Parikshak NN, Geschwind DH. Gene expression in human brain implicates sexually dimorphic pathways in autism spectrum disorders. *Nat Commun* 2016;7:10717.
102. Frahm KA, Pepper ME, Zhang JY, et al. Research resource: the dexamethasone transcriptome in hypothalamic embryonic neural stem cells. *Mol Endocrinol* 2016;30:144–54.
103. Ronald A, Pennell CE, Whitehouse AJ. Prenatal maternal stress associated with ADHD and autistic traits in early childhood. *Front Psychol* 2010;1:223.
104. Grossi E, Veggo F, Narzisi A, Compare A, Muratori F. Pregnancy risk factors in autism: a pilot study with artificial neural networks. *Pediatr Res* 2016;79:339–47.
105. Gao L, Xi QQ, Wu J, et al. Association between prenatal environmental factors and child autism: a case control study in Tianjin, China. *Biomed Environ Sci* 2015;28:642–50.
106. Visser JC, Rommelse N, Vink L, et al. Narrowly versus broadly defined autism spectrum disorders: differences in pre- and perinatal risk factors. *J Autism Dev Disord* 2013;43:1505–16.
107. Van den Bergh BR, Mennes M, Oosterlaan J, et al. High antenatal maternal anxiety is related to impulsivity during performance on cognitive tasks in 14- and 15-year-olds. *Neurosci Biobehav Rev* 2005;29:259–69.
108. van den Bergh BR, Mennes M, Stevens V, et al. ADHD deficit as measured in adolescent boys with a continuous performance task is related to antenatal maternal anxiety. *Pediatr Res* 2006;59:78–82.
109. Markham JA, Koenig JI. Prenatal stress: role in psychotic and depressive diseases. *Psychopharmacology (Berl)* 2011;214:89–106.
110. Buss C, Davis EP, Hobel CJ, Sandman CA. Maternal pregnancy-specific anxiety is associated with child executive function at 6-9 years age. *Stress* 2011;14:665–76.
111. Laplante DP, Brunet A, Schmitz N, Ciampi A, King S. Project Ice Storm: prenatal maternal stress affects cognitive and linguistic functioning in 5 ½-year-old children. *J Am Acad Child Adolesc Psychiatry* 2008;47:1063–72.
112. Davis EP, Sandman CA. The timing of prenatal exposure to maternal cortisol and psychosocial stress is associated with human infant cognitive development. *Child Dev* 2010;81:131–48.
113. Yehuda R, Bell A, Bierer LM, Schmeidler J. Maternal, not paternal, PTSD is related to increased risk for PTSD in offspring of Holocaust survivors. *J Psychiatr Res* 2008;42:1104–11.
114. Betts KS, Williams GM, Najman JM, Alati R. Maternal depressive, anxious, and stress symptoms during pregnancy predict internalizing problems in adolescence. *Depress Anxiety* 2014;31:9–18.
115. Austin MP, Hadzi-Pavlovic D, Leader L, Saint K, Parker G. Maternal trait anxiety, depression and life event stress in pregnancy: relationships with infant temperament. *Early Hum Dev* 2005;81:183–90.
116. Blair MM, Glynn LM, Sandman CA, Davis EP. Prenatal maternal anxiety and early childhood temperament. *Stress* 2011;14:644–51.
117. Salisbury AL, O'Grady KE, Battle CL, et al. The roles of maternal depression, serotonin reuptake inhibitor treatment, and concomitant benzodiazepine use on infant neurobehavioral functioning over the first postnatal month. *Am J Psychiatry* 2016;173:147–57.
118. Monk C, Feng T, Lee S, Krupka I, Champagne FA, Tycko B. Distress during pregnancy: epigenetic regulation of placenta glucocorticoid-related genes and fetal neurobehavior. *Am J Psychiatry* 2016;173:705–13.
119. Davis EP, Glynn LM, Waffarn F, Sandman CA. Prenatal maternal stress programs infant stress regulation. *J Child Psychol Psychiatry* 2011;52:119–29.
120. Dorringston S, Zammit S, Asher L, Evans J, Heron J, Lewis G. Perinatal maternal life events and psychotic experiences in children at twelve years in a birth cohort study. *Schizophr Res* 2014;152:158–63.
121. Betts KS, Williams GM, Najman JM, Alati R. The relationship between maternal depressive, anxious, and stress symptoms during pregnancy and adult offspring behavioral and emotional problems. *Depress Anxiety* 2015;32:82–90.
122. Malaspina D, Corcoran C, Kleinhaus KR, et al. Acute maternal stress in pregnancy and schizophrenia in offspring: a cohort prospective study. *BMC Psychiatry* 2008;8:71.
123. Corcoran C, Perrin M, Harlap S, et al. Incidence of schizophrenia among second-generation immigrants in the Jerusalem perinatal cohort. *Schizophr Bull* 2009;35:596–602.
124. Jutapakdeegul N, Afadl S, Polaboon N, Phansuwan-Pujito P, Govitrapong P. Repeated restraint stress and corticosterone injections during late pregnancy alter GAP-43 expression in the hippocampus and prefrontal cortex of rat pups. *Int J Dev Neurosci* 2010;28:83–90.
125. Afadl S, Polaboon N, Surakul P, Govitrapong P, Jutapakdeegul N. Prenatal stress alters presynaptic marker proteins in the hippocampus of rat pups. *Neurosci Lett* 2010;470:24–7.
126. Antonow-Schlörke I, Helgert A, Gey C, et al. Adverse effects of antenatal glucocorticoids on cerebral myelination in sheep. *Obstet Gynecol* 2009;113:142–51.
127. Raschke C, Schmidt S, Schwab M, Jirikowski G. Effects of betamethasone treatment on central myelination in fetal sheep: an electron microscopical study. *Anat Histol Embryol* 2008;37:95–100.
128. Zucchi FC, Yao Y, Ward ID, et al. Maternal stress induces epigenetic signatures of psychiatric and neurological diseases in the offspring. *PLoS One* 2013;8:e56967.
129. Uchida T, Furukawa T, Iwata S, Yanagawa Y, Fukuda A. Selective loss of parvalbumin-positive GABAergic interneurons in the cerebral cortex of maternally stressed Gad1-heterozygous mouse offspring. *Transl Psychiatry* 2014;4:e371.

130. Stevens HE, Su T, Yanagawa Y, Vaccarino FM. Prenatal stress delays inhibitory neuron progenitor migration in the developing neocortex. *Psychoneuroendocrinology* 2013;38:509–21.
131. Ehrlich DE, Neigh GN, Bourke CH, et al. Prenatal stress, regardless of concurrent escitalopram treatment, alters behavior and amygdala gene expression of adolescent female rats. *Neuropharmacology* 2015;97:251–8.
132. Petit B, Boissy A, Zanella A, et al. Stress during pregnancy alters dendritic spine density and gene expression in the brain of new-born lambs. *Behav Brain Res* 2015;291:155–63.
133. Bennett GA, Palliser HK, Shaw JC, Walker D, Hirst JJ. Prenatal stress alters hippocampal neuroglia and increases anxiety in childhood. *Dev Neurosci* 2015;37:533–45.
134. Goelman G, Ilinca R, Zohar I, Weinstock M. Functional connectivity in prenatally stressed rats with and without maternal treatment with lisdostigil, a brain-selective monoamine oxidase inhibitor. *Eur J Neurosci* 2014;40:2734–43.
135. Skelin I, Needham MA, Molina LM, Metz GA, Gruber AJ. Multigenerational prenatal stress increases the coherence of brain signaling among cortico-striatal-limbic circuits in adult rats. *Neuroscience* 2015;289:270–8.
136. Ehrlich DE, Rainnie DG. Prenatal stress alters the development of socio-emotional behavior and amygdala neuron excitability in rats. *Neuropsychopharmacology* 2015;40:2135–45.
137. Barzegar M, Sajjadi FS, Taleai SA, Hamidi G, Salami M. Prenatal exposure to noise stress: anxiety, impaired spatial memory, and deteriorated hippocampal plasticity in postnatal life. *Hippocampus* 2015;25:187–96.
138. Negrón-Oyarzo I, Neira D, Espinosa N, Fuentealba P, Aboitiz F. Prenatal stress produces persistence of remote memory and disrupts functional connectivity in the hippocampal-prefrontal cortex axis. *Cereb Cortex* 2015;25:3132–43.
139. Holloway T, Moreno JL, Umali A, et al. Prenatal stress induces schizophrenia-like alterations of serotonin 2A and metabotropic glutamate 2 receptors in the adult offspring: role of maternal immune system. *J Neurosci* 2013;33:1088–98.
140. Rini CK, Dunkel-Schetter C, Wadhwa PD, Sandman CA. Psychological adaptation and birth outcomes: the role of personal resources, stress, and sociocultural context in pregnancy. *Health Psychol* 1999;18:333–45.
141. Davis EP, Sandman CA, Buss C, Wing DA, Head K. Fetal glucocorticoid exposure is associated with preadolescent brain development. *Biol Psychiatry* 2013;74:647–55.
142. Chen L, Pan H, Tuan TA, et al.; Gusto Study Group. Brain-derived neurotrophic factor (BDNF) Val66Met polymorphism influences the association of the methylenetetrahydrofolate cycle with maternal anxiety and neonatal brain volumes. *Dev Psychopathol* 2015;27:137–50.

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>

© The Author(s) (2016)