


AUTHOR CORRECTION OPEN



Correction: Suberoylanilide hydroxamic acid suppresses axonal damage and neurological dysfunction after subarachnoid hemorrhage via the HDAC1/HSP70/TDP-43 axis

Kui Luo, Zhifei Wang, Kai Zhuang, Shishan Yuan, Fei Liu and Aihua Liu


© The Author(s) 2022

Experimental & Molecular Medicine (2022) 54:1064-1065; https://doi.org/10.1038/s12276-022-00783-3

Correction to: Experimental & Molecular Medicine https://doi.org/10.1038/s12276-022-00761-9, published online 02 May 2022

In this article the unit in the Vertical Axis in figure 1A should be corrected; the figure should have appeared as shown below.

The original article has been corrected.



Published online: 4 July 2022

Open Access This article is licensed under a Creative Commons International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2022