

CORRECTION



Correction to: Revisiting the definition of bronchopulmonary dysplasia in premature infants at a single center quaternary neonatal intensive care unit

Faith Kim , David A. Bateman, Nimrod Goldshtrom , Rakesh Sahni, Jen Tien Wung and Aaron Wallman-Stokes

© The Author(s), under exclusive licence to Springer Nature America, Inc. 2021

Journal of Perinatology (2021) 41:2575; <https://doi.org/10.1038/s41372-021-01140-4>

Correction to: *Journal of Perinatology*

<https://doi.org/10.1038/s41372-021-00980-4>

Table 3 of this article represents the prevalence and not the incidence of BPD among each of the weight groups, which is indeed an error. Table 3 should have appeared as shown below. The original article has been corrected.

Table 3. Incidence of bronchopulmonary dysplasia using different definitions according to birth weight cohort.

Birth weight (g)	Total: 352 N (%)	BW (g)	GA (weeks)	VON N (%)	NIH 2001 N (%)	NIH 2018 N (%)
≤750	92 (25)	618 ± 97	26 ± 2	20 (22)	57 (62)*	57 (62)*
751–1000	85 (24)	883 ± 79	27 ± 2	7 (8)	32 (38)*	36 (42)*
1001–1500	176 (50)	1251 ± 69	30 ± 2	6 (3) ^a	12 (7)	26 (15)
All ≤1500	352	993 ± 285	28 ± 2	33 (9)	101 (29)	119 (33)

* $P < 0.001$ comparing incidence using NIH 2018 and NIH 2001 definitions to the VON definition, respectively.

^aOne infant included since transferred prior to 36 weeks PMA on supplemental oxygen.