CORRECTION

Correction: High-mobility group box 1 induces endoplasmic reticulum stress and activates hepatic stellate cells

Qin He¹ · Yu Fu² · Xiangming Ding¹ · Dongxiao Li¹ · Zi Wang¹ · Dean Tian¹ · Wei Yan¹

Published online: 31 October 2018

© United States & Canadian Academy of Pathology 2018

Correction to: *Laboratory Investigation*; <https://doi.org/10.1038/s41374-018-0085-9>; Article published online 29 June 2018

Following publication of the original article, the authors noticed a few errors in Figs. 1–3:

1. In Fig. 1c2, incorrect molecular weight and name of lamin was provided. The correct molecular weight should be 74 KD and the name of reference gene should be laminA/C.

2. In Fig. 2a, on the western blot image of the R-HSC cells, the wrong reference gene “GAPDH” was listed, which is the same as α -SMA.

3. In Fig. 2b, statistical analysis for the R-HSC cells was omitted.

4. In Fig. 2c, the incorrect image of HSC-T6 with 10 ng/ml HMGB1 was published, which is the same as NC.

5. In Fig. 3a, the wrong gene “GRP78” was listed, which is the same as IRE1 α . The western blot of GRP78 is incorrect. Please see correct figures (1–3) below.

The original article can be found online at <https://doi.org/10.1038/s41374-018-0085-9>.

✉ Dean Tian
datian@tjh.tjmu.edu.cn

✉ Wei Yan
yanwei@tjh.tjmu.edu.cn

¹ Department of Gastroenterology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

² Department of Gastroenterology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

Fig. 1

Fig. 2

Fig. 3