








CORRECTION



Correction: Adeno-associated virus (AAV) 9-mediated gene delivery of Nurr1 and Foxa2 ameliorates symptoms and pathologies of Alzheimer disease model mice by suppressing neuro-inflammation and glial pathology

Yunseon Yang , Min-Jong Seok , Ye Eun Kim, Yunjung Choi, Jae-Jin Song, Yanuar Alan Sulistio , Seong-hoon Kim, Mi-Yoon Chang, Soo-Jin Oh, Min-Ho Nam , Yun Kyung Kim , Tae-Gyun Kim, Heh-In Im , Seong-Ho Koh and Sang-Hun Lee 

© The Author(s), under exclusive licence to Springer Nature Limited 2023

Molecular Psychiatry (2023) 28:5375–5377; <https://doi.org/10.1038/s41380-023-02169-x>

Correction to: *Molecular Psychiatry* <https://doi.org/10.1038/s41380-022-01693-6>, published online 29 July 2022

1. In the figure legend of Fig. 4I (page 8, line 7), ‘hippocampal lysates’ need to be corrected to ‘cortical lysates’.
2. In Figs. 4I and 5I, in the originally published version of this article, the western blot images for β -ACTIN bands were incorrectly cropped. The bands were replaced by the correct ones. In addition, the graphs for the western blot data were revised.

Figure 4





Figure 5



The original article has been corrected.