

ARTICLE



Pollutants enhance IgE sensitization in the gut via local alteration of vitamin D-metabolizing enzymes

Eunsoo Kim¹, Astrid Bonnegarde-Bernard¹, Stephen O. Opiyo², Marisa R. Joldrichsen¹, Zayed Attia¹, Brian H. Ahmer³, Estelle Cormet-Boyaka¹ and Prosper N. Boyaka^{1,3,4}✉

© The Author(s), under exclusive licence to Society for Mucosal Immunology 2021

Mechanisms linking ingested pollutants to increased incidence of allergy are poorly understood. We report that mice exposed to low doses of cadmium develop higher IgE responses following oral allergen sensitization and more severe allergic symptoms upon allergen challenge. The environmentally relevant doses of this pollutant also induced oxidative/inflammatory responses in the gut of SPF, but not germ-free mice. Interestingly, the increased IgE responses correlated with stimulation of the vitamin D₃-metabolizing enzymes CYP27B1 and CYP24A1 in the gut and increased luminal levels of oxidized vitamin D₃ metabolites that are not ligands of the vitamin D receptor. Inhibition of CYP27B1 and CYP24A1 via oral administration of pharmacological inhibitors reduced IgE responses induced in mice orally exposed to cadmium. Our findings identify local alteration of vitamin D signaling as a new mechanism for induction of IgE responses by environmental pollutants. They also identify vitamin D₃-metabolizing enzymes as therapeutic targets for the treatment of allergy.

Mucosal Immunology (2022) 15:143–153; <https://doi.org/10.1038/s41385-021-00440-4>

INTRODUCTION

Food allergies are IgE-mediated hypersensitivity reactions with clinical symptoms that are capable of affecting multiple tissues, including the skin, airways, and gastrointestinal tract. The increased incidence of allergic diseases over the last few decades demonstrates that both genetic predisposition [1] and nongenetic factors, including the excessive use of antibiotics, especially during early life, contribute to the development of allergies [2, 3]. Environmental factors, such as pollutants and drugs, are also viewed as major contributors to the growing number of allergic individuals [4, 5]; however, how environmental pollutants affect allergic sensitization and the development of allergen-specific IgE responses and/or immune mechanisms during the effector phase of allergic responses remains poorly understood.

Cadmium (Cd) is a cationic heavy metal that is commonly used for metal plating, or for making pigments and batteries. Cd can remain in the air, soil, and water [6], and can be ingested through contaminated water, leafy vegetables, fish, or grains [6, 7]. The average daily intake of Cd through water and food has been estimated to be around 8–25 µg [7]. Approximately 20% of arable land in China is reportedly contaminated with heavy metals [8]. Furthermore, Cd content in contaminated rice paddy fields was found to range between 5 and 145 mg/kg, with Cd accumulation in brown rice between 1.9 and 9.4 mg/kg [9]. Because Cd has a low rate of excretion and a half-life in the body greater than 15 years [6], the chronic ingestion of subtoxic doses may have important health implications. Exposure to toxic doses of Cd has been shown to induce inflammatory responses [10, 11] and impair immune cell function via mechanisms involving the

production of reactive oxygen species (ROS) [12]. With the exception of occupational exposure, the general adult population is more likely to be exposed to subtoxic doses of Cd. However, how Cd affects T helper cell differentiation, immunoglobulin (Ig) class switching, and production of IgE remain largely unknown.

Here, we report that chronic ingestion of subtoxic, and environmentally relevant, doses of Cd [6, 13, 14], enhances the production of IgE and the severity of subsequent allergic symptoms. We also show that Cd enhances the expression of the vitamin D₃-oxidizing enzymes CYP27B1 and CYP24A1. The subsequent increased levels of oxidized vitamin D₃ metabolites that are not ligands of vitamin D receptor drive IgE responses in hosts exposed to subtoxic doses of this heavy metal.

RESULTS

Chronic ingestion of subtoxic doses of Cd enhances IgE responses

The gut is the primary target of ingested and inhaled Cd [7]. To mimic chronic exposure to subtoxic and environmentally relevant doses of Cd [13–15], groups of mice were given drinking water containing 10 µM CdCl₂ (Cd10) or 25 µM CdCl₂ (Cd25) for 28 days. Compared to controls (NoCd), mice exposed to Cd ingested the same amount of water and had similar body weight. They also had similar numbers of T and B lymphocytes in tissues (Fig. S1A), suggesting that the doses of Cd ingested had no major effect on the overall number of cells of adaptive immunity in gut-associated lymphoid tissues or in the spleen.

¹Department of Veterinary Biosciences, The Ohio State University, Columbus, OH, USA. ²Molecular, Cellular Imaging Center—Columbus, The Ohio State University, Columbus, OH, USA. ³Department of Microbial Infection and Immunity, The Ohio State University, Columbus, OH, USA. ⁴Infection Diseases Institute, The Ohio State University, Columbus, OH, USA. ✉email: boyaka.1@osu.edu




Fig. 1 Prior chronic ingestion of subtoxic doses of Cd enhances IgE responses and allergy severity. **a** Experimental scheme. Mice ($n = 5$ per group) were provided CdCl₂ [10 μ M (Cd10) or 25 μ M (Cd25)] in drinking water for 28 days and sensitized by the oral administration of OVA (1 mg) and cholera toxin (CT, 15 μ g). **b** OVA-specific serum Ab titers 1 week after the last sensitization (Day 14). **c** Allergen-specific CD4⁺ T cell responses. Splensens were collected on day 14 and restimulated in vitro with 1 mg/mL of OVA before flow cytometry analysis of CD4⁺ T cell cytokine responses. **d** Surface body temperatures of anesthetized mice before and 1 h after nasal allergen challenge with OVA (200 μ g). **e** Lung inflammation and mucus production, and **(f)** quantification of mucus production. Data represent one of at least four independent experiments with five mice per group. Data are expressed as the mean \pm SD. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ compared to No Cd.

To determine whether such exposure to Cd could modulate allergic sensitization and promote IgE responses, age-matched controls and mice exposed to Cd for 28 days were sensitized by oral administration of ovalbumin (OVA) and the Th2-promoting adjuvant cholera toxin (CT) (Fig. 1a). Cd-exposed mice produced higher levels of OVA-specific serum IgE antibody (Ab) responses than control No Cd mice (Fig. 1b). However, antigen-specific IgG1 and IgA responses were unchanged in Cd-exposed mice, indicating that Ig classes were differentially affected by this pollutant (Fig. 1b). We also found that 4 weeks of exposure of mice to Cd doses lower than Cd10 (i.e., Cd4) did not enhance IgE responses in this experimental model (Fig. S1B). The allergen-specific Ab responses in Cd-treated mice were associated with elevated frequencies of IL-4⁺CD4⁺, IFN γ ⁺CD4⁺, IL-17⁺CD4⁺, and IL-21⁺ CD4⁺ T cells in the splensens (Figs. 1c and S1C).

Chronic ingestion of subtoxic doses of Cd enhances allergic responses

To determine whether ingestion of cadmium could affect the severity of allergic responses, mice orally-sensitized were nasally challenged with the allergen. This nasal allergen challenge triggered a greater drop in body temperature in Cd-exposed mice than in controls, indicating more severe signs of allergy (Fig. 1d). Allergen challenge also increased the frequencies of CD4⁺ T cells, SSC^{hi}CD11c^{lo}GR1^{lo}CD3^{lo}CD19⁻ eosinophils and macrophages, including CD38⁺ M1 macrophages, in the bronchoalveolar lavage fluid (BALF) of mice exposed to Cd (Fig. S2A). The

increased number of eosinophils in the lungs of Cd-treated mice was further confirmed by the levels of eosinophil-associated *Siglec-F* and peroxidase (*Epx*) mRNA (Fig. S2B). When T cells collected from BALF were re-stimulated with an allergen in vitro, they showed an increased frequency of Th2 (IL-4), and Th1 (IFN γ) cells (Fig. S2C). This profile was consistent with elevated levels of IL-4, IFN γ , and TNF α measured in the BALF of mice exposed to Cd10 (Fig. S2D) which also exhibited lung inflammation as indicated by increased leukocyte infiltrates and mucus production after nasal allergen challenge (Fig. 1e, f).

We addressed whether stimulation of IgE responses by ingested Cd was mediated by events that occurred in mucosal tissues of the gut or the systemic compartment. In contrast with mice sensitized orally, mice sensitized via the systemic route (i.e., intraperitoneal (i.p.) route) by injection of OVA and CT as adjuvant showed no increase in OVA-specific IgE responses or other Ig isotype responses (Fig. S2E). Together, these data indicate that chronic ingestion of subtoxic doses of Cd enhances the magnitude of IgE responses via events that occur locally in gut tissues.

Chronic ingestion of subtoxic doses of Cd promotes oxidative stress and inflammatory responses in the intestines

Consistent with reports that Cd induces ROS production and oxidative stress responses [12], intestines of mice chronically exposed to subtoxic doses of Cd had higher mRNA levels of dual oxidase 2 (*Duox2*) and dual oxidase maturation factor 2 (*Duoxa2*)




Fig. 2 Chronic ingestion of subtoxic doses of Cd induces oxidative stress and inflammatory responses. Mice ($n = 5$ per group) were provided CdCl₂ [10 μ M (Cd10) or 25 μ M (Cd25)] in drinking water for 28 days. **a** Expression of mRNA levels of the oxidase subunits *Duox1*, *Duox2*, *Duoxa1*, and *Duoxa2*. cDNAs were generated from small intestines, and mRNA responses were analyzed by qRT-PCR. **b, c** Expression of NRF2. Sections of small intestine. **c** Quantification of NRF2 staining. **d** Cytokine mRNA responses in intestinal tissues. qRT-PCR data represent one of at least four independent experiments with five mice per group and are expressed as the mean \pm SD. **e–g** Expression of PGE₂ in small intestinal tissues. **e** Representative immunofluorescence staining. Sections of small intestines were stained with anti-PGE₂ antibody and counterstained with DAPI to visualize nuclei. **f** Quantification of PGE₂ staining. **g** Analysis of PGE₂ levels in small intestinal tissues by ELISA ($n = 5$ per group). * $p < 0.05$; ** $p < 0.01$, *** $p < 0.001$ compared to NoCd.

than controls (Fig. 2a). Antioxidants and phase II-detoxifying enzymes are under the control of the transcription factor nuclear factor erythroid 2-related factor 2 (NRF2) [12], and dysregulation of the NRF2 pathway stimulates the expression of inflammatory cytokines. Accordingly, mice exposed to subtoxic doses of Cd exhibited a dose-dependent induction of NRF2 in intestinal epithelial cells (Fig. 2b, c), and increased proinflammatory (i.e., *Ifn γ* , *IL-1 β* , *Mip2*, and *KC*) and anti-inflammatory (i.e., *IL-10* and *Tgf β*) mRNA responses (Fig. 2d). Furthermore, exposure of mice to Cd

does not increase the expression of Th2 (i.e., *IL-4* and *IL-5*) cytokines, but *IL-33* and *TSLP* were elevated by Cd exposure (Fig. 2d). Oxidative stress responses were recently shown to induce PGE₂ [16], and PGE₂ can promote IgE in vivo [17, 18]. Thus, we examined PGE₂ production in intestinal tissues by immunofluorescence staining (Fig. 2e, f). The higher levels of PGE₂ detected in tissues of mice treated with Cd10 and to a lesser extent Cd25 was confirmed by quantification of PGE₂ in small intestinal tissues by ELISA (Fig. 2g).




Fig. 3 IgE-promoting effects of low doses of Cd are lost in the absence of gut microbiota. **a** Scheme of experiments. Germ-free mice ($n = 5$ per group) were provided CdCl₂ [10 μM (Cd10) or 25 μM (Cd25)] in drinking water for 28 days. **b** Frequency of B and T lymphocytes in mucosal [Peyer's patches (PP); mesenteric lymph node (MLN)] and peripheral (spleen) lymphoid tissues from germ-free mice exposed to subtoxic doses of Cd. **c–e** Allergen-specific antibody responses after oral sensitization. **c** Allergen-specific serum IgE responses. **d** Allergen-specific serum IgA responses. **e** Allergen-specific serum IgG subclass responses. **f–i** Gut innate responses before sensitization. **f** Expression of NRF2. Sections of small intestines were stained with anti-NRF2 antibody and counterstained with DAPI to visualize nuclei. **g** Cytokine mRNA responses in intestinal tissues. **h, i** Expression of PGE₂. **h** Sections of small intestines were stained with anti-PGE₂ antibody and DAPI. **i** Quantification of PGE₂ levels in small intestinal tissues by ELISA. Data are expressed as mean ± SD. * $p < 0.05$; ** $p < 0.01$ and represent one of at least four independent experiments with five mice per group. Immunofluorescence images are representative of five mice per group.

IgE-promoting effects of low doses of Cd are lost in the absence of gut microbiota

We next used germ-free (GF) mice to address whether the increased IgE responses observed in Cd-treated mice reflected direct host response to Cd or indirect effects mediated through the gut microbiota (Fig. 3a). As observed in conventional specific pathogen-free (SPF) mice, chronic exposure to subtoxic doses of Cd did not change the frequency of T and B cells in lymphoid tissues of GF mice (Fig. 3b). Interestingly, prior exposure to subtoxic doses of Cd did not enhance allergen-specific IgE responses in GF mice (Fig. 3c). However, allergen-specific serum IgA (Fig. 3d) and IgG subclass (Fig. 3e) responses were increased in Cd-treated mice, suggesting that IgE responses are differentially regulated by Cd than other Ig isotypes. To better understand the inability of Cd to enhance IgE responses in germ-free mice, we examined oxidative stress and inflammatory responses in their gut tissues prior to allergic sensitization. NRF2 was not induced (Fig. 3f) and we detected minimal levels of inflammatory and type 2 cytokine mRNA responses with the exception of *Tnfa* mRNA responses which were significantly enhanced (Fig. 3g). Finally, Cd-treated and control germ-free mice failed to express high levels of PGE₂ in gut tissues (Fig. 3h, i).

Chronic ingestion of subtoxic doses of Cd alters the gut microbiome

Increased intestinal permeability is known to facilitate allergic sensitization. We found no difference between the absorption of orally administered FITC-dextran by gut tissues of control and Cd-treated mice either at the basal level (no CT, Fig. 4a). Furthermore, Cd-treated mice did not show higher intestinal permeability than control mice after oral administration of the adjuvant CT (CT 16 h, Fig. 4a). These results indicate that ingestion of these levels of Cd does not increase intestinal permeability. Although Cd has no known nutritional role in bacterial growth, it may interfere with divalent cationic metal ions that limit the growth of certain bacteria. Microbial colonization is the single most important event that controls intestinal secretory IgA (SIgA) levels [19, 20], and reduction of SIgA levels is a potential marker of dysbiosis [21, 22]. Intestinal SIgA levels were reduced after 2 weeks of ingestion of subtoxic doses of Cd and remained low after 28 days of exposure (Fig. 4b). Proteomic analyses of intestinal contents showed that mice treated with low doses of Cd (Cd10) expressed higher levels of the antimicrobial peptide regenerating islet-derived protein 3 beta (Fig. S3A). Other luminal proteins increased after Cd10 ingestion, including ferritin, laminin, and histone (Figs. 4c and S3A). In contrast, Cd exposure reduced overall levels of mucins,




Fig. 4 Chronic ingestion of subtoxic doses of Cd alters the gut microbiome, and luminal metabolite and proteomic profiles. Mice ($n = 5$ per group) were provided CdCl_2 [$10 \mu\text{M}$ (Cd10) or $25 \mu\text{M}$ (Cd25)] in drinking water for 28 days. **a** Intestinal permeability was analyzed by measuring serum FITC-dextran (4 kDa). **b** Fecal IgA levels. Total IgA levels in freshly emitted feces were analyzed by ELISA. The amounts of IgA were normalized by total protein content. **c** Relative proteins expression in mice treated with Cd10 vs. NoCd mice. **d** Bacterial load in fecal extracts. Dilution of fecal pellets was normalized by mass (g) of feces and plated on blood agar, and bacteria (CFU) were counted after 48 h of culture. **e** Relative abundance of Firmicutes to Bacteroidetes (upper) and other less dominant phyla (lower). **f** Metabolites in fecal extracts. **g** Principal component analysis of metabolite profiles. **g** Heat map of relative expression of metabolites with significantly different concentrations in samples from Cd-treated mice vs. NoCd mice ($*p < 0.05$). Data are expressed as the mean \pm SD. $*p < 0.05$; $**p < 0.01$ (4–5 mice per group).

kallikrein-1, and proteins associated with the digestive function of epithelial cells (e.g., aminopeptidase, alkaline phosphatase, sucrose isomaltase) (Figs. 4c, S3A, and S3B). Proteomics data also showed that Cd stimulates antimicrobial responses and impairs the function of goblet cells. Reduced levels of mucus may facilitate the function of goblet cells. Reduced levels of mucus has been reported to deliver regulatory signals that enhance oral tolerance [23]. Kallikrein-1 cleaves kininogens into kinins that enhance the permeability of capillaries and epithelial cell layers [24]. Thus, reduced levels of kallikrein-1 in fecal samples of mice exposed to Cd support the conclusions of

our FITC-dextran absorption studies (Fig. 4a), that intestinal permeability was not increased in these mice.

Ingestion of Cd did not affect the number of cultivable bacteria in the gastrointestinal tract (Fig. 4d) but did induce dysbiosis (Fig. 4e). The relative abundance of gut microbiota (Figs. 4e, S3C, and S3D) and linear discriminant analysis (LDA) (Fig. S3E) showed that Cd most significantly reduced bacteria of the genus *Prevotella*, whereas bacteria in genera *Tannerella* and *Turicibacter* increased the most in mice exposed to Cd10 and Cd25, respectively.




Fig. 5 Cd-induced gut microenvironment enhances IgE responses via stimulation of PGE₂. **a** Experimental scheme for the transfer of fecal materials from Cd-treated mice. **b, c** Allergen-specific serum IgE (**b**) and IgG1 (**c**) responses after oral sensitization of recipient mice not exposed to Cd. **d** IgM and IgE production in culture supernatants of naïve murine spleen cells cultured for 4 days in the presence of IL-4 (10 ng/ml), anti-CD40 (1 µg/ml), and bacteria-free fecal extracts ($n = 4$). **e** *Baff* and *April* mRNA responses in *Myd88* KO macrophages cultured 24 h in the presence of bacteria-free fecal extracts ($n = 5$). **f** Expression of BAFF in small intestinal tissues of naïve mice or mice exposed to subtoxic doses of Cd for 28 days ($n = 5$). **g** PGE₂ secretion from HT-29 cells 4 h after the addition of bacteria-free fecal extracts ($n = 5$). **h** *Baff*, *April*, and *Ptges* mRNA responses by MLN lymphocytes (2×10^5 cells/mL) stimulated for 24 h with anti-CD40 and IL-4 in the presence of bacteria-free fecal extracts only, or together with the COX2 inhibitor celecoxib (10 µM) ($n = 3$). **i** Effects of celecoxib on IgE production by MLN lymphocytes stimulated as described in (**h**) ($n = 3$). Data are from at least four independent experiments and are expressed as the mean \pm SD. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$ compared to NoCd. ## $p < 0.01$; ### $p < 0.001$; #### $p < 0.0001$ compared to no celecoxib.

Chronic ingestion of subtoxic doses of Cd alters metabolite composition and protein expression in the gut

Consistent with the notion that dysbiosis is generally associated with changes in the composition of metabolites in the gut, Cd-treated mice exhibited profiles of metabolites that differed from control mice (Fig. 4f, g). Statistical differences in metabolite composition between NoCd mice and at least one of the Cd-treated groups are depicted in Fig. 4g. In general, samples from Cd-treated mice exhibited reduced levels of metabolites related to food consumption and amino acid metabolism. The main feature of Cd-treated mice was a major increase in the levels of lipids and lipid-like molecules (Fig. 4g), principally, the vitamin D₃ metabolite 24,25,26,27-tetranor-23-oxo-hydroxyvitamin D₃ (oxo-VD₃) (Fig. 4g and Table S2). This finding was of high significance, since mice exposed to Cd10, which contain the highest levels of oxo-VD₃ in the gut, were also those that exhibited the highest IgE responses

upon oral sensitization (Fig. 2b). In this regard, vitamin D₃ metabolites were shown to be associated with increased production of PGE₂ [25], and PGE₂ promotes IgE production [17, 18].

IgE-enhancing effects of Cd can be transferred by fecal transplantation

To address the role of Cd-induced luminal metabolites/proteins and/or bacteria in increasing allergic sensitization, fecal materials from Cd-treated mice were transferred to groups of SPF mice, which were subsequently orally sensitized by the administration of OVA and CT as adjuvant (Fig. 5a). OVA-specific serum IgE responses (Fig. 5b), but not OVA-specific IgG subclasses or IgA (Figs. 5c and S4A), were enhanced in recipients of fecal material transplantation (FMT) from mice exposed to Cd. Furthermore, subsequent nasal allergen challenge resulted in higher lung




Fig. 6 Vitamin D₃-metabolizing enzymes and oxidized vitamin D₃ metabolites regulate IgE production in mice exposed to Cd. **a** *Cyp27b1* and *Cyp24a1* mRNA responses in small intestines from control mice or mice exposed to Cd for 28 days ($n = 5$ per group). **b** COX2 inhibitor (ASA; aspirin)- or antioxidant (NAC; N-acetyl cysteine)-mediated regulation of cytokine (*Baff* and *April*) and vitamin D₃-metabolizing enzyme (*Cyp24a1*) mRNAs in mice exposed to subtoxic doses of Cd. *Ptges* and *Duox2* mRNAs were measured to confirm inhibitor specificity. **c**, **d** Numbers of IgE⁺CD19⁺ cells in cultures of cells from spleen (**c**) and mesenteric lymph node (**d**) from IgE Verigem mice stimulated in vitro with anti-CD40 and IL-4 in the presence of bacteria-free fecal extracts from mice exposed to Cd, mice exposed to Cd and treated with a COX2 inhibitor (Cd + ASA), or mice exposed to Cd and treated with an antioxidant (Cd + NAC). **e** Antigen-specific IgE responses in Cd-treated mice treated with a COX2 inhibitor before allergic sensitization ($n = 4$). **f** In vivo regulation of *Baff*, *April*, and *Ptges* mRNA responses in mice exposed to subtoxic doses of Cd by ketoconazole, an inhibitor of vitamin D₃ metabolism ($n = 5$). **g** Antigen-specific IgE responses in Cd-treated mice treated with an inhibitor of vitamin D₃-metabolizing enzymes. Data are from at least four independent experiments and are expressed as mean \pm SD. * $p < 0.05$; ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ compared to control NoCd mice. ## $p < 0.01$, ### $p < 0.001$, #### $p < 0.0001$ compared to Cd alone.

inflammatory responses and mucus production in recipients of FMT from mice chronically exposed to Cd (Fig. S4B). In the reverse experiment (Fig. S4C–H), fecal material from control untreated mice reduced allergen-specific IgE responses in recipient Cd-treated mice (Fig. S4D) but had no effect on the other Ig isotypes (Fig. S4E). In addition, Cd-treated mice, which received fecal material from untreated mice, developed fewer signs of allergy after nasal allergen challenge, as manifested by a less significant drop in body temperature (Fig. S4F) and lower numbers of IFN γ ⁺CD4⁺ T and IL-4⁺CD4⁺ T cells in the BALF (Fig. S4G, H). Taken together, these results show that fecal materials of Cd-treated mice contain factors that regulate IgE responses.

The Cd-induced gut microenvironment enhances IgE responses via stimulation of PGE₂

The FMT experiments described above did not elucidate whether bacteria, bacteria product, or host molecule mediated the IgE-promoting effects. To elucidate the mechanism(s) driving IgE responses in Cd-treated mice, bacteria-free fecal extracts were added to spleen cells stimulated in vitro with anti-CD40 in the presence of IL-4, and secretion of IgE was assessed in cell culture

supernatants. Cells cultured in the presence of fecal extracts from Cd-treated mice produced more IgE than cells exposed to fecal extracts from mice that did not receive Cd (Fig. 5d). This effect was unlikely due to cytokines, which were either unchanged or reduced (IL-6 and GM-CSF) in the fecal extracts from Cd-treated mice (Fig. S5A). Fecal extracts from Cd-treated mice increased the mRNA levels of *April* and *Baff* members of the TNF family that support Ig class switching [26–28], in macrophages from *Myd88* KO mice (Fig. 5e) and enhanced *APRIL* and *C4BP* mRNA responses in human epithelial HT-29 cells (Fig. S5B). These findings indicate that the luminal contents of Cd-treated mice stimulate *APRIL*/*BAFF* expression via molecules distinct from microbial PAMPs and can induce similar responses in human epithelial cells. Finally, *BAFF* expression was upregulated in vivo in small intestinal tissues of naive mice exposed for 28 days to subtoxic doses of Cd (Fig. 5f).

Gut expansion of fungi was reported to stimulate PGE₂ and promote allergic inflammation [29] and PGE₂ supports the production of IgE but not IgG1 [18]. Control and Cd-treated mice had similar levels of PGE₂ in their feces (Fig. S5C) and showed no evidence of *Candida* overgrowth (Fig. S5D). On the other hand, bacteria-free fecal extracts from Cd-treated mice enhanced PGE₂




Fig. 7 Mechanisms of enhanced IgE sensitization by ingested cadmium. Chronic ingestion of subtoxic doses of Cd stimulates expression of the vitamin D metabolizing enzymes CYP27B1 and CYP24A1 leading to increased levels of oxidized vitamin D₃ metabolites. This results in enhanced PGE₂ responses to allergic sensitization and higher IgE responses.

production in both human HT-29 cells (Fig. 5g) and in LPS-unresponsive Trif/Tram double-KO murine macrophages (Fig. S5E). To determine the link between PGE₂, BAFF/APRIL, and IgE production, we added the specific COX2 inhibitor celecoxib to cultures of spleen and mesenteric lymph node cells (MLN) stimulated with anti-CD40 in the presence of IL-4 and bacteria-free fecal extracts from Cd-treated mice. Celecoxib suppressed the transcription of PGE₂ synthase (*Ptges*), *Baff*, and *April* (Figs. 5h, S5F, and S5G) and the production of IgE by these cells (Fig. 5i). Together, these results indicate that a product(s) found in fecal extracts stimulated the production of PGE₂ in vivo by intestinal epithelial cells and myeloid cells and that inhibition of PGE₂ response by COX-2 inhibitor limits the production of BAFF and APRIL and associated high IgE responses

Host vitamin D₃-metabolizing enzymes play key roles in the Cd-mediated increase in IgE responses

The main feature of fecal samples obtained from Cd-treated mice was the presence of high levels of oxo-VD₃, an oxidized vitamin D₃ metabolite. Oxidized vitamin D₃ metabolites can be induced by oxidation of active vitamin D₃ by 25-hydroxyvitamin D₃ 1-alpha-hydroxylase (CYP27B1) and the steroid 21-hydroxylase (CYP24A1) (Fig. S5H) [30, 31]. Interestingly, *Cyp27b1* mRNA levels were significantly enhanced in the intestines of mice chronically exposed to subtoxic doses of Cd (Fig. 6a). An inverse relationship has been reported between vitamin D₃ receptor (VDR) signaling and PGE₂ production [32]. Consistent with the fact that oxo-VD₃ is not a ligand of the VDR, and that expression of VDR is stimulated by its ligands [32], the addition of bacteria-free fecal extracts from Cd-treated mice to J774 macrophages cultures reduced *Vdr* mRNA levels (Fig. S5I).

To determine the sequence of events that led to the increase in IgE responses in mice exposed to Cd, control and Cd-treated mice were orally treated with inhibitors of oxidative stress (N-acetyl cysteine, NAC) or PGE₂ (the COX2 inhibitor aspirin, ASA). Treatment with NAC reduced the mRNA levels of *Ptges*, *Baff*, *April*, and *Cyp24a1* in Cd-treated mice (Fig. 6b). The COX2 inhibitor downregulated the mRNA levels of *Ptges*, *Baff*, and *April* in the intestines of Cd-treated mice but failed to reduce the expression of *Cyp24a1* (Fig. 6b). We next used in vitro cultures of the spleen, MLN, or Peyer's patch cells from IgE Verlgem mice to determine whether treatment with these inhibitors affects the IgE-promoting effects of the gut luminal microenvironment of Cd-treated mice. As depicted in Fig. 6c, NAC completely suppressed the

IgE-promoting effects of fecal materials from Cd-treated mice while Cd-treated mice that had been administered ASA orally retained their IgE-promoting capabilities in vitro. These findings indicate that products of the vitamin D₃-metabolizing enzymes that were present in the fecal materials of mice given ASA drove IgE responses. Furthermore, oral sensitization with OVA in the presence of CT induced the same levels of IgE in control and Cd-exposed mice that were treated with aspirin prior to sensitization (Fig. 6e). These results demonstrate that PGE₂ is a central checkpoint, downstream of vitamin D₃-metabolizing enzymes, in the increase in IgE responses in vivo in hosts exposed to subtoxic doses of Cd.

To conclusively demonstrate the role of vitamin D₃-metabolizing enzymes, mice exposed to Cd were treated orally with ketoconazole, a pharmacological inhibitor of CYP27B1 and CYP24A1 (Fig. 6f). This treatment suppressed the mRNA expression of *Cyp27b1*, *Cyp24a1*, *Ptges*, *Baff*, and *April* in the intestinal tissues of Cd-treated mice (Fig. 6f). It also reduced the levels of allergen-specific IgE induced after oral sensitization to levels seen in control mice that were not exposed to Cd (Fig. 6g). These results show that vitamin D₃-metabolizing enzymes are previously unknown regulators of allergic sensitization in hosts chronically exposed to low, but environmentally relevant, doses of this heavy metal pollutant (Fig. 7).

DISCUSSION

Extensive use of antibiotics in early life is often mentioned as a leading factor driving the increased prevalence of allergic diseases. But other factors including diet and environmental pollutants are also believed to play roles in the steady increase of allergic diseases worldwide [3, 4, 33]. Data presented herein reveal a previously unknown mechanism for increased allergic sensitization in the gut and possibly other mucosal tissues. More specifically, using chronic exposure to a subtoxic dose of Cd as an experimental system that induces oxidative responses, we have identified the induction of vitamin D₃-metabolizing enzymes in gut tissues and the subsequent increased luminal levels of vitamin D₃-oxidized metabolites which are not ligands of the VDR as central mechanisms for selective enhancement of IgE responses.

Production of IgE is initiated by IL-4 and IL-13, which provide the signals for the transcription of the IgE (ε) germline [34]. Engagement of CD40 on the B cell surface by CD40L expressed on activated T cells further enables Ig isotype switching to IgE and

subsequent IgE production. In vitro studies have shown that exposure of B cells to low doses of Cd (0.1 μM) inhibits IgE production [35]. We show that oral allergen sensitization of mice exposed to environmentally relevant doses of Cd actually increased IgE responses. Cd-treated mice also had increased allergen-specific Th1 and Th17 cells, which further enhance histological and clinical signs of allergy after allergen challenge. It was reported that splenocytes of the offspring from C57BL/6 mice exposed to 10 ppm Cd, produced lower amounts of IFN γ upon in vitro stimulation than cells from control mice [36]. Nonetheless, upon vaccination, these offspring developed higher levels of both T cell-dependent and -independent serum antibodies compared to control animals [36]. Although other Ig isotypes can contribute to allergic responses, in our study, Cd did not alter IgG1 or IgA responses. This suggests that IgE-promoting signals are selectively targeted/stimulated by this heavy metal in the gastrointestinal tract.

It is important that Cd did not alter intestinal barrier function or increase intestinal permeability as indicated by absorption of FITC-dextran and reduced levels of kallikrein-1 in the intestinal contents of mice exposed to Cd. On the other hand, subtoxic doses of Cd-induced oxidative stress and increased the levels of the oxidative stress-sensing transcription factor NRF2 in the small intestinal tissues. These oxidative stress responses are associated with the stimulation of inflammatory cytokines and PGE $_2$ production. These findings are consistent with the reported ability of Cd to induce oxidative stress [12], and that oral exposure to high doses of Cd through drinking water in early life promotes systemic inflammation and subsequent fat accumulation in male mice [37]. Interestingly, the effects of Cd on oxidative stress, inflammatory responses, and PGE $_2$ were not observed in germ-free mice. Accordingly, exposure to Cd failed to enhance the IgE responses in germ-free mice. In this regard, *Duox2* expression in the intestine was shown to depend on microbial colonization [38, 39] and to be partially induced by TRIF and canonical NF- κ B signaling [39]. The cationic metal manganese was recently shown to increase the sensitivity of cGAS to dsDNA and STING activation [40]. Thus, Cd, which is also a cationic metal that can accumulate in the cytosol, likely enhanced the reactivity of intestinal epithelial cells to TLR ligands from commensal microbes.

It was previously reported that PGE $_2$ produced by *Candida* overgrowth in the gut tissues of mice treated with antibiotics increases allergic airway inflammation via the induction of M2 macrophages [29]. The low doses of Cd used in our study did not affect the bacterial load or result in *Candida* overgrowth or PGE $_2$ secretion in fecal materials. On the other hand, overall fecal IgA levels were reduced by Cd ingestion. The two main events that regulate the levels of fecal IgA are the bacterial load in the intestine and the composition of the microbiome [19, 21, 22]. Accordingly, Cd-induced dysbiosis was characterized by increased levels of the genera *Tannerella* and *Turicibacter* in mice exposed to Cd10 and Cd25, respectively. Our findings are consistent with previous reports that oral exposure to 100 ppm of Cd reduces the proportion of *Bifidobacterium* and *Lactobacillus* [11] and increases the numbers of *Lactobacillaceae* and *Erysipelotrichaceae*, especially *Turicibacter* in the *Erysipelotrichaceae* family [41]. However, it is unlikely that these changes alone could selectively enhance IgE responses.

The most striking and unexpected findings from this work were that Cd (a) stimulates the expression of the vitamin D $_3$ -metabolizing enzymes CYP27B1 and CYP24A1 in the gut and (b) enhances the levels of oxidized vitamin D $_3$ metabolites, that are not ligands of the VDR. Since bacteria-free fecal extracts from Cd-treated mice downregulate *Vdr* mRNA expression by macrophage, collectively our data suggest that exposure to subtoxic doses of Cd mimics vitamin D deficiency. In this regard, epidemiologic data and animal studies have linked vitamin D

deficiency to atopy, asthma, and food allergy [42–44], and vitamin D is believed to protect against allergic diseases via inhibition of T cell proliferation, IFN γ and Th17 responses [45], and ILC2 [46], or stimulation of Treg [47]. Vitamin D supplementation has been investigated as a strategy to mitigate or prevent allergic diseases. The timing and dose of Vitamin D used for supplementation were found to be crucial. Thus, vitamin D is only protective when it was given before the allergy develops (i.e., before or early after birth) [42]. Furthermore, high-dose vitamin D supplementation failed to prevent allergic sensitization of infants [48].

Our study suggests that activation of vitamin D-metabolizing enzymes is perhaps the missing link in our understanding of allergic sensitization and the efficacy of vitamin D supplementation therapy. In fact, we show that conditions of vitamin D deficiency and enhanced allergic sensitization can be induced in vitamin D-sufficient hosts by the activation of vitamin D-metabolizing enzymes and thus, depletion of functional vitamin D. The fact that these enzymes are inducible and that their expression is regulated by the concentration of vitamin D could explain the reported failure of high-dose vitamin D supplementation to prevent allergic sensitization [48]. It is also important to note that Cd-induced oxidative stress responses were the main drivers of CYP27B1 and CYP24A1 expression in gut tissues. Diesel exhaust particles, which contain heavy metals and induce oxidative stress responses in the airways [49], are known to promote Th2-mediated allergic asthma [50, 51]. Thus, induction of CYP27B1 and CYP24A1 could represent a common mechanism to increase allergic sensitization after oxidative stress responses in mucosal tissues. Taken together, our findings reveal a new mechanism for the promotion of IgE responses by environmental pollutants (Fig. 7). These insights also have important implications for the prevention and treatment of allergic diseases since they identify CYP27B1 and CYP24A1 as potential targets for the prevention of allergic sensitization.

MATERIALS AND METHODS

Mice

SPF C57BL/6 mice (Jackson Laboratory (Bar Harbor, ME)) were maintained at the Ohio State University animal care facility. Germ-free C57BL/6 mice were obtained by cesarean derivation and maintained in sterile isolators. The Verigem IgE reporter (IgE Verigem) mice were obtained from Dr. Christopher Allen (the University of California at San Francisco). All animal experiments were approved by the OSU Animal Care and Use Committee.

Exposure of mice to subtoxic doses of Cd in vivo

Mice aged 8–12 weeks received cadmium chloride (CdCl $_2$, MW = 183.3; Sigma-Aldrich, St. Louis, MO) in drinking water for 4 weeks. Cd was given at the environmentally relevant doses of 10 μM (Cd10) or 25 μM (Cd25) (equivalent to 2 or 5 ppm ($\mu\text{g/L}$)), based on previous studies [10].

Allergen sensitization and allergen challenge

Mice were sensitized orally on days 0 and 7 by intragastric gavage of 1 mg of ovalbumin (OVA) and 15 μg CT in 250 μL of phosphate-buffered saline (PBS). Blood samples were collected on day 14 for analysis of serum IgE and other immunoglobulin isotypes. Nasal antigen challenges (200 μg of OVA in PBS 100 μL) were performed on days 15, 16, and 19 on mice anesthetized by intraperitoneal injection of ketamine/xylazine. The hypothermia associated with allergic responses was measured on the skin with a digital thermometer (Heat Spy infrared thermal imaging camera, Wahl, Culver City, CA).

In vivo treatment with pharmacological inhibitors of specific pathways

To address the role of oxidative stress, PGE $_2$, and Vitamin D $_3$ -metabolizing enzymes, mice received orally by intragastric gavage 250 μL of saline containing 250 mg/kg of N-acetyl L-cysteine, 25 mg/kg of aspirin, or 10 mg/kg of ketoconazole (Sigma-Aldrich, Saint Louis, MO).

Statistical analyses

Results are expressed as mean \pm SD. Statistical significance was determined by one- or two-way ANOVA, followed by Tukey's multiple range test. All statistical analyses were performed with the StataSE 12.0 software (StataCorp LLC, College Station, TX) and Prism 7 software (GraphPad Software, La Jolla, CA).

Additional methods

Additional methods are in the Supplemental Information which includes 2 tables and 5 figures.

REFERENCES

- Hong, X., Tsai, H. J. & Wang, X. Genetics of food allergy. *Curr. Opin. Pediatr.* **21**, 770–6 (2009).
- Liu, A. H. Revisiting the hygiene hypothesis for allergy and asthma. *J. Allergy Clin. Immunol.* **136**, 860–5 (2015).
- Wesemann, D. R. & Nagler, C. R. The microbiome, timing, and barrier function in the context of allergic disease. *Immunity* **44**, 728–38 (2016).
- Park, S., Lee, E. H. & Kho, Y. The association of asthma, total IgE, and blood lead and cadmium levels. *J. Allergy Clin. Immunol.* **138**, 1701–3 e6 (2016).
- Yang, S. N., Hsieh, C. C., Kuo, H. F., Lee, M. S., Huang, M. Y. & Kuo, C. H. et al. The effects of environmental toxins on allergic inflammation. *Allergy Asthma Immunol. Res.* **6**, 478–84 (2014).
- Faroon O, Ashizawa A, Wright S, Tucker P, Jenkins K, Ingerman L, et al. *Toxicological Profile for Cadmium*. Atlanta (GA); 2012.
- Jarup, L. & Akesson, A. Current status of cadmium as an environmental health problem. *Toxicol. Appl. Pharm.* **238**, 201–8 (2009).
- Wei, B. & Yang, L. A review of heavy metal contamination in urban roads dusts and agricultural soils from China. *Microchemical J.* **94**, 99–107 (2010).
- Wang, S., Huang, D. Y., Zhu, Q. H., Zhu, H. H., Liu, S. L. & Luo, Z. C. et al. Speciation and phytoavailability of cadmium in soil treated with cadmium-contaminated rice straw. *Environ. Sci. Pollut. Res. Int.* **22**, 2679–86 (2015).
- Kundu, S., Sengupta, S., Chatterjee, S., Mitra, S. & Bhattacharyya, A. Cadmium induces lung inflammation independent of lung cell proliferation: a molecular approach. *J. Inflamm.* **6**, 19 (2009).
- Liu, Y., Li, Y., Liu, K. & Shen, J. Exposing to cadmium stress cause profound toxic effect on microbiota of the mice intestinal tract. *PLoS ONE* **9**, e85323 (2014).
- Wu, K. C., Liu, J. J. & Klaassen, C. D. Nrf2 activation prevents cadmium-induced acute liver injury. *Toxicol. Appl. Pharmacol.* **263**, 14–20 (2012).
- Hogervorst, J., Plusquin, M., Vangronsveld, J., Nawrot, T., Cuypers, A. & Van Hecke, E. et al. House dust as possible route of environmental exposure to cadmium and lead in the adult general population. *Environ. Res.* **103**, 30–7 (2007).
- Nawrot, T., Plusquin, M., Hogervorst, J., Roels, H. A., Celis, H. & Thijs, L. et al. Environmental exposure to cadmium and risk of cancer: a prospective population-based study. *Lancet Oncol.* **7**, 119–26 (2006).
- Ke, S., Cheng, X. Y., Zhang, J. Y., Jia, W. J., Li, H. & Luo, H. F. et al. Estimation of the benchmark dose of urinary cadmium as the reference level for renal dysfunction: a large sample study in five cadmium polluted areas in China. *BMC Public Health* **15**, 656 (2015).
- Hu, Y. P., Peng, Y. B., Zhang, Y. F., Wang, Y., Yu, W. R. & Yao, M. et al. Reactive oxygen species mediated prostaglandin E2 contributes to acute response of epithelial injury. *Oxid. Med. Cell Longev.* **2017**, 4123854 (2017).
- Gao, Y., Zhao, C., Wang, W., Jin, R., Li, Q. & Ge, Q. et al. Prostaglandins E2 signal mediated by receptor subtype EP2 promotes IgE production in vivo and contributes to asthma development. *Sci. Rep.* **6**, 20505 (2016).
- Kuroda, E., Ishii, K. J., Uematsu, S., Ohata, K., Coban, C. & Akira, S. et al. Silica crystals and aluminum salts regulate the production of prostaglandin in macrophages via NALP3 inflammasome-independent mechanisms. *Immunity* **34**, 514–26 (2011).
- Hapfelmeier, S., Lawson, M. A., Slack, E., Kirundi, J. K., Stoel, M. & Heikenwalder, M. et al. Reversible microbial colonization of germ-free mice reveals the dynamics of IgA immune responses. *Science* **328**, 1705–9 (2010).
- Planer, J. D., Peng, Y., Kau, A. L., Blanton, L. V., Ndao, I. M. & Tarr, P. I. et al. Development of the gut microbiota and mucosal IgA responses in twins and gnotobiotic mice. *Nature* **534**, 263–6 (2016).
- Wu, W., Sun, M., Chen, F., Cao, A. T., Liu, H. & Zhao, Y. et al. Microbiota metabolite short-chain fatty acid acetate promotes intestinal IgA response to microbiota which is mediated by GPR43. *Mucosal Immunol.* **10**, 946–56 (2017).
- Zhang, X. Y., Liu, Z. M., Zhang, H. F., Li, Y. S., Wen, S. H. & Shen, J. T. et al. TGF- β 1 improves mucosal IgA dysfunction and dysbiosis following intestinal ischaemia-reperfusion in mice. *J. Cell Mol. Med.* **20**, 1014–23 (2016).
- Shan, M., Gentile, M., Yeiser, J. R., Walland, A. C., Bornstein, V. U. & Chen, K. et al. Mucus enhances gut homeostasis and oral tolerance by delivering immunoregulatory signals. *Science* **342**, 447–53 (2013).
- D'Mello, R. J., Caldwell, J. M., Azouz, N. P., Wen, T., Sherrill, J. D. & Hogan, S. P. et al. LRRc31 is induced by IL-13 and regulates kallikrein expression and barrier function in the esophageal epithelium. *Mucosal Immunol.* **9**, 744–56 (2016).
- Jones, S. A., Schneider, H. & Challis, J. R. Vitamin D3 metabolites stimulate prostaglandin production by human fetal membranes and placenta in vitro. *Am. J. Perinatol.* **6**, 138–41 (1989).
- Castigli, E., Wilson, S. A., Scott, S., Dedeoglu, F., Xu, S. & Lam, K. P. et al. TAC1 and BAFF-R mediate isotype switching in B cells. *J. Exp. Med.* **201**, 35–9 (2005).
- Castigli, E., Scott, S., Dedeoglu, F., Bryce, P., Jabara, H. & Bhan, A. K. et al. Impaired IgA class switching in APRIL-deficient mice. *Proc. Natl Acad. Sci. USA* **101**, 3903–8 (2004).
- He, B., Santamaria, R., Xu, W., Cols, M., Chen, K. & Puga, I. et al. The transmembrane activator TAC1 triggers immunoglobulin class switching by activating B cells through the adaptor MyD88. *Nat. Immunol.* **11**, 836–45 (2010).
- Kim, Y. G., Udayanga, K. G., Totsuka, N., Weinberg, J. B., Nunez, G. & Shibuya, A. Gut dysbiosis promotes M2 macrophage polarization and allergic airway inflammation via fungi-induced PGE(2). *Cell Host Microbe* **15**, 95–102 (2014).
- Jacobs, E. T., Van Pelt, C., Forster, R. E., Zaidi, W., Hibler, E. A. & Galligan, M. A. et al. CYP24A1 and CYP27B1 polymorphisms modulate vitamin D metabolism in colon cancer cells. *Cancer Res.* **73**, 2563–73 (2013).
- Kundu, R., Chain, B. M., Coussens, A. K., Khoo, B. & Noursadeghi, M. Regulation of CYP27B1 and CYP24A1 hydroxylases limits cell-autonomous activation of vitamin D in dendritic cells. *Eur. J. Immunol.* **44**, 1781–90 (2014).
- Liu, X., Nelson, A., Wang, X., Farid, M., Gunji, Y. & Ikari, J. et al. Vitamin D modulates prostaglandin E2 synthesis and degradation in human lung fibroblasts. *Am. J. Respir. Cell Mol. Biol.* **50**, 40–50 (2014).
- Greer, F. R., Sicherer, S. H. & Burks, A. W. American Academy of Pediatrics Committee on N, American Academy of Pediatrics Section on A, and Immunology Effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, timing of introduction of complementary foods, and hydrolyzed formulas. *Pediatrics* **121**, 183–91 (2008).
- Geha, R. S., Jabara, H. H. & Brodeur, S. R. The regulation of immunoglobulin E class-switch recombination. *Nat. Rev. Immunol.* **3**, 721–32 (2003).
- Jelovcan, S., Gutsch, A., Kleinhappl, B., Sedlmayr, P., Barth, S. & Marth, E. Effects of low concentrations of cadmium on immunoglobulin E production by human B lymphocytes in vitro. *Toxicology* **188**, 35–48 (2003).
- Hanson, M. L., Holaskova, I., Elliott, M., Brundage, K. M., Schafer, R. & Barnett, J. B. Prenatal cadmium exposure alters postnatal immune cell development and function. *Toxicol. Appl. Pharmacol.* **261**, 196–203 (2012).
- Ba Q, Li M, Chen P, Huang C, Duan X, Lu L, et al. Gender-dependent effects of cadmium exposure in early life on gut microbiota and fat accumulation in mice. *Environ. Health Perspect.* 2016.
- Grasberger, H., Gao, J., Nagao-Kitamoto, H., Kitamoto, S., Zhang, M. & Kamada, N. et al. Increased expression of DUOX2 is an epithelial response to mucosal dysbiosis required for immune homeostasis in mouse intestine. *Gastroenterology* **149**, 1849–59 (2015).
- Sommer, F. & Backhed, F. The gut microbiota engages different signaling pathways to induce Duox2 expression in the ileum and colon epithelium. *Mucosal Immunol.* **8**, 372–9 (2015).
- Wang, C., Guan, Y., Lv, M., Zhang, R., Guo, Z. & Wei, X. et al. Manganese increases the sensitivity of the cGAS-STING pathway for double-stranded DNA and is required for the host defense against DNA viruses. *Immunity* **48**, 675–87 e7 (2018).
- Breton, J., Massart, S., Vandamme, P., De Brandt, E., Pot, B. & Foligne, B. Ecotoxicology inside the gut: impact of heavy metals on the mouse microbiome. *BMC Pharmacol. Toxicol.* **14**, 62 (2013).
- Bolcas, P. E., Brandt, E. B., Zhang, Z., Biagini Myers, J. M., Ruff, B. P. & Khurana Hershey, G. K. Vitamin D supplementation attenuates asthma development following traffic-related particulate matter exposure. *J. Allergy Clin. Immunol.* **143**, 386–94 e3 (2019).
- Hollams, E. M., Hart, P. H., Holt, B. J., Serralha, M., Parsons, F. & de Klerk, N. H. et al. Vitamin D and atopy and asthma phenotypes in children: a longitudinal cohort study. *Eur. Respir. J.* **38**, 1320–7 (2011).
- Hufnagl, K. & Jensen-Jarolim, E. Vitamin A and D in allergy: from experimental animal models and cellular studies to human disease. *Allergo J. Int.* **27**, 72–8 (2018).
- Cantorna, M. T., Snyder, L., Lin, Y. D. & Yang, L. Vitamin D and 1,25(OH)₂D regulation of T cells. *Nutrients* **7**, 3011–21 (2015).
- Ruiter, B., Patil, S. U. & Shreffler, W. G. Vitamins A and D have antagonistic effects on expression of effector cytokines and gut-homing integrin in human innate lymphoid cells. *Clin. Exp. Allergy* **45**, 1214–25 (2015).
- Chambers, E. S., Suwannasaen, D., Mann, E. H., Urry, Z., Richards, D. F. & Lertmongsolkolchai, G. et al. 1 α ,25-dihydroxyvitamin D3 in combination with transforming growth factor- β increases the frequency of Foxp3(+) regulatory

- T cells through preferential expansion and usage of interleukin-2. *Immunology* **143**, 52–60 (2014).
48. Rosendahl, J., Pelkonen, A. S., Helve, O., Hauta-Alus, H., Holmlund-Suila, E. & Valkama, S. et al. High-dose vitamin D supplementation does not prevent allergic sensitization of infants. *J. Pediatr.* **209**, 139–45 e1 (2019).
49. Lodovici, M. & Bigagli, E. Oxidative stress and air pollution exposure. *J. Toxicol.* **2011**, 487074 (2011).
50. Brandt, E. B., Biagini Myers, J. M., Acciani, T. H., Ryan, P. H., Sivaprasad, U. & Ruff, B. et al. Exposure to allergen and diesel exhaust particles potentiates secondary allergen-specific memory responses, promoting asthma susceptibility. *J. Allergy Clin. Immunol.* **136**, 295–303 e7 (2015).
51. De Grove, K. C., Provoost, S., Hendriks, R. W., McKenzie, A. N. J., Seys, L. J. M. & Kumar, S. et al. Dysregulation of type 2 innate lymphoid cells and TH2 cells impairs pollutant-induced allergic airway responses. *J. Allergy Clin. Immunol.* **139**, 246–57 e4 (2017).

ACKNOWLEDGEMENTS

This work was supported by NIH grants R01AI043197, R01DK101323, and R01AI145144, and the UL1TR001070 award from the National Center for Advancing Translational Sciences. The authors thank Arpad Somogyi and Matthew Bernier of the OSU Center for Clinical and Translational Science Mass Spectrometry & Proteomics Core for assistance with metabolomics and proteomics studies. The Mass Spectrometry & Proteomics Core was supported by NIH grant P30 CA016058 and the Fusion Orbitrap instrument was supported by NIH grant S10 OD018056.

AUTHOR CONTRIBUTIONS

E.K. conducted the experiments, analyzed the data, and wrote the manuscript. A.B., M.R.J., and Z.A. conducted the experiments and analyzed the data. S.O. performed the statistical analyses. B.H.A. designed the experiments and reviewed the manuscript. E.C. designed the experiments, analyzed the data, and reviewed the manuscript. P.N.B. designed the experiments, analyzed the data, and wrote the manuscript. Current address for AB: Zoetis, Louvain-la-Neuve, Belgium. Current address for ZA: Nationwide Children Hospital, Columbus, Ohio.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Correspondence and requests for materials should be addressed to Prosper N. Boyaka.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.