



## ARTICLE

# Increased functional coupling of the mu opioid receptor in the anterior insula of depressed individuals

Pierre-Eric Lutz<sup>1,6</sup>, Daniel Almeida<sup>1</sup>, Dominique Filliol<sup>2</sup>, Fabrice Jollant<sup>1,3,4,7</sup>, Brigitte L. Kieffer<sup>1,2,5,8</sup> and Gustavo Turecki<sup>1</sup>

The mu opioid receptor (MOR) is a G protein-coupled receptor that plays an essential role in reward and hedonic processes, and that has been implicated in disorders such as depression and addiction. Over the last decade, several brain imaging studies in depressed patients have consistently found that dysregulation of MOR function occurs in particular in the anterior insular cortex, an important brain site for the perception of internal states and emotional regulation. To investigate molecular mechanisms that may underlie these effects, here we assessed genetic polymorphisms, expression, and functional G-protein coupling of MOR in a large post-mortem cohort ( $N = 95$ ) composed of depressed individuals who died by suicide, and healthy controls. Results indicated that depression, but not comorbid substance use disorder or acute opiate consumption, was associated with increased MOR activity. This effect was partly explained by a specific increase in expression of the inhibitory alpha G-protein subunit GNAI2. Consistent with previous neuroimaging studies, our findings support the notion that enhanced endogenous opioidergic tone in the anterior insula may buffer negative affective states in depressed individuals, a mechanism that could potentially contribute to the antidepressant efficacy of emerging opioid-based medications.

*Neuropsychopharmacology* (2021) 46:920–927; <https://doi.org/10.1038/s41386-021-00974-y>

## INTRODUCTION

Depression is a recurring and often chronic brain disorder that stems from maladaptive processes affecting multiple neurotransmitter systems. While the monoaminergic theory of depression received considerable attention, and fueled intense research efforts over the last 50 years, another theory, dating back to the early 1900s, posits the implication of the opioid system in regulating mood [1–3]. Euphoric properties of opiates acting at the mu opioid receptor (MOR), such as morphine and heroin, have been recreationally exploited for centuries, and were once considered a potential ‘opioid cure’ for depression.

Over the last two decades, both clinical and animal researches have further strengthened the notion that adaptations at the level of MOR signaling may contribute to depression pathophysiology [1]. Recently, functional neuroimaging has allowed studies of MOR in patients experiencing various mood states [2]. Positron emission tomography (PET) with the MOR selective radiotracer [<sup>11</sup>C]-carfentanil found that the induction of a sadness state, or of sustained pain, was associated in both healthy [3, 4] and depressed [5] women with changes in MOR bioavailability across several brain regions. Among these regions, the insular cortex [6] appears as a major site where MOR modulates affective states, in particular in relation to social experiences, such as social acceptance and rejection, in both healthy volunteers [7] and depressed [8] subjects.

While the PET approach allows studying MOR availability in living individuals, it does not allow to differentiate, at baseline, between variation in receptor concentration and variation in occupancy by endogenous opioid peptides (which requires an additional releasing challenge [9]), complicating interpretation of findings. Since MOR belongs to the large family of inhibitory G protein-coupled receptors (GPCR), radioligand binding assays in post-mortem cohorts offer a complementary approach to assess receptor level and signaling. The five previous studies available for MOR in relation to depression were conducted between 1990 and 2012 [10–14], focused on relatively small cohorts, and investigated the temporal and frontal cortices, caudate nucleus, and thalamus. Importantly, no investigation was conducted on the cortical structure most consistently identified by the aforementioned brain imaging studies, the insular cortex.

In this context, the goal of the present work was to test the hypothesis that major depressive disorder (MDD) may associate with dysregulated MOR function in the anterior insular cortex (AI). Using post-mortem samples from a large and well-characterized cohort ( $N = 95$  total), we compared subjects diagnosed with MDD ( $N = 55$ ), and healthy controls ( $N = 40$ ). Expression of MOR was measured using quantitative PCR, while MOR coupling was assessed using agonist-stimulated [<sup>35</sup>S]-GTPγS binding, and the expression of Gi proteins measured using Nanostring. In addition, a series of single nucleotide

<sup>1</sup>McGill Group for Suicide Studies, Douglas Mental Health Research Centre, McGill University, 6875 LaSalle Boulevard, Verdun, Montréal, QC, Canada; <sup>2</sup>Translational Medicine and Neurogenetics, Institut de Génétique et de Biologie Moléculaire et Cellulaire, INSERM U-964, CNRS UMR-7104, Université de Strasbourg, Illkirch, France; <sup>3</sup>Université de Paris, Paris, France; <sup>4</sup>CHU Nîmes, Nîmes, France and <sup>5</sup>Douglas Mental Health Research Centre, Department of Psychiatry, McGill University, Montréal, QC H4H 1R3, Canada  
Correspondence: Pierre-Eric Lutz ([pierreeric.lutz@gmail.com](mailto:pierreeric.lutz@gmail.com)) or Gustavo Turecki ([gustavo.turecki@mcgill.ca](mailto:gustavo.turecki@mcgill.ca))

<sup>6</sup>Present address: Centre National de la Recherche Scientifique, Université de Strasbourg, Fédération de Médecine Translationnelle de Strasbourg, Institut des Neurosciences Cellulaires et Intégratives UPR3212, Strasbourg, France

<sup>7</sup>Present address: Clinique des Maladies Mentales et de l'Encéphale (CMME), CH Sainte-Anne, GHU Paris, Psychiatrie et Neurosciences, Paris, France

<sup>8</sup>Present address: INSERM U1114, Department of Psychiatry, University Hospital of Strasbourg, Strasbourg, France

Received: 4 October 2020 Accepted: 14 January 2021

Published online: 2 February 2021

polymorphisms located along the MOR gene and potentially contributing to variability in receptor expression or function were characterized. Overall, our results indicate that MDD associates in the anterior insula with increased MOR coupling, which appears partly mediated by increased expression of the alpha(i2) G-protein subunit, suggesting a molecular substrate that may contribute to disease pathophysiology.

## MATERIAL AND METHODS

### Cohort

This study was approved by the Research Ethics Board at the Douglas Mental Health University Institute, Verdun, Quebec. Brain tissue was obtained from the Douglas-Bell Canada Brain Bank (<http://douglasbrainbank.ca/>). We compared two groups of subjects matched for sex, age, post-mortem interval (PMI), pH and RNA Integrity Numbers (RIN; see Table 1): (1) One group of individuals who died by suicide during a major depressive episode (MDD group), and (2) A second group of psychiatrically healthy individuals who died suddenly from accidental causes, and who constituted the control group (C). Information on psychopathology, socio-demographic variables and development was obtained by way of psychological autopsies performed by trained clinicians, with the informants best acquainted with the deceased, as described elsewhere [15]. The process employed by our group in the collection of information used in psychological autopsies has been extensively investigated and found to produce valid information, especially in the context of observable behaviors and major life events [16–18]. Both cases and controls were characterized by the same psychological autopsy methods, avoiding the occurrence of systematic biases. Psychopathology diagnoses were obtained using DSM-IV criteria by means of SCID-I interviews adapted for psychological autopsies. Brain samples were dissected from each subject from 0.5 cm-thick coronal brain sections by expert brain bank staff following standard dissection procedures, with the aid of a human brain atlas (see [19] and [thehumanbrain.info/brain/bn\\_brain\\_atlas/brain.html](http://thehumanbrain.info/brain/bn_brain_atlas/brain.html)). Samples were carefully dissected at 4 °C after having been flash-frozen in isopentane at –80 °C. AI tissue was dissected from the superior part of the insula, just beneath the overlying opercula in the brain section containing the striatum, at the level of nucleus accumbens.

### Relationship to previous studies

Our previous study on a partly overlapping but smaller cohort ( $N=59$ ) focused on the kappa opioid receptor [20] and its epigenetic regulation, while also reporting mRNA expression for the MOR. The present study investigated an enlarged cohort

( $N=95$ ), and comprehensively characterized MOR for gene expression, genetic variation, and signaling efficiency.

### Quantitative polymerase chain reaction (qPCR)

RNA was extracted from homogenized brain tissue samples using the RNeasy Lipid Tissue Mini Kit (Qiagen). RNA quantity and quality were measured by Nanodrop® and Agilent 2100 Bioanalyzer technologies, and only samples with an RNA integrity number (RIN)  $\geq 5$  were used (resulting in  $N=40$  C and  $N=55$  MDD subjects). Extracted RNA was reverse-transcribed using M-MLV reverse transcriptase (Invitrogen™). Messenger RNA (mRNA) levels were quantified by real time polymerase chain reaction (RT-PCR) using SYBR® Green DNA intercalating dye and master mix (Bio-Rad) and the ABI 7900HT PCR machine. Primers for targeted genes were designed using Primer-BLAST (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>) and validated by gel migration and dissociation curves. Complementary DNAs (cDNA) from every subject were pooled and used to prepare calibration curves, from which cDNA quantities of target genes were calculated in each sample according to the measured RT-PCR threshold cycle (Ct value), in quintuplicates. Relative expression levels for each gene of interest were calculated by dividing the cDNA quantity for the gene of interest by the arithmetic mean of cDNA quantities for two reference housekeeping genes, GAPDH and  $\beta$ -actin. MOR, forward primer: TTC CGT ACT CCC CGA AAT GC; reverse primer: ACA ATC TAT GGA ACC TTG CCT G. GAPDH, forward primer: TTG TCA AGC TCA TTT CCT GG; reverse primer: TGT GAG GAG GGG AGA TTC AG.  $\beta$ -actin, forward primer: AAG ACC TGT ACG CCA ACA CA; reverse primer: GCA GTG ATC TCC TTC TGC ATC.

### Genotyping

Genomic DNA extracted from brain tissue was first amplified, via PCR, using a Platinum Taq DNA Polymerase (Invitrogen®). Forward and reverse primers were designed using NCBI's Primer-BLAST tool and synthesized by Integrated DNA Technologies (IDT). Final reactions contained 1  $\mu$ L of gDNA, 1  $\mu$ L of 10x PCR buffer, 0.2  $\mu$ L of 10 mM dNTP mix, 0.6  $\mu$ L of 50 mM MgCl<sub>2</sub>, 0.1  $\mu$ L of Q-solution, 1  $\mu$ L of 10 nM forward and reverse primers, 0.1  $\mu$ L of DNA Polymerase, and 6  $\mu$ L of H<sub>2</sub>O. PCR reactions were carried out using the following conditions: 95 °C for 5 min, then 35 cycles of [95 °C for 30 s, 60 °C for 30 s, 72 °C for 40 s], final extension at 72 °C for 7 min. Individual amplicons were then purified using a 0.7x AMPure bead mix. Purified amplicons were sent for Sanger sequencing, on a 3730xl DNA Analyzer (Applied Biosystems), at the McGill University and Genome Quebec Innovation Center. Sequence trace files in ABI format, as well as reference sequences in FASTA format, were loaded into the novoSNP 3.0.1 program for Windows users [21]. Variation at each location was determined by following the procedures outlined in Weckx et al. [21].

### Agonist-stimulated [<sup>35</sup>S]-GTP $\gamma$ S binding activity of MOR

The functional activity of MOR was assessed by the [<sup>35</sup>S]-GTP $\gamma$ S binding assay, as described previously [22]. The method is based on the ability of the radiolabeled GTP analog, [<sup>35</sup>S]-GTP $\gamma$ S, to specifically bind to G-proteins. Here, 100 mg of AI tissue was used for each subject for membrane preparation. Membranes were prepared by homogenizing the tissue in ice-cold 0.25 M sucrose solution 10 vol (ml/g wet weight of tissue). Samples were then centrifuged at 1100  $\times$  g for 10 min. Supernatants were collected and diluted 5 times in buffer containing 50 mM Tris-HCl, pH 7.4, and 1 mM EDTA, and were then centrifuged at 35,000  $\times$  g for 30 min. The pellets were homogenized in 2 ml of ice-cold sucrose solution (0.32 M), aliquoted and kept at –80 °C until further use. For each [<sup>35</sup>S]-GTP $\gamma$ S binding assay, 2.5  $\mu$ g of membrane proteins were used per well. Samples were incubated with the MOR agonist DAMGO (using 12 increasing concentrations, from 10<sup>–4</sup> to 10<sup>–10</sup> M) for 1 h at 25 °C in assay buffer 50 mM Tris-HCl (pH 7.4), 3 mM MgCl<sub>2</sub>, 100 mM NaCl, 0.2 mM EGTA containing 30  $\mu$ M GDP

**Table 1.** Sample demographics.

	Controls (C)	Major depressive disorder (MDD)
<i>N</i>	40	55
Sex, M/F (%)	32 / 8 (80.0 / 20.0%)	45 / 10 (81.8 / 18.2%)
Age (years)	45.65 $\pm$ 2.98	45.73 $\pm$ 2.15
RNA integrity number (RIN)	6.89 $\pm$ 0.16	7.15 $\pm$ 0.14
PMI (hours)	27.3 $\pm$ 3.52	25.8 $\pm$ 2.51
pH	6.44 $\pm$ 0.05	6.56 $\pm$ 0.04
Alcohol/Drug Abuse	14 (35.0%)	20 (36.4%)

Healthy controls (C) and subjects who died by suicide during a major depressive episode (MDD) were matched for sex, age, post-mortem interval (PMI), and pH. There were no differences, between groups, in the number of male and female subjects, the presence of alcohol or drug abuse, or RIN values (see main text). Age, RIN, PMI and pH values are shown as mean  $\pm$  sem.

and 0.1 nM [<sup>35</sup>S]-GTPγS. Incubation was terminated by rapid filtration and washing in ice-cold buffer (50 mM Tris-HCl, 5 mM MgCl<sub>2</sub>, 50 mM NaCl, pH 7.4). Bound radioactivity was quantified using a liquid scintillation counter (Perkin Elmer Top Count). Non-specific binding was defined as binding in the presence of 10 μM GTPγS, and basal binding was assessed in the absence of DAMGO. Assays were performed in triplicates in different membrane preparations. Non-linear regressions were performed on individual concentration-response curves of DAMGO-induced stimulations (Prism, GraphPad Software, San Diego, CA, USA). Data are expressed as the maximal net stimulation ( $E_{max}$ , stimulated minus basal levels) expressed in fmol.mg<sup>-1</sup> protein. The concentration that elicited half-maximal effect ( $EC_{50}$ ) was obtained and normalized as log  $EC_{50}$  values.

#### Quantification of inhibitory G-proteins alpha subunits expression using Nanostring

Nanostring experiments were performed as described previously [23] at the Jewish General Hospital Molecular Pathology Centre (Montréal, QC, Canada) using Nanostring nCounter targeted gene expression profiling [24]. Briefly, 5 μl of total RNA (20 ng/μl) was hybridized with the reporter and capture probes at 65 °C in a thermocycler for 19–20 h. Probes were designed against 4 genes of interest (GNAI1, GNAI2, GNAO1 and GNAZ) and against 4 housekeeping control genes highly expressed in the AI: GAPDH, β-Actin, Tubulin and β-2-Globulin. The samples were processed with the nCounter Prep Station to purify the hybridized targets and affix them to the cartridge. After transfer to the nCounter Digital Analyzer, barcodes were counted and tabulated for each target molecule. The data were analyzed using the nSolver version 2.6.43 following the manufacturer's recommendations.

#### Statistical analysis


Statistical analyses were carried out using R and Prism v8.4.1. Frequencies of SNPs were analyzed using  $\chi^2$  test to test for

Hardy-Weinberg equilibrium. For MOR expression and [<sup>35</sup>S]-GTPγS binding, a general linear model (GLM) was used to analyze group differences. Potential correlations between variables that commonly act as confounders (pH, PMI, RIN, sex and age) and gene expression or [<sup>35</sup>S]-GTPγS binding were assessed separately, and only those variables showing a significant Pearson correlation were included in each final GLM model. Statistical significance was set at  $p < 0.05$ , and Bonferroni multiple testing corrections were applied to take into account the number of SNPs investigated in MOR genotyping experiments ( $n = 16$ ), and the number of Gi alpha-subunit genes analyzed using Nanostring ( $n = 4$ ).

## RESULTS

### MOR gene expression in the anterior insula

We first used qPCR to quantify MOR mRNA levels in the two groups of C and MDD subjects. Groups did not differ with respect to the subjects' sex (Fisher's exact test,  $p > 0.99$ ) nor RIN ( $t = 1.25$ ,  $p = 0.21$ ; see Table 1). MOR expression was similar across men and women ( $t = 1.30$ ,  $p = 0.20$ ) and, as expected, significantly correlated with RIN values (Fig. S1A). Interestingly, it also increased with age (Fig. 1A;  $r = 0.37$ ,  $N = 90$ ,  $p = 3E-04$ ), in contrast with previous studies that reported a significant decrease in the medio-dorsal thalamus [20], with no changes in three other structures (the caudal part of the anterior cingulate cortex [20], the ventral posterior part of the lateral thalamus [25], and the secondary somato-sensory cortex [25]). This could indicate brain region-specific patterns of MOR regulation during aging. Importantly, a GLM comparing our two main groups, C and MDD, and considering age and RIN as significant covariates, did not reveal a difference between groups (Fig. 1B;  $F(1,86) = 1.11$ ,  $p = 0.27$ ). Further, while we previously reported that a history of child abuse, a major distal risk factor for MDD, associated in the AI with decreased expression of the kappa opioid receptor [20], no such effect was found for MOR ( $t = 0.83$ ,  $p = 0.41$ ).



**Fig. 1** Relative real-time quantitative PCR (RT-PCR) expression of the mu opioid receptor (MOR) in the anterior insula by age and clinical group. **A** MOR expression was quantified by RT-PCR in quintuplicates, using calibration curves and two reference housekeeping genes: GAPDH and β-actin (see methods for details). The scatter plot displays a significant positive linear correlation between MOR expression and age, in all subjects, independent of clinical group ( $r = 0.37$ ,  $p = 3E-04$ ). **B** Expression of MOR in healthy controls and subjects who died during a major depressive episode (MDD). There was no significant difference between the two clinical groups ( $F(1,86) = 1.11$ ,  $p = 0.27$ ). Mean ± sem, as well as individual expression values, are shown.

**Table 2.** Single nucleotide polymorphisms (SNP) profiled in this study in the *OPRM1* gene encoding the mu opioid receptor.

SNP	Position	Feature	Sample count major/minor	Sample frequency major/minor (%)	1000 genomes frequency major/minor (%)
rs147391824	chr6:154,348,457	Promoter	<b>A</b> (179)/ <b>C</b> (7)	<b>A</b> (96.2)/ <b>C</b> (3.8)	<i>A</i> (98.5)/ <i>C</i> (1.5)
rs7751510	chr6:154,348,488	Promoter	<b>G</b> (176)/ <b>C</b> (10)	<b>G</b> (94.6)/ <b>C</b> (5.4)	<i>G</i> (89.3)/ <i>C</i> (10.7)
rs117514803	chr6:154,348,559	Promoter	<b>C</b> (172)/ <b>T</b> (14)	<b>C</b> (92.5)/ <b>T</b> (7.5)	<i>C</i> (98.0)/ <i>T</i> (2.0)
rs7776341	chr6:154,348,605	Promoter	<b>A</b> (177)/ <b>C</b> (9)	<b>A</b> (95.2)/ <b>C</b> (4.8)	<i>A</i> (89.4)/ <i>C</i> (10.6)
rs1074287	chr6:154,348,809	Promoter	<b>A</b> (137)/ <b>G</b> (49)	<b>A</b> (73.6)/ <b>G</b> (26.4)	<i>A</i> (71.8)/ <i>G</i> (28.2)
rs60509363	chr6:154,348,978	Promoter	<b>G</b> (177)/ <b>A</b> (9)	<b>G</b> (95.2)/ <b>A</b> (4.8)	<i>G</i> (85.9)/ <i>A</i> (14.1)
rs17181003	chr6:154,360,444	Exon 1	<b>A</b> (184)/ <b>G</b> (0)	<b>A</b> (100)/ <b>G</b> (0)	<i>A</i> (98.9)/ <i>G</i> (1.1)
rs6912029	chr6:154,360,508	Exon 1	<b>G</b> (175)/ <b>T</b> (9)	<b>G</b> (95.1)/ <b>T</b> (4.9)	<i>G</i> (92.7)/ <i>T</i> (7.3)
rs17174638	chr6:154,360,569	Exon 1	<b>C</b> (184)/ <b>T</b> (0)	<b>C</b> (100)/ <b>T</b> (0)	<i>C</i> (96.7)/ <i>T</i> (3.3)
rs1799972	chr6:154,360,696	Exon 1	<b>C</b> (184)/ <b>T</b> (0)	<b>C</b> (100)/ <b>T</b> (0)	<i>C</i> (92.9)/ <i>T</i> (7.1)
rs1799971	chr6:154,360,797	Exon 1	<b>A</b> (159)/ <b>G</b> (25)	<b>A</b> (86.4)/ <b>G</b> (13.6)	<i>A</i> (77.7)/ <i>G</i> (22.3)
rs12209447	chr6:154,361,048	Intron 1	<b>C</b> (175)/ <b>T</b> (9)	<b>C</b> (95.1)/ <b>T</b> (4.9)	<i>C</i> (96.0)/ <i>T</i> (4.0)
rs9479757	chr6:154,411,344	Intron 2	<b>G</b> (164)/ <b>A</b> (16)	<b>G</b> (91.1)/ <b>A</b> (8.9)	<i>G</i> (92.0)/ <i>A</i> (8.0)
rs17181199	chr6:154,411,710	Intron 2	<b>T</b> (184)/ <b>A</b> (0)	<b>T</b> (100)/ <b>A</b> (0)	<i>T</i> (98.6)/ <i>A</i> (1.4)
rs17181213	chr6:154,411,847	Intron 2	<b>C</b> (186)/ <b>T</b> (0)	<b>C</b> (100)/ <b>T</b> (0)	<i>C</i> (98.5)/ <i>T</i> (1.5)
rs2075572	chr6:154,412,004	Intron 2	<b>C</b> (113)/ <b>G</b> (67)	<b>C</b> (62.8)/ <b>G</b> (37.2)	<i>C</i> (61.2)/ <i>G</i> (38.8)

SNPs were determined using Sanger sequencing (see methods). Genomic position (in the Human GRCh37/hg19 reference genome), gene feature (promoter/exon/intron), observed counts and percentages of major and minor alleles in our cohort, and expected percentages (according to the 1000 Genomes Project) are shown.


All drugs of abuse, including opiates, trigger rewarding effects at least in part by recruiting the MOR [26], which has been conceptualized as a molecular gateway to addiction [27]. Also, animal studies have reported changes in MOR mRNA expression following acute or chronic receptor activation (with mostly increased expression, but also inconsistent and even opposite effects across distinct brain regions, MOR splicing isoforms, sex, or time-points following opiate exposure [28–30]). We therefore explored whether MOR expression varied as a function of comorbid substance use disorder (SUD), or ante-mortem acute opiate consumption. Results showed that SUD diagnosis, present at similar frequency in C and MDD groups (Fisher's exact test,  $p > 0.99$ ; Table 1), had no significant effect (Fig. S1B,  $t = 0.32$ ,  $p = 0.75$ ). Toxicological screenings (conducted on blood, urine, and ocular liquid samples) indicated that all drugs of abuse considered together (opiates, cocaine, amphetamines, ethanol, and tetrahydrocannabinol) were detected in 35 subjects, while opiates were detected in 8 subjects only. Both categorizations significantly correlated with SUD diagnosis, as expected ( $p < 0.05$ ), but, importantly, did not associate with any change in MOR expression ( $p > 0.05$ ; Fig. S1C). Altogether, MOR expression in the AI was not modulated by MDD, SUD, recent opiate consumption, or history of child abuse, consistent with our previous findings for this receptor in the anterior cingulate cortex, the mediodorsal thalamus, and a smaller cohort of AI samples [20].

MOR gene expression and single nucleotide polymorphisms (SNP) We investigated a series of SNP in the MOR gene (see observed frequencies in Table 2). These were chosen based on previous literature [25, 31, 32], and notably included the A118G (rs1799971), a well-studied SNP in relation to MOR function and SUD [33]. The distribution of the A and G alleles for A118G was in Hardy-Weinberg equilibrium (HWE), as assessed using a  $\chi^2$  test ( $\chi^2 = 0.52$ ,  $p = 0.47$ ), with a low frequency of the G allele (~13.6%), as expected in a Quebec population of European descent, and similar to previous studies [34, 35]. When results from the present work were pooled with those we obtained previously in two other brain regions (the caudal part of the anterior cingulate cortex, and medio-dorsal thalamus [20]), the A118G genotype was found to

have a moderately significant effect on MOR mRNA expression, with higher expression in carriers of the minor G allele (Fig. S2; 2-way ANOVA: genotype effect  $F(1,184) = 6.30$ ,  $p = 0.013$ ; brain region effect  $F(2,184) = 1.53$ ,  $p = 0.22$ ). Previous studies that investigated this same SNP reported no correlation with MOR expression, neither in the thalamus nor the somato-sensory [25] or dorsolateral prefrontal cortices [36], suggesting a brain region-specific impact of this SNP. For the 15 other SNPs that we investigated, no relationship with MOR expression was observed ( $p > 0.05$  for all).

#### MOR signaling efficiency

We next used [ $^{35}$ S]-GTP $\gamma$ S to determine MOR G-protein coupling. Using DAMGO as a well-validated agonist, we generated dose-response curves for each subject, which were then used to evaluate maximal receptor stimulation ( $E_{\max}$ ), and the agonist concentration eliciting half-maximal effect (log EC $_{50}$ ). First, we found a significant correlation between inter-individual variability of MOR mRNA (qPCR) and protein (MOR  $E_{\max}$ ) levels ( $r = 0.28$ ,  $p = 0.028$ ; Fig. 2A). Because it is generally considered that good correlation between mRNA and protein abundance indicates local synthesis, while lack of such correlation points towards protein transport [37], this result suggests that MOR signaling in the AI is at least partly explained by its expression from local neurons, and also provides technical validation of our [ $^{35}$ S]-GTP $\gamma$ S procedure. Second, we found that basal [ $^{35}$ S]-GTP $\gamma$ S binding (defined as the constitutive activity of all GPCRs, measured in the absence of DAMGO), was slowly decreasing as a function of age ( $r = (-0.23)$ ,  $p = 0.028$ ; Fig. S3A), consistent with previous reports [38]. MOR  $E_{\max}$  was not affected by either sex ( $p = 0.42$ ) or age ( $p = 0.31$ ), but significantly correlated with brain tissue pH ( $r = 0.34$ ,  $p = 7.2E-05$ ; Fig. S3B). Finally, we observed that MOR affinity for DAMGO, measured as the log EC $_{50}$ , showed a significant negative correlation with age ( $r = -0.31$ ,  $p = 0.0023$ ; Fig. 2B), corresponding to a leftward shift of concentration-response curves, and indicating enhanced DAMGO efficacy with aging. This is consistent with results from a previous study conducted in the prefrontal cortex (Brodmann Area 9, see [38]), although the biological significance of this effect is elusive.




**Fig. 2 Mu opioid receptor (MOR) DAMGO agonist-stimulated [<sup>35</sup>S]-GTPγS binding in the anterior insular cortex of healthy controls and subjects with major depressive disorder (MDD).** Coupling efficiency of MOR was evaluated using [<sup>35</sup>S]-GTPγS binding and 12 increasing concentrations of the potent and specific agonist DAMGO, followed by non-linear regression of concentration-response curves, to compute E<sub>max</sub> (maximal net stimulation of MOR; [<sup>35</sup>S]-GTPγS net stimulation over basal binding) and logEC<sub>50</sub> (agonist concentration for half-maximal effect, log transformed) values. **A** The scatter plot displays a significant positive linear correlation ( $r = 0.28$ ,  $p = 0.028$ ) between MOR mRNA levels (measured by RT-PCR) and protein signaling (E<sub>max</sub>). **B** Scatter plot displaying a significant negative linear correlation ( $r = (-0.31)$ ,  $p = 0.0023$ ) between age and MOR affinity for DAMGO (logEC<sub>50</sub>). **C, D** Concentration-response curves of [<sup>35</sup>S]-GTPγS stimulation by log<sub>10</sub>[DAMGO]. **C** Dose-response curve in non-carriers (broken line; AA genotype) or carriers (filled-in line; AG genotype) of the minor G allele for the MOR single-nucleotide polymorphism rs1799971 (A118G). The two genotypes did not show any significant differences in MOR E<sub>max</sub> ( $p = 0.64$ ) or logEC<sub>50</sub> ( $p = 0.97$ ) values. **D** Dose-response curve in control (broken line) or MDD (filled-in line) subjects. Subjects diagnosed with MDD showed a significant increase in MOR coupling efficiency in the anterior insular cortex ( $F(1,94) = 4.97$ ,  $p = 0.028$ ; controlling for the significant effect of pH, see main text). Values are mean ± sem, \* $p < 0.05$ .

We next wondered whether MOR coupling was modulated by genetic variation, as per our SNP data. We used dominant negative models and linear regression to investigate relationships between MOR E<sub>max</sub> and observed allelic frequencies. Results showed a nominally significant correlation for rs1074287, a SNP located upstream the canonical MOR transcription start site and within the first intron of rare transcript variants (see Fig. S4A). Accordingly, a higher potency of the DAMGO agonist was observed in carriers of the minor G allele ( $p = 0.017$ ; Fig. S4B), suggesting higher MOR protein abundance in these subjects. This effect, however, did not stand multiple testing correction ( $n = 16$  SNPs; Bonferroni-corrected  $p$  value = 0.27), and will require replication in a larger cohort. In addition, no significant effect of the minor G allele was observed on MOR mRNA expression ( $p = 0.88$ ; Fig. S4C). For other SNPs investigated, no significant correlation was observed ( $p > 0.05$ ; data not shown), including for the A118G SNP (Fig. 2C). Regarding the later, to our knowledge, two previous studies similarly compared MOR coupling across A and G carriers [25, 39], both using brain tissue from the secondary somatosensory cortex (SII) and ventral posterior part of the lateral thalamus. The first study [39] (total sample size  $N = 33$ ) found lower DAMGO efficacy in G carriers in

the SII, but not in the thalamus, while the second study [25] (total sample size  $N = 60$ ), conducted in another cohort, found lower efficacy in G carriers in both brain regions. Combined with our findings, this suggests that the A118G might exert a brain-region specific effect on MOR coupling. This hypothesis is consistent with results from an animal study conducted in genetically modified mice that model the human polymorphism, which reported similar brain-region specificity (with no significant effect of genotype in the AI, see [40]).

Finally, we investigated MOR coupling in relation to psychopathology. Similar to aforementioned analysis of MOR mRNA expression, we first wondered whether the [<sup>35</sup>S]-GTPγS assay was impacted by ante-mortem consumption of drugs of abuse, or SUD diagnosis. Results showed that none of these factors had any effect on MOR E<sub>max</sub> (SUD:  $p = 0.55$ , Fig. S3C; acute opiate:  $p = 0.74$ , Fig. S3D). We then built a GLM model to search for an association with MDD, taking into account the effect of pH as covariate (see above and Fig. S3B). Results indicated that depressed subjects showed significantly higher E<sub>max</sub> ( $F(1,94) = 4.97$ ,  $p = 0.028$ ; Fig. 2D, an effect that remained significant even when correcting for the effect of rs1074287;  $p = 0.0447$ ). This was not accompanied by any change in logEC<sub>50</sub> as a function of MDD ( $p = 0.58$ ), indicating that



**Fig. 3 Relative expression of major inhibitory alpha G-protein subunits in the anterior insula of healthy controls and subjects with major depressive disorder (MDD).** NanoString was used to quantify the mRNA relative expression of 4 inhibitory alpha G-protein subunits. No difference was found between controls and depressed individuals for 3 subunits: GNAI1, GNAO1 and GNAZ. In contrast, subjects with MDD showed a significant increase in the expression of the GNAI2 subunit, compared to controls ( $F(1,89) = 7.42, p = 0.0077$ ). Mean  $\pm$  sem, as well as individual expression values, are shown,  $*p < 0.05$ .

MOR protein signaling, but not affinity, is affected by MDD status. Overall, these results show that coupling of the MOR to inhibitory G-proteins is potentiated in individuals diagnosed with MDD, independently of comorbid SUD.

**Expression of inhibitory alpha G-protein subunits**  
MOR belongs to the family of inhibitory G protein-coupled receptors that activate Gi alpha-subunits, thereby inhibiting adenylate cyclase activity and lowering cAMP levels. To determine whether changes in expression of these proteins might contribute to altered MOR coupling in MDD subjects, we used Nanostring to quantify the expression of the four major corresponding genes: GNAI1, GNAI2, GNAO1, and GNAZ. We first looked at potential effects of clinical and technical covariates, and found none for sex or SUD diagnosis ( $p > 0.05$ ), while RIN affected GNAI1 only ( $p = 0.029$ ). Of note, aging associated with a significant decrease in GNAZ expression ( $p = 0.016$ ; Fig. S5), an effect that may contribute to the decrease in basal binding (i.e., in the absence of MOR agonist) previously observed in [ $^{35}$ S]-GTP $\gamma$ S experiments (see above, and Fig. S4A). More importantly, we investigated potential association with MDD, and observed a selective increase in the expression of GNAI2 in depressed individuals (GLM,  $F(1,89) = 7.42, p = 0.0077$ ; Fig. 3D), which remained significant even when controlling for the number of genes investigated by Nanostring (Bonferroni-corrected  $p$ -value=0.031). Therefore, our results suggest that increased expression of the GNAI2 alpha subunit is a molecular adaptation that may contribute to enhanced MOR coupling in subjects with MDD.

**DISCUSSION**

Inspired by recent imaging studies reporting convergent evidence for MOR dysfunction in the AI of individuals with MDD, the present post-mortem study investigated its molecular expression and function at this brain site.

We first quantified MOR expression using qPCR, and found no significant change as a function of MDD. To our knowledge, only two previous studies quantified MOR mRNA in depressed individuals compared to healthy controls, and reported increased levels in the prefrontal cortex (BA9, see [41]), or no change in the

mediodorsal thalamus or anterior cingulate cortex (BA24/32, [20]), altogether suggesting that MDD-related changes in MOR expression may be subtle, and specific to particular brain regions.

We then used [ $^{35}$ S]-GTP $\gamma$ S to assess MOR signaling. In the past, four human studies have investigated MOR density using the radiolabelled agonist [ $^3$ H]-DAMGO [10, 12–14], while only one publication reported agonist-stimulated [ $^{35}$ S]-GTP $\gamma$ S measures of receptor affinity and density [22]. The two oldest studies, although conducted in small cohorts ([12],  $N = 2-7$  subjects/group; [13],  $N = 15$ /group), provided preliminary evidence for increased MOR abundance in the frontal and temporal cortices, and caudate nucleus. However, the three subsequent studies, performed on frontal cortex tissue (BA9 and BA24) in larger cohorts ([10],  $N = 9-10$ /group; [22],  $N = 28$ /group; [14],  $N = 20-51$ /group), failed to report consistent findings, as none of them found differences between subjects who died by suicide (mostly diagnosed with mood disorders) and controls. In the present study, we investigated a larger cohort ( $N = 40-55$ /group) and focused instead on the AI, previously not investigated using such assays. Our findings indicate that MDD was associated in this brain region with a significant increase in MOR activity, an effect that occurred in the absence of any change in receptor affinity, and that was not accounted for by other confounders such as opiate consumption, SUD, or clinical covariates. This suggests, therefore, that the AI is a prominent brain site where MDD-related changes affecting MOR may contribute to, or alternatively reflect, disease pathophysiology.

To understand molecular adaptations potentially underlying enhanced MOR coupling, and because MOR is an inhibitory G protein-coupled receptor, we next used Nanostring to quantify expression of the 4 major inhibitory alpha G-protein subunits. Results showed a specific increase in the expression of GNAI2 in depressed individuals. Because [ $^{35}$ S]-GTP $\gamma$ S  $E_{max}$  measures at saturating agonist concentration reflect maximal signaling when all receptors are recruited, this increased GNAI2 expression likely contributes to the increase in  $E_{max}$  detected in the GTP $\gamma$ S assay.

Importantly, these findings appear consistent with imaging studies published over the last 15 years [42], mostly conducted using positron emission tomography (PET) and the MOR selective radiotracer [ $^{11}$ C]-Carfentanil. These studies first indicated that

emotionally salient acute stimuli (whether the recall of sad autobiographical events [4], exposure to social rejection [7], or viewing pictures of appetizing food [43]) led in healthy volunteers and depressed individuals [8] to rapid changes in MOR availability, notably at the level of the AI, reflecting the well-characterized notion that MOR activity modulates hedonic tone. Among other findings, MOR availability typically increased in healthy subjects during self-imposed sadness, but decreased during external negative cues, such as social rejection or pain (with variable effects among distinct brain regions; see [44] for review). Beyond these fluctuations, prolonged low mood in MDD also seems to affect MOR function. Accordingly, while an initial PET study reported lower MOR availability in depressed women, which reached significance in the posterior thalamus only (possibly due to modest statistical power,  $N = 14/\text{group}$ ) [5], a more recent study found convergent results by investigating a large cohort characterized for subclinical depressive symptoms ( $N = 135$ , with 113 males; see [2]). In the later work, widespread reductions in MOR availability were observed as a function of symptom severity across numerous cortical regions, with the insula, precuneus, and temporoparietal regions most strongly affected. Importantly, such reduced availability can be interpreted in PET experiments as reflecting either increased occupancy of the MOR by endogenous opioid peptides or, alternatively, lower receptor density or affinity. Results from the present post-mortem study argue in favor of the former possibility. Because increased MOR signaling was observed in our study, in the absence of any change in receptor expression at mRNA level, we hypothesize that depression associates in the AI with higher MOR coupling efficiency ( $E_{\text{max}}$ ), which at least partly stems from higher GNAI2 expression. In addition, PET studies suggest enhanced endogenous release of opioid peptides (lower MOR availability) in MDD patients. Therefore, 2 distinct pre-opioid release and post-synaptic (receptor coupling) mechanisms might converge to potentiate the insular opioidergic tone in depressed individuals. Importantly, considering the wealth of animal and human data supporting the notion that acute MOR signaling has positive effects on mood, and buffers negative emotions associated with social rejection or physical pain, in particular in the AI [1, 42–45], we favor the interpretation that these adaptations reflect a compensatory mechanism that counteracts low affect. Alternatively, because MOR-dependent regulation of mood fluctuations remain incompletely resolved (with e.g., potentially opposite effects across distinct cortical MOR pools [46], or following acute or chronic receptor activation [1]), it is nevertheless possible that increased MOR signaling in the AI may represent a pathological mechanism contributing to depressed mood. Future studies will be necessary to differentiate the 2 hypotheses.

While MOR-mediated regulation of reward and mood has been primarily investigated in animal models in structures from the mesolimbic dopaminergic system, such as the ventral tegmental area and basal ganglia [1, 47], it has been left relatively unexplored in cortical structures. Studies in human, however, clearly indicate that MOR signaling in the AI contributes to important physiological function, in particular the processing of emotional aspects of pain that may arise from somatosensory stimuli (physical pain), or from situations that threatens the organism's survival, internal homeostasis, or social position (i.e., psychological or social pain) [44, 45]. Our findings are consistent with this framework, and suggest that opiate medications that recruit the MOR and are currently under intense examination for the management of depression and suicidal ideas (e.g., buprenorphine, frequently combined with samidorphan) [48, 49], may act at least in part by potentiating in the AI the beneficial impact of MOR on social and emotional distress.

In conclusion, the present work brings new knowledge that, in combination with imaging studies, indicates that the AI is a critical brain site where increased MOR signaling occurs in the context of

MDD and suicide. Future studies will be necessary to determine whether similar adaptations may affect other brain regions where MOR contributes to reward and affective regulation.

## FUNDING AND DISCLOSURES

The authors reported no biomedical financial interests or potential conflicts of interest. PEL was supported by fellowships from the Fondation Fyssen, Fondation Bettencourt-Schueller, Canadian Institutes of Health Research (201311MFE-320636-218885), American Foundation for Suicide Prevention (PDF-0-081-13), and Fondation pour la Recherche Médicale (ARF20160936006). This work was also supported by the Agence Nationale de la Recherche (ANR grant JCJC SignOp, PEL).

## AUTHOR CONTRIBUTIONS

PEL, FJ, BLK and GT conceived the study. PEL performed RNA extractions, qPCR, and Nanostring experiments. DA conducted SNP genotyping. DF, PEL and BLK conceived and performed [ $^{35}\text{S}$ ]-GTP $\gamma$ S experiments. PEL prepared the manuscript, and all authors approved its final version.

## ADDITIONAL INFORMATION

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41386-021-00974-y>.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## REFERENCES

1. Lutz PE, Kieffer BL. Opioid receptors: distinct roles in mood disorders. *Trends Neurosci.* 2013;36:195–206.
2. Nummenmaa L, Karjalainen T, Isojärvi J, Kantonen T, Tuisku J, Kaasinen V, et al. Lowered endogenous mu-opioid receptor availability in subclinical depression and anxiety. *Neuropsychopharmacology.* 2020. <https://doi.org/10.1038/s41386-020-0725-9>.
3. Zubieta JK, Smith YR, Bueller JA, Xu Y, Kilbourn MR, Jewett DM, et al. Regional mu opioid receptor regulation of sensory and affective dimensions of pain. *Science.* 2001;293:311–5.
4. Zubieta JK, Ketter TA, Bueller JA, Xu Y, Kilbourn MR, Young EA, et al. Regulation of human affective responses by anterior cingulate and limbic mu-opioid neurotransmission. *Arch Gen Psychiatry.* 2003;60:1145–53.
5. Kennedy SE, Koeppe RA, Young EA, Zubieta JK. Dysregulation of endogenous opioid emotion regulation circuitry in major depression in women. *Arch Gen Psychiatry.* 2006;63:1199–208.
6. Namkung H, Kim S-H, Sawa A. The insula: an underestimated brain area in clinical neuroscience, psychiatry, and neurology. *Trends Neurosci.* 2017;40:200–7.
7. Hsu DT, Sanford BJ, Meyers KK, Love TM, Hazlett KE, Wang H, et al. Response of the mu-opioid system to social rejection and acceptance. *Mol Psychiatry.* 2013;18:1211–7.
8. Hsu DT, Sanford BJ, Meyers KK, Love TM, Hazlett KE, Walker SJ, et al. It still hurts: altered endogenous opioid activity in the brain during social rejection and acceptance in major depressive disorder. *Mol Psychiatry.* 2015;20:193–200.
9. Spagnolo PA, Kimes A, Schwandt ML, Shokri-Kojori E, Thada S, Phillips KA, et al. Striatal dopamine release in response to morphine: a [ $^{11}\text{C}$ ]Raclopride positron emission tomography study in healthy men. *Biol Psychiatry.* 2019;86:356–64.
10. Zalsman G, Molcho A, Huang Y, Dwork A, Li S, Mann JJ. Postmortem mu-opioid receptor binding in suicide victims and controls. *J Neural Transm.* 2005;112:949–54.
11. Meana JJ, Gonzalez-Maeso J, Garcia-Sevilla JA, Guimon J. mu-opioid receptor and alpha2-adrenoceptor agonist stimulation of [ $^{35}\text{S}$ ]GTP $\gamma$  binding to G-proteins in postmortem brains of opioid addicts. *Mol Psychiatry.* 2000;5:308–15.
12. Gross-Isseroff R, Dillon KA, Israeli M, Biegon A. Regionally selective increases in mu opioid receptor density in the brains of suicide victims. *Brain Res.* 1990;530:312–6.
13. Gabilondo AM, Meana JJ, Garcia-Sevilla JA. Increased density of mu-opioid receptors in the postmortem brain of suicide victims. *Brain Res.* 1995;682:245–50.
14. Scarr E, Money TT, Pavey G, Neo J, Dean B. Mu opioid receptor availability in people with psychiatric disorders who died by suicide: a case control study. *BMC Psychiatry.* 2012;12:126.

15. Dumais A, Lesage AD, Alda M, Rouleau G, Dumont M, Chawky N, et al. Risk factors for suicide completion in major depression: a case-control study of impulsive and aggressive behaviors in men. *Am J Psychiatry*. 2005;162:2116–24.
16. Brent DA, Perper JA, Moritz G, Allman CJ, Roth C, Schweers J, et al. The validity of diagnoses obtained through the psychological autopsy procedure in adolescent suicide victims: use of family history. *Acta Psychiatr Scand*. 1993;87:118–22.
17. Conner KR, Conwell Y, Duberstein PR. The validity of proxy-based data in suicide research: a study of patients 50 years of age and older who attempted suicide. II. Life events, social support and suicidal behavior. *Acta Psychiatr Scand*. 2001; 104:452–7.
18. Kelly TM, Mann JJ. Validity of DSM-III-R diagnosis by psychological autopsy: a comparison with clinician ante-mortem diagnosis. *Acta Psychiatr Scand*. 1996; 94:337–43.
19. Mai JK, Paxinos G, Voss T. Atlas of the human brain. Academic Press Inc, 3rd ed. Elsevier Science. 2007.
20. Lutz PE, Gross JA, Dhir SK, Maussion G, Yang J, Bramouille A, et al. Epigenetic regulation of the kappa opioid receptor by child abuse. *Biol Psychiatry*. 2018; 84:751–61.
21. Weckx S, Del-Favero J, Rademakers R, Claes L, Cruts M, De Jonghe P, et al. novoSNP, a novel computational tool for sequence variation discovery. *Genome Res*. 2005;15:436–42.
22. Gonzalez-Maeso J, Rodriguez-Puertas R, Meana JJ, Garcia-Sevilla JA, Guimon J. Neurotransmitter receptor-mediated activation of G-proteins in brains of suicide victims with mood disorders: selective supersensitivity of alpha(2A)-adrenoceptors. *Mol Psychiatry*. 2002;7:755–67.
23. Lutz PE, Tanti A, Gasecka A, Barnett-Burns S, Kim JJ, Zhou Y, et al. Association of a history of child abuse with impaired myelination in the anterior cingulate cortex: convergent epigenetic, transcriptional, and morphological evidence. *Am J Psychiatry*. 2017;174:1185–94.
24. Geiss GK, Bumgarner RE, Birditt B, Dahl T, Dowidar N, Dunaway DL, et al. Direct multiplexed measurement of gene expression with color-coded probe pairs. *Nat Biotechnol*. 2008;26:317–25.
25. Oertel BG, Doehring A, Roskam B, Kettner M, Hackmann N, Ferreiros N, et al. Genetic-epigenetic interaction modulates mu-opioid receptor regulation. *Hum Mol Genet*. 2012;21:4751–60.
26. Lutz PE, Kieffer BL. The multiple facets of opioid receptor function: implications for addiction. *Curr Opin Neurobiol*. 2013;23:473–9.
27. Contet C, Kieffer BL, Befort K. Mu opioid receptor: a gateway to drug addiction. *Curr Opin Neurobiol*. 2004;14:370–8.
28. Xu J, Faskowitz AJ, Rossi GC, Xu M, Lu Z, Pan YX, et al. Stabilization of morphine tolerance with long-term dosing: association with selective upregulation of mu-opioid receptor splice variant mRNAs. *Proc Natl Acad Sci USA*. 2015;112:279–84.
29. Verzillo V, Madaia PA, Liu N-J, Chakrabarti S, Gintzler AR. Mu-opioid receptor splice variants: sex-dependent regulation by chronic morphine. *J Neurochem*. 2014;130: 790–6.
30. Castelli MP, Melis M, Mameli M, Fadda P, Diaz G, Gessa GL. Chronic morphine and naltrexone fail to modify mu-opioid receptor mRNA levels in the rat brain. *Brain Res Mol Brain Res*. 1997;45:149–53.
31. Bilbao A, Robinson JE, Heilig M, Malanga CJ, Spanagel R, Sommer WH, et al. A pharmacogenetic determinant of mu-opioid receptor antagonist effects on alcohol reward and consumption: evidence from humanized mice. *Biol Psychiatry*. 2015; 77:850–8.
32. Belzeaux R, Gorgievski V, Fiori LM, Lopez JP, Grenier J, Lin R, et al. GPR56/ADGRG1 is associated with response to antidepressant treatment. *Nat Commun*. 2020;11:1635.
33. Mague SD, Blendy JA. OPRM1 SNP (A118G): involvement in disease development, treatment response, and animal models. *Drug Alcohol Depend*. 2010;108:172–82.
34. Cimino S, Carola V, Cerniglia L, Bussone S, Bevilacqua A, Tambelli R. The mu-opioid receptor gene A118G polymorphism is associated with insecure attachment in children with disruptive mood regulation disorder and their mothers. *Brain Behav*. 2020;10:e01659.
35. Troisi A, Frazzetto G, Carola V, Di Lorenzo G, Coviello M, Siracusanò A, et al. Variation in the mu-opioid receptor gene (OPRM1) moderates the influence of early maternal care on fearful attachment. *Soc Cogn Affect Neurosci*. 2012;7:542–7.
36. Hancock DB, Levy JL, Gaddis NC, Glasheen C, Saccone NL, Page GP, et al. Cis-expression quantitative trait loci mapping reveals replicable associations with heroin addiction in OPRM1. *Biol Psychiatry*. 2015. <https://doi.org/10.1016/j.biopsych.2015.01.003>.
37. Mansour A, Fox CA, Akil H, Watson SJ. Opioid-receptor mRNA expression in the rat CNS: anatomical and functional implications. *Trends Neurosci*. 1995;18:22–29.
38. Gonzalez-Maeso J, Torre I, Rodriguez-Puertas R, Garcia-Sevilla JA, Guimon J, Meana JJ. Effects of age, postmortem delay and storage time on receptor-mediated activation of G-proteins in human brain. *Neuropsychopharmacology*. 2002;26:468–78.
39. Oertel BG, Kettner M, Scholich K, Renne C, Roskam B, Geisslinger G, et al. A common human micro-opioid receptor genetic variant diminishes the receptor signaling efficacy in brain regions processing the sensory information of pain. *J Biol Chem*. 2009;284:6530–5.
40. Wang YJ, Huang P, Blendy JA, Liu-Chen LY. Brain region- and sex-specific alterations in DAMGO-stimulated [(35) S]GTPgammaS binding in mice with Oprm1 A112G. *Addict Biol*. 2014;19:354–61.
41. Escriba PV, Ozaita A, Garcia-Sevilla JA. Increased mRNA expression of alpha2A-adrenoceptors, serotonin receptors and mu-opioid receptors in the brains of suicide victims. *Neuropsychopharmacology*. 2004;29:1512–21.
42. Lutz PE, Courtet P, Calati R. The opioid system and the social brain: implications for depression and suicide. *J Neurosci Res*. 2018. <https://doi.org/10.1002/jnr.24269>.
43. Nummenmaa L, Saanijoki T, Tuominen L, Hirvonen J, Tuulari JJ, Nuutila P, et al. mu-opioid receptor system mediates reward processing in humans. *Nat Commun*. 2018;9:1500.
44. Pecina M, Karp JF, Mathew S, Todtenkopf MS, Ehrlich EW, Zubieta JK. Endogenous opioid system dysregulation in depression: implications for new therapeutic approaches. *Mol Psychiatry*. 2018. <https://doi.org/10.1038/s41380-018-0117-2>.
45. Eisenberger NI. Social ties and health: a social neuroscience perspective. *Curr Opin Neurobiol*. 2013. <https://doi.org/10.1016/j.conb.2013.01.006>.
46. Castro DC, Berridge KC. Opioid and orexin hedonic hotspots in rat orbitofrontal cortex and insula. *Proc Natl Acad Sci USA*. 2017;114:E9125–E9134.
47. Russo SJ, Nestler EJ. The brain reward circuitry in mood disorders. *Nat Rev Neurosci*. 2013;14:609–25.
48. Fava M, Thase ME, Trivedi MH, Ehrlich E, Martin WF, Memisoglu A, et al. Opioid system modulation with buprenorphine/samidorphan combination for major depressive disorder: two randomized controlled studies. *Mol Psychiatry*. 2020;25:1580–91.
49. Yovell Y, Bar G, Mashiah M, Baruch Y, Briskman I, Asherov J, et al. Ultra-low-dose buprenorphine as a time-limited treatment for severe suicidal ideation: a randomized controlled trial. *Am J Psychiatry*. 2016;173:491–8.