CORRECTION

Open Access

Correction: Impact of energy turnover on the regulation of glucose homeostasis in healthy subjects

Franziska Büsing¹, Franziska Anna Hägele¹, Alessa Nas², Mario Hasler³, Manfred James Müller¹ and Anja Bosy-Westphal¹

Correction to: Nutrition & Diabetes

<https://doi.org/10.1038/s41387-019-0089-6>
published online 08 August 2019

This has been corrected in both the PDF and HTML versions of the Article.

Published online: 07 October 2019

Since publication of this article the authors noted that the legend for Table 1 was incomplete, as the subtitle was missing. The complete table should appear as given below:

Correspondence: Anja Bosy-Westphal (abosyw@nutrition.uni-kiel.de)

¹Institute of Human Nutrition and Food Science, Christian-Albrechts University of Kiel, Kiel, Germany

²Institute of Nutritional Medicine, University of Hohenheim, Stuttgart, Germany

³Applied Statistics, Faculty of Agricultural and Nutritional Sciences, Christian-Albrechts University of Kiel, Kiel, Germany

© The Author(s) 2019

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Table 1 Physical activity and daily energy-, macronutrient- and glycemic load intake during caloric restriction, zero energy balance and overfeeding with low, medium and high energy turnover (ET)

	Caloric restriction			Energy balance			Overfeeding		
	Low ET	Med ET	High ET	Low ET	Med ET	High ET	Low ET	Med ET	High ET
Physical activity									
PAL	1.26 ± 0.06	1.52 ± 0.05	1.74 ± 0.07	1.30 ± 0.03	1.57 ± 0.04	1.75 ± 0.05	1.34 ± 0.05	1.55 ± 0.04	1.71 ± 0.05
Steps [counts/d]	545 ± 902	17 618 ± 557	34 527 ± 1847	464 ± 912	17 710 ± 219	34 414 ± 1539	436 ± 847	17 790 ± 382	34 821 ± 1568
Diet									
Energy intake [kcal/d]	1 792 ± 144	2 156 ± 78	2 494 ± 90	2 390 ± 97	2 863 ± 72	3 325 ± 177	2 986 ± 99	3 570 ± 183	4 146 ± 320
CHO intake [g/d]	219.4 ± 20.8	255.1 ± 37.3	308.2 ± 12.0	292.7 ± 13.9	350.1 ± 6.0	406.5 ± 20.3	366.2 ± 11.8	437.2 ± 22.1	506.7 ± 37.9
Fat intake [g/d]	70.5 ± 4.9	84.7 ± 2.6	97.7 ± 5.0	93.7 ± 3.3	112.8 ± 3.9	131.0 ± 7.5	117.3 ± 3.9	140.8 ± 7.0	163.8 ± 12.5
Protein intake [g/d]	66.1 ± 4.0	79.4 ± 3.3	92.0 ± 5.6	88.4 ± 0.0	106.1 ± 5.6	120.3 ± 9.3	109.8 ± 3.0	131.0 ± 8.6	153.3 ± 13.0
Glycemic load [g/d]	124.0 ± 8.9	149.2 ± 4.7	173.8 ± 7.0	166.0 ± 6.5	197.8 ± 5.7	228.9 ± 14.1	206.4 ± 7.0	247.4 ± 13.0	285.9 ± 24.0

Values are means ± SDs; $n = 16$; linear mixed model with multiple contrast tests, results of physical activity were compared at the same level of ET between all energy balance conditions (all $p > 0.05$) and within all energy balance conditions at three levels of ET (all significantly different at $p < 0.001$); results of dietary intake were compared at the same level of ET between all energy balance conditions (all significantly different at $p < 0.001$) and within all energy balance conditions at three levels of ET (all significantly different at $p < 0.001$); ET, energy turnover; CHO, carbohydrate