



Correction: Energy-dependent nucleolar localization of p53 in vitro requires two discrete regions within the p53 carboxyl terminus

O Karni-Schmidt¹ · A Friedler^{2,3} · A Zupnick¹ · K McKinney¹ · M Mattia¹ · R Beckerman¹ · P Bouvet⁴ · M Sheetz¹ · A Fersht² · C Prives¹

Published online: 1 August 2018
© Springer Nature Limited 2018

Correction to: Oncogene (2007) 26, 3878–3891; <https://doi.org/10.1038/sj.onc.1210162>; published online: 22 Jan 2007

Following the publication of this article the authors noted that two images were duplicated in Figure 2B. The corrected figure 2B is below. The authors wish to apologize for any inconvenience caused.

✉ C Prives
clp3@columbia.edu

¹ Department of Biological Sciences, Columbia University, New York, USA

² Department of MRC-CPE, Center for Protein Engineering, Medical Research Council, Cambridge, UK

³ Department of Organic Chemistry, The Hebrew University of Jerusalem, Givat Ram, Jerusalem, Israel

⁴ Laboratoire de Biologie Moléculaire de la Cellule/UMR 5161, Ecole Normale Supérieure de Lyon 46, Allée d'Italie, 69364 Lyon Cedex, France




Fig. 2b .