Correction: Signal peptidase complex 18, encoded by SEC11A, contributes to progression via TGF- α secretion in gastric cancer

N Oue¹ · Y Naito¹ · T Hayashi¹ · M Takigahira² · A Kawano-Nagatsuma³ · K Sentani¹ · N Sakamoto¹ · H Z Oo¹ · N Uraoka¹ · K Yanagihara⁴ · A Ochiai³ · H Sasaki⁵ · W Yasui¹

Published online: 7 June 2019

© The Author(s), under exclusive licence to Springer Nature Limited 2019

Correction to: Oncogene

<https://doi.org/10.1038/onc.2013.364>

Published online 2 September 2013

This has been corrected from Zarni Oo H to Oo HZ in both the HTML and PDF versions of the Article.

Following publication of this article the author Htoo Zarni Oo noted that their name had been published incorrectly.

The original article can be found online at <https://doi.org/10.1038/onc.2013.364>.

✉ W Yasui
wyasui@hiroshima-u.ac.jp

- ¹ Department of Molecular Pathology, Hiroshima University Institute of Biomedical and Health Sciences, Hiroshima, Japan
- ² Investigative Treatment Division, Research Center for Innovative Oncology, National Cancer Center Hospital East, Kashiwa, Japan
- ³ Pathology Division, and Research Center for Innovative Oncology, National Cancer Center Hospital East, Kashiwa, Japan
- ⁴ Division of Translational Research, Research Center for Innovative Oncology, National Cancer Center Hospital East, Kashiwa, Japan
- ⁵ Division of Genetics, National Cancer Center Research Institute, Tokyo, Japan