



Correction: The topoisomerase I- and p53-binding protein topors is differentially expressed in normal and malignant human tissues and may function as a tumor suppressor

Ahamed Saleem¹ · Jayeeta Dutta¹ · Diptee Malegaonkar¹ · Farheena Rasheed¹ · Zeshaan Rasheed¹ · Rajeev Rajendra¹ · Henderson Marshall¹ · Minjie Luo² · Honghua Li² · Eric H Rubin^{1,3}

Published online: 11 July 2019
© Springer Nature Limited 2019

Correction to: *Oncogene*

<https://doi.org/10.1038/sj.onc.1207700>

Following publication of this article the author Minjie Luo noted that their name had been published incorrectly as Minje Luo.

The original article can be found online at <https://doi.org/10.1038/sj.onc.1207700>.

✉ Eric H Rubin
ehrubin@umdnj.edu

¹ Department of Pharmacology, the Cancer Institute of New Jersey, Robert Wood Johnson Medical School, University of Medicine and Dentistry of New Jersey, New Brunswick, NJ, USA

² Department of Molecular Genetics and Microbiology, the Cancer Institute of New Jersey, Robert Wood Johnson Medical School, University of Medicine and Dentistry of New Jersey, New Brunswick, NJ, USA

³ Department of Medicine, the Cancer Institute of New Jersey, Robert Wood Johnson Medical School, University of Medicine and Dentistry of New Jersey, New Brunswick, NJ, USA