CORRECTION

Correction: Effect on splanchnic oxygenation of breast milk, fortified breast milk, and formula milk in preterm infants

Carlo Dani, Caterina Coviello, Simona Montano, Giulia Remaschi, Chiara Petrolini, Maria Chiara Strozzi, Elena Maggiora, Miriam Sabatini and Diego Gazzolo

Correction to: *Pediatric Research* https://doi.org/10.1038/s41390-020-0935-1, published online 7 May 2020

In the original version of this article, Table 2 reported data as mean values and standard variation, while the *P*-values were calculated using median values and interquartile ranges (IQR). Table 2 has now been corrected to display data as median values and IQR. The original, incorrect version of Table 2 is displayed below for reference.

Additionally, in the section "Statistical analysis" it was incorrectly stated that serial measurements of rSO_2S and FOES, rather than median values of rSO_2S and FOES, were compared by Kruskal-Wallis one-way ANOVA and Dunn's method. This has been corrected in both the PDF and HTML versions of the article.

Table 2. Changes of rSO2S and FOES before (T0), during (T1) and after feeding (T2) in MOM-, FBM-, and PTF-fed infants. Mean ± (SD).

	T_0	T_1	T_2
MOM group ($n = 18$)			
rSO_2S (%)	57.4 ± 7.9	60.4 ± 8.5	61.9 ± 3.7
Difference vs. baseline		3.4 ± 11.3[§]	4.5 ± 11.8[§§]
FOES	0.35 ± 0.13	0.32 ± 0.17	0.27 ± 0.05
Difference vs. baseline		-0.03 ± 0.10	-0.08 ± 0.11[§]
FBM group ($n = 18$)			
rSO_2S (%)	57.6 ± 10.9	55.1 ± 11.4*	58.3 ± 13.3**
Difference vs. baseline		-2.4 ± 6.3	0.7 ± 11.7
FOES	0.43 ± 0.15	0.41 ± 0.13*	0.38 ± 0.14**
Difference vs. baseline		-0.01 ± 0.13	-0.04 ± 0.12
PTF ($n = 18$)			
rSO_2S (%)	57.9 ± 15.1	51.1 ± 13.8	46.9 ± 17.8*
Difference vs. baseline		-6.8 ± 7.4	-11.0 ± 10.6
FOES	0.44 ± 0.21	0.47 ± 0.13*	0.50 ± 0.18*[#]
Difference vs. baseline		0.02 ± 0.10	0.06 ± 0.13

*$P < 0.001$ vs. T_0; **$P < 0.001$ vs. T_2; [#]$P < 0.05$ vs. T_1.
[§]$P < 0.005$ vs. PTF; [§§]$P < 0.00_1$ vs. PTF.