

COMMENT



How hypoxia slows fetal growth: insights from high altitude

Lorna G. Moore¹✉

© The Author(s), under exclusive licence to the International Pediatric Research Foundation, Inc 2021

A continuous oxygen supply is vital for fueling fetal growth, yet we know surprisingly little as to the precise mechanisms by which hypoxia reduces fetal growth. Animal models, while important, are limited by the enormous variability in mammalian physiologic responses to pregnancy. One of the earliest models for understanding hypoxia-induced reductions in fetal growth came from studies at high altitude that showed high altitude to have one of the strongest depressant effects on fetal growth. But all populations are not equally affected. Dolma et al. have added valuable information now showing that that Ladakhis, like Tibetans and Andeans, have a lower-than-expected frequency of small-for-gestational age (SGA) infants. Consistent with previous reports, these authors also found that uterine artery diameters at mid pregnancy (weeks 26–28) were larger in women giving birth to appropriate-for-gestational age (AGA) than SGA infants at high but, interestingly, not at low altitude. Much remains to be learned about the physiologic pathways by which hypoxia impairs fetal growth. The variability among high-altitude populations and increasing sophistication of tools for investigating causal mechanisms have the potential to expand our presently limited means for identifying new treatments for hypoxia-related complications of pregnancy and fetal life.

Pediatric Research (2022) 91:17–18; <https://doi.org/10.1038/s41390-021-01784-0>

Together with their Indian colleagues, this University College London team headed by Hugh Montgomery and Sara Hillman have added valuable information for persons interested in mechanisms underlying hypoxia-associated fetal growth restriction. Consistent with previous studies in multigenerational Himalayan residents, their results show that residents of high altitude (conventionally defined as >2500 m) in Leh, Ladakh (3524 m, $n = 316$) give birth to heavier than expected babies (3.15 kg), especially when of Tibetan ancestry (3.62 kg), and correspondingly have fewer small-for-gestational age (SGA) infants than residents of Delhi (216 m, $n = 101$) (14 vs. 19%, respectively). Maternal risk factors (e.g., age, body mass index (BMI), socioeconomic indicators) were similar in the two groups as were gestational age at delivery and baby sex, thus implicating protection from fetal growth restriction in the Ladakhi residents as the cause of the birth weight differences. Associated with such protection were greater, but still within the normal range, maternal age and BMI and greater uterine artery diameter at weeks 26–28. Interestingly, while similar relationships with birth weight were seen for maternal age and BMI at low altitude, there were no differences in uterine artery diameter between the mothers of SGA vs. appropriate-for-gestational age (AGA) infants at low altitude.

Birth weights have long been observed to fall at high altitude, decreasing on average 100 g per 1000 m altitude gain.¹ The effect of high altitude is independent of other known risk factors (e.g., maternal age, weight gain, access to prenatal care, smoking) and has the greatest effect on birth weight next to gestational age. While the consensus is that reduced oxygen availability (hypoxia) rather than some other attribute of the high-altitude environment is responsible, the precise mechanisms by which hypoxia slows fetal growth remain unknown. Nor are all women or populations

equally affected. As noted in the present report, multigenerational high-altitude residents are protected relative to women of low-altitude, European, Han, or as now shown here, Indo-Aryan ancestry. This protective effect is seen even when the persons of low-altitude ancestry were themselves born and raised at high altitude, thereby implicating genetic rather than developmental factors.² Ladakhis, like Tibetans, have descended from Mongolian populations but also have ancestral roots among the Dards, a group living in the Hindu Kush mountains of northern Pakistan. The women in the present report had >5 generations of high-altitude residence but perhaps less than the >15,000-to-25,000-year duration estimated for Andeans or Tibetans, respectively. If so, the comparatively short time frame (in evolutionary terms) would be consistent with the likelihood that uterine vascular responses to pregnancy were subject to strong selective pressure and/or involved relatively few genes.

These authors' observation of lower uterine artery diameters at weeks 26–28 in high-altitude residents giving birth to SGA vs. AGA babies is consistent with reports of less pregnancy-associated increase in uterine artery diameter or blood flow at weeks 20–36 in 3100 vs. 1600 m Colorado residents of European ancestry and greater uterine artery diameters and blood flow in multigenerational Andean or Tibetan residents of ~3600 m than in women of European or Han ancestry, respectively, living at the same altitude.^{3–5} Dolma et al.'s observation that such differences in uterine artery diameter at weeks 26–28 is important because this is before growth begins to measurably slow at high altitude⁶ and, in turn, suggests that greater blood flow enables the maintenance of normal fetal growth under conditions of hypoxia. But exactly how hypoxia acts to slow fetal growth remains unclear. Experimental animal and isolated human myometrial artery

¹Department of Obstetrics and Gynecology, University of Colorado Denver School of Medicine, MS 8613, 12700 E 19th Avenue, Room 3004, Aurora, CO 80045, USA.
✉email: Lorna.Moore@cuanschutz.edu

vasoreactivity studies indicate that hypoxia, whether due to altitude or the pregnancy complications of fetal growth restriction or preeclampsia,⁷ impairs normal maternal uterine vasodilator and/or growth responses to pregnancy. Pregnancy profoundly vasodilates the entire maternal systemic (and pulmonary) circulation but has even greater vasodilator effects in the utero-placental vasculature, which thereby directs the majority of the ~30% rise in cardiac output to that vascular bed by term. Uterine vasodilation is due not only to trophoblast remodeling of the end (spiral) arteries but also to upstream effects as demonstrated by the enhanced vasodilator response and/or marked growth seen during pregnancy in uterine, arcuate, basal, and myometrial arteries.^{8,9}

As remarked by the authors, the availability of multiple populations with and without genetic adaptations to high altitude makes high altitude a unique natural laboratory for investigating hypoxia-associated fetal as well as maternal complications of pregnancy. There are presently no treatments for such disorders, apart from early delivery that is costly and can itself have unfavorable effects. As noted, promising results have been reported by our group using adenosine monophosphate kinase (AMPK) or peroxisome proliferator-activated receptor gamma activators in vitro in resistance-sized human uterine vessels and in vivo in experimental animals. Noteworthy was that the enhanced vasodilation in response to AMPK activation was selective for the maternal uteroplacental circulation, unlike the body-wide vasodilation seen using sildenafil whose clinical trial as a preeclampsia therapy was stopped due to adverse fetal effects.^{10,11}

Additional studies are needed for identifying the ways in which uterine blood flow can be raised selectively as well as for defining the mechanisms by which hypoxia influences fetal growth. Future studies at high altitudes can be valuable not only because the increased frequency of fetal growth restriction (and preeclampsia, another pregnancy disorder associated with uteroplacental ischemia) reduces the number needed to treat¹² but also because most high-altitude residents live in low- to middle-income countries where 98–99% of maternal and perinatal deaths occur.¹³

REFERENCES

- Jensen, G. M. & Moore, L. G. The effect of high altitude and other risk factors on birthweight: independent or interactive effects? *Am. J. Public Health* **87**, 1003–1007 (1997).
- Julian, C. G., Hageman, J. L., Wilson, M. J., Vargas, E. & Moore, L. G. Lowland origin women raised at high altitude are not protected against lower uteroplacental O₂ delivery during pregnancy or reduced birth weight. *Am. J. Hum. Biol.* **23**, 509–516 (2011).
- Julian, C. G. et al. Lower uterine artery blood flow and higher endothelin relative to nitric oxide metabolite levels are associated with reductions in birth weight at high altitude. *Am. J. Physiol.* **295**, R906–R915 (2008).
- Julian, C. G. et al. Augmented uterine artery blood flow and oxygen delivery protect Andeans from altitude-associated reductions in fetal growth. *Am. J. Physiol.* **296**, R1564–R1575 (2009).
- Zamudio, S. et al. Maternal oxygen delivery is not related to altitude- and ancestry-associated differences in human fetal growth. *J. Physiol.* **582**, 883–895 (2007).
- Unger, C., Weiser, J. K., McCullough, R. E., Keefer, S. & Moore, L. G. Altitude, low birth weight, and infant mortality in Colorado. *JAMA* **259**, 3427–3432 (1988).
- Lorca, R. et al. Chronic hypoxia reduces NO-dependent myometrial artery vasodilator response during pregnancy. *Hypertension* **73**, 1319–1326 (2019).
- Keyes, L. E., Majack, R., Dempsey, E. C. & Moore, L. G. Pregnancy stimulation of DNA synthesis and uterine blood flow in the guinea pig. *Pediatr. Res.* **41**, 708–715 (1997).
- Osol, G. & Moore, L. G. Maternal uterine vascular remodeling during pregnancy. *Microcirculation* **21**, 38–47 (2014).
- Hale, S. A. et al. Sildenafil increases uterine blood flow in nonpregnant nulliparous women. *Reprod. Sci.* **17**, 358–365 (2010).
- Pels, A. et al. Maternal sildenafil vs placebo in pregnant women with severe early-onset fetal growth restriction: a randomized clinical trial. *JAMA Netw. Open* **3**, e205323 (2020).
- Askie, L. M., Duley, L., Henderson-Smart, D. J., Stewart, L. A. & Group, P. C. Anti-platelet agents for prevention of pre-eclampsia: a meta-analysis of individual patient data. *Lancet* **369**, 1791–1798 (2007).
- Duley, L. Maternal mortality associated with hypertensive disorders of pregnancy in Africa, Asia, Latin America and the Caribbean. *Br. J. Obstet. Gynaecol.* **99**, 547–553 (1992).

AUTHOR CONTRIBUTIONS

L.G.M. conceived of, drafted, revised, and approved the commentary submitted.

FUNDING INFORMATION

This work was funded by NIH HD088590 and NIH HL138181.

COMPETING INTERESTS

The author declares no competing interests.

ADDITIONAL INFORMATION

Correspondence and requests for materials should be addressed to Lorna G. Moore.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.