(Check for updates

POPULATION STUDY ARTICLE Prenatal smoking and drinking are associated with altered newborn autonomic functions

Ayesha Sania^{1,2,8 \Bigsty}, Michael M. Myers^{1,2,8}, Nicolò Pini^{1,2}, Maristella Lucchini^{1,2}, J. David Nugent^{1,2}, Lauren C. Shuffrey^{1,2}, Shreya Rao³, Jennifer Barbosa¹, Jyoti Angal^{4,5}, Amy J. Elliott^{4,5}, Hein J. Odendaal⁶, William P. Fifer^{1,2,7} and for the PASS Network

© The Author(s), under exclusive licence to the International Pediatric Research Foundation, Inc 2022

BACKGROUND: Prenatal smoking and drinking are associated with sudden infant death syndrome and neurodevelopmental disorders. Infants with these outcomes also have altered autonomic nervous system (ANS) regulation. We examined the effects of prenatal smoking and drinking on newborn ANS function.

METHODS: Pregnant women were enrolled in Northern Plains, USA (NP) and Cape Town (CT), South Africa. Daily drinking and weekly smoking data were collected prenatally. Physiological measures were obtained during sleep 12–96 h post-delivery. **RESULTS:** In all, 2913 infants from NP and 4072 from CT were included. In active sleep, newborns of mothers who smoked throughout pregnancy, compared to non-smokers, had higher breathing rates (2.2 breaths/min; 95% CI: 0.95, 3.49). Quit-early smoking was associated with reductions in beat-to-beat heart rate variability (HRV) in active (-0.08 s) and quiet sleep (-0.11 s) in CT. In girls, moderate-high continuous smoking was associated with increased systolic (3.0 mmHg, CI: 0.70, 5.24) and diastolic blood pressure (2.9 mmHg, CI: 0.72, 5.02). In quiet sleep, low-continuous drinking was associated with slower heart rate (-4.5 beat/min). In boys, low-continuous drinking was associated with a reduced ratio of low-to-high frequency HRV (-0.11, CI: -0.21, -0.02). **CONCLUSIONS:** These findings highlight potential ANS pathways through which prenatal drinking and smoking may contribute to neurodevelopment outcomes.

Pediatric Research (2023) 93:242-252; https://doi.org/10.1038/s41390-022-02060-5

IMPACT:

- In this prospective cohort study of 6985 mother–infant dyads prenatal drinking and smoking were associated with multiple ANS parameters.
- Smoking was associated with increased neonatal breathing rates among all infants, and heart rate variability (HRV) and blood pressure (BP) among girls.
- Drinking was associated with reductions in HR and BP among all newborns, and reductions in the ratio of low to-high frequency HRV among boys.
- These findings suggest that prenatal smoking and drinking alter newborn ANS which may presage future neurodevelopmental disorders.

INTRODUCTION

Although many women modify their drinking and smoking behavior following pregnancy recognition,^{1,2} heightened risk of adverse outcomes associated with prenatal alcohol (PAE) and tobacco (PTE) exposures remains a public health concern, especially in low-resource settings and in populations with high prevalence of alcohol and tobacco use.^{3,4} There is well documented evidence of role of prenatal drinking and smoking on increased fetal loss, stillbirth, preterm birth, intrauterine growth restriction, sudden infant death syndrome (SIDS), and other neurodevelopmental disorders.^{5–7} Recent studies also linked PAE and PTE to

dysregulation of the fetal and infant autonomic nervous system (ANS)^{8,9} and the newborn central nervous system.¹⁰

Alcohol and nicotine can directly affect the developing fetal brain leading to autonomic dysfunction and poor neurodevelopment.^{11,12} Alcohol passed through the placenta is known to cause oxidative stress, mitochondrial damage, and alterations in the developing brain including malformations in the corpus callosum and cerebellum.^{13–15} Nicotine alters fetal brain development by effects on cell proliferation and differentiation, specific neurotransmitters receptors, and disruption of neural activity in the brain.¹⁶ Tobacco smoking and alcohol can also

Received: 3 October 2021 Revised: 22 March 2022 Accepted: 24 March 2022 Published online: 19 April 2022

¹Department of Psychiatry, Columbia University Irving Medical Center, New York, NY 10032, USA. ²Division of Developmental Neuroscience, New York State Psychiatric Institute, New York, NY 10032, USA. ³Department of Statistics, Columbia University, New York, NY 10032, USA. ⁴Avera Research Institute, Sioux Falls, SD 57108, USA. ⁵Department of Pediatrics, University of South Dakota School of Medicine, Sioux Falls, SD 57069, USA. ⁶Department of Obstetrics and Gynaecology, Faculty of Medicine and Health Science, Stellenbosch University, Cape Town, Western Cape 7530, South Africa. ⁷Department of Pediatrics, Columbia University Irving Medical Center, New York, NY 10032, USA. ⁸These authors contributed equally: Ayesha Sania, Michael M. Myers. A complete list of non-author contributors appears in the Acknowledgments. ^{\Beginet} established university.

Impairment in the ANS regulation (cardiac, respiratory, and vascular) has been associated with SIDS, and adverse neurodevelopmental outcomes such as autism and ADHD.^{5,19,20} Because of the increased incidence of SIDS among infants exposed to prenatal drinking and smoking,²¹ autonomic alterations associated with these exposures are important to understand. In addition to previous reports on alterations in SIDS victims' brainstem regions involved in autonomic regulation,²² members of the Prenatal and Alcohol in SIDS and Stillbirth (PASS) research team have reported on smoking and drinking associated alterations in the brainstem locations involved in cardiorespiratory functions and arousal in SIDS infants.²³ Other studies have shown that infants who subsequently die of SIDS have altered patterns of ANS activity during sleep including higher heart rate (HR), reduced HR variability (HRV), and abnormalities in the beat-to-beat dynamics.^{24–30} Inadequate responses to hypotension while asleep may also play a role in the fatal SIDS event.^{31,32}

Previous studies have reported significant associations of prenatal drinking and smoking with reduced HRV in the fetus, and diminished HR responses to head-up tilting in newborns.^{8,33,34} These studies have some methodological limitations including small sample sizes and limited exposure and covariate data. In the present study, we examined the role of prenatal drinking and smoking on parameters of ANS control in newborn infants in two large ethnically diverse cohorts of mother–infant dyads from the USA and South Africa. We hypothesized that prenatal drinking and smoking would affect newborn autonomic function and these effects will be contingent on timing, pattern, and quantity of exposure. However, given the paucity of data with detailed measures of exposure as in the current study. we had no specific directional hypotheses.

METHODS

Study population and data collection

Data for this study come from the Prenatal and Alcohol in SIDS and Stillbirth (PASS) Safe Passage Study. The Safe Passage Study was a prospective cohort study designed to evaluate the role of prenatal drinking and smoking on adverse pregnancy outcomes including stillbirth, SIDS, and fetal alcohol spectrum disorders (FASD). Participants were enrolled from antenatal clinics in Northern Plains, USA (South and North Dakota) and Cape Town, South Africa between August 2007 and January 2015. Pregnant women aged 16 years or older, at or after 6 weeks of gestation, carrying one or two fetuses were eligible to participate. Data on socioeconomic status, demographic, obstetric and medical history, periconceptional drinking and smoking were collected at the enrollment interview. Details on the study design and characteristics of the study population are published elsewhere.³⁵

Exposures assessment and characterization

Data on alcohol exposure were collected using a validated modification of the timeline follow back method,³⁶ where, at each study visit, participants reported detailed information on daily drinking ±15 days of their last menstrual period at enrollment visit and 30 days prior to their last known drinking day at each study visit. Participants also reported on the number of days they smoked per week and number of cigarettes smoked per day at enrollment and during the prenatal visits. Depending on the number and timing of the study visits for each participant, some data on daily drinking and smoking were missing. Missing data were imputed using a non-parametric algorithm called K nearest neighbor algorithm (k-NN). Using clustering methods, based on the amount and timing of drinking and smoking, the participants were divided into 4 smoking and 10 drinking groups.³⁸ Because several of the drinking groups had a small number of subjects they were collapsed into four groups for the current analyses. Women were categorized as moderate-to-high continuous smoking, lowcontinuous smoking, quit-early smoking, and non-smoking; and moderateto-high continuous drinking, low-continuous drinking, quit-early drinking, and non-drinking groups (see Supplementary Methods on exposure imputation and exposure clustering).

Table 1. Characteristics of the study population (n = 6915) by site.

| | Cape Town, South Africa (N = 4072) | Northern Plains, USA (<i>N</i> = 2913) |
|---|--|--|
| | N ¹ (%) or mean (SD) | N ¹ (%) or mean (SD) |
| Maternal characteristics | | |
| Maternal age, years | 24.7 (5.8) | 26.8 (5.3) |
| Marital status | | |
| Married/cohabiting | 2195 (53.9) | 2290 (78.6) |
| Unmarried | 1877 (46.1) | 623 (21.4) |
| Education | | |
| Primary school education | 313 (7.7) | 46 (1.6) |
| Some high school education | 2738 (67.2) | 480 (16.5) |
| High school completed | 876 (21.5) | 531 (18.2) |
| Higher than high school | 142 (3.5) | 1855 (63.7) |
| Employment status | | |
| Employed | 1183 (29.1) | 1914 (65.1) |
| Unemployed | 2889 (70.9) | 999 (34.3) |
| BMI, kg/m ² | | |
| <18.5 | 287 (7.1) | 61 (2.1) |
| 18.5–25 | 2010 (49.4) | 1045 (35.9) |
| 25–30 | 957 (23.5) | 877 (30.1) |
| 30+ | 818 (20.1) | 930 (31.9) |
| Parity | | |
| Nulliparous | 1597 (38.8) | 935 (32.1) |
| Parity 1–2 | 1265 (31.1) | 937 (32.2) |
| Parity >3 | 739 (18.2) | 546 (18.7) |
| Mode of delivery | | |
| Normal and assisted delivery | 3650 (89.70) | 2234 (76.72) |
| Cesarean section | 419 (10.30) | 678 (23.28) |
| Crowding index | 1.6 (0.9) | 0.7 (0.6) |
| Depression, Edinburgh Depression Score | 12.77 (5.92) | 5.46 (4.34) |
| Infant characteristics | | |
| Birth weight, g | 3107.7 (447.3) | 3520. 9 (457.3) |
| Gestational age, weeks | 39.4 (1.2) | 39.4 (1.1) |
| Sex | | |
| Female | 1985 (48.8) | 1469 (50.4) |
| Male | 2087 (51.2) | 1444 (49.6) |
| Hours of life at measurement | 60.26 (51.27) | 29.93 (47.56) |

ANS data measurement and data processing

Details on ANS measurement, processing and normative values in our study population have been published previously.³⁹ Newborns were tested 12–96 h after delivery. Electrocardiograms (ECG) and respiration rates were collected using custom hardware and software, and blood pressure (BP) was collected using a clinical monitor. The standard protocol involved recording ECG, respiration rate, and BP signals during a 10-min baseline period and in response to three rapid (~3–5 s) 45° head-up tilts while the infant was in the prone position. The resting state measures allowed the characterization of autonomic traits, whereas changes in autonomic state in response to tilt assessed the infant's capacity to respond to an

244

environmental challenge. As cardiorespiratory variables differ between sleep states, the records were subdivided into epochs coded as either active or quiet sleep states.^{40,41}

Baseline parameters. Baseline physiological variables were computed for 10 1-minute epochs prior to two baseline BP measurements. For each minute, the median the median R-wave to R-wave interval (RRi) was computed, and from the median RRi, HR was computed. The standard deviations of RRi (SD-RRi) and the square root of the mean of the squared successive differences in RRis (rMSSD) were computed as were lowfrequency HRV (LF; HRV between 0.04 and 0.40 Hz) and high-frequency HRV (HF; HRV between 0.50 and 1.50 Hz) power from Fourier spectral analyses. The LF/HF ratio was also computed. For each minute, the median breath-to-breath interval, and breathing rate were computed. The longterm variability parameters (SD, LF power) evaluate a combination of sympathetic and parasympathetic nervous systems contributions to HRV. In contrast, measures of beat-to-beat variability (RMSSD, HF) are influenced largely by the parasympathetic nervous system, tied to rapid ANS reactivity. The LF/HF ratio has been interpreted as a measure of sympatho-vagal balance, with higher values indicating sympathetic predominance and predominance.⁴² indicating lower values parasympathetic

Parameters in responses to head-up tilting. For HR, acute responses to tilts were computed as maximum HR-minimum HR during the first 15 s in the head-up position. The median maximum-minimum HR over the three tilts were analyzed. The pre-tilt values for systolic and diastolic BP were recorded 90 s before tilts and head-up BP values were recorded during the period just before returning to the horizontal position, 90 s after head-up tilting. Median changes in BP over the three tilts were analyzed.

Statistical analysis

Linear regression models were used to estimate associations of each prenatal drinking groups (reference: non-drinkers) and smoking groups (reference: non-smokers) with ANS parameters with separate models for each of the ANS outcomes. All models included infant sex, hours of life at assessment and gestational age (GA) at birth as covariates. Models included maternal education, household crowding index, prenatal depression scores measured with the Edinburgh scale, and study site as potential confounders. Missing indicators were included in the multivariate models when covariate data were missing. Outcome variables that were not normally distributed were log transformed (SD-RRi, RMSSD). Analyses combining participants across sites were performed first, and then, separately for each of the two sites. Additional sex stratified analyses were also performed. Because there were site differences in key characteristics of the participants, study site was included as a covariate in site combined models. Additional analyses stratified by site were also obtained. The overlap between exposure clusters (four smoking groups and four drinking groups) was insufficient to examine the effects of interaction between drinking and smoking on ANS outcomes. Because we have related outcomes and covariates, adjustment for multiple comparisons (such as the Bonferroni method) was not appropriate.⁴ We have reported all analyses and estimates (and confidence intervals) with equal emphasis and detail in the manuscript. All analyses were performed in SAS software version 9.4 (SAS Institute, Cary NC).

Ethics statement

Written informed consent was obtained from all participants. Ethical approval was obtained from Stellenbosch University, Avera Health, Sanford Health, the Indian Health Service and from participating Tribal Nations.

RESULTS

The final analyses included 6985 singletons born at term, of them 2913 (42%) were from Northern Plains and 4072 (58%) were from South Africa. We excluded infants with missing exposure data, preterm and post-term infants, and infants whose mothers used psychiatric medication during pregnancy (Supplementary Fig. S1). South African women had poorer socioeconomic status, as reflected in lower educational attainment, higher unemployment rates, higher crowding index, and higher proportion living without a partner. Although mean GAs at birth were similar in both sites, South African newborns were about 400 g lighter on average (Table 1). ANS parameters were obtained later in South Africa (60. 3 h, SD 51.3) compared to the Northern Plains site (29.9 h, SD 47.3). This difference was due the hospital policy in South Africa of discharging infants within 24 h of delivery, so they needed to be brought in for assessment later.

The prevalence of smoking was higher in South Africa (61%) than in Northern Plains (24%). In South Africa, 23% pregnant women continued smoking in moderate to high amount and 35% continued with a low amount throughout the pregnancy while only 17% woman in Northern Plains continued. Women in the moderate-to-high continuous, mild continuous, and quit early groups smoked 48.31 (SD 21.71), 15.72 (SD 10.28), and 8.81 (SD 9.92) cigarettes/week, on average in first trimester, respectively (Fig. 1 and Supplementary Table 1), with a declining number of cigarettes smoked in subsequent trimesters in all groups. Approximately half of the pregnant women in both sites reported drinking during pregnancy. The majority (43%) of NP participants guit drinking in the first trimester, only 8% drank throughout pregnancy, while 29% participants from SA continued to drink throughout (Supplementary Table S3). Women in the high moderate-to-continuous, mild continuous, and guit early groups drank 40.9 (SD 60.1), 2.4 (SD 3.8), and 8.9 (SD 7.4) drinks in first trimester, respectively (Fig. 1 and Supplementary Table 3). Women in the moderate-to-high continuous drinking group reported 2 (interguartile range (IQR) 0-5), 1 (IQR 0-2), and 0 (IQR 0-1) binge episodes in trimester 1, 2, and 3 respectively. Women in low continuous and quit early group rarely binged.



Fig. 1 Magnitude of smoking and drinking in each trimester by clusters.

| | Non-smoker | Quit early | Low continuous | Moderate to high continuous |
|--|-------------------------|-----------------------------|-----------------------------|--------------------------------|
| | Mean (95% CI) | Mean difference (95% CI) | Mean difference (95% Cl) | Mean difference (95% Cl) |
| | | P value | P value | P value |
| HR (bpm) | 126.13 (125.41, 126.84) | 1.07 (-0.72, 2.87) | 0.096 (-0.88, 1.07) | -0.019 (-1.21, 1.16) |
| | | (0.24) | (0.84) | (0.97) |
| log ₁₀ SD-RRi (s) | -1.61 (-1.62, -1.59) | -0.005 (-0.04, 0.02) | 0.002 (-0.014, 0.02) | 0.02 (-0.001, 0.04) |
| | | (0.72) | (0.20) | (0.06) |
| log ₁₀ RMSSD (s) | -1.95 (-1.96, -1.94) | -0.027 (-0.060, 0.0005) | -0.0002 (0.02, 0.01) | 0.015 (-0.007, 0.04) |
| | | (0.10) | (0.98) | (0.17) |
| Breaths/min | 56.05 (55.29, 56.82) | -1.09 (-3.03, 0.84) | 0.79 (-0.25, 1.85) | 2.22 (0.95, 3.49) |
| | | (0.26) | (0.13) | (0.0006) |
| Systolic blood pressure (mmHg) | 83.12 (81.96, 84.28) | 0.88 (-2.15, 3.91) | 0.44 (-1.11, 1.98) | 1.24 (-0.65, 3.14) |
| | | (0.56) | (0.57) | (0.19) |
| Diastolic blood pressure (mmHg) | 50.82 (49.86, 51.77) | 1.44 (-1.03, 3.92) | 0.87 (-0.39, 2.14) | 1.53 (-0.02, 3.09) |
| | | (0.25) | (0.17) | (0.053) |
| Low-frequency power of HRV | -5.88 (-5.90, -5.86) | 0.0003 (-0.059, 0.059) | 0.002 (-0.029, 0.034) | 0.03 (-0.003, 0.07) |
| | | (0.992) | (0.881) | (0.075) |
| High-frequency power of HRV | -6.83 (-6.85, -6.80) | -0.077 (-0.14, -0.011) | 0.009 (-0.026, 0.045) | 0.04 (-0.0013, 0.085) |
| | | (0.02) | (0.59) | (0.057) |
| Ratio of low and high-frequency power of | 0.94 (0.92, 0.97) | 0.05 (-0.007, 0.11) | 0.006 (-0.025, 0.038) | 0.013 (-0.025, 0.051) |
| HRV | | (0.08) | (0.709) | (0.504) |
| Quiet sleep | | | | |
| HR (bpm) | 120.53 (119.39, 121.67) | -0.49 (-3.15, 2.17) | 0.58 (-1.04, 2.19) | 1.03 (-1.04, 3.11) |
| | | (0.71) | (0.48) | (0.32) |
| log ₁₀ SD-RRi (s) | -1.74 (-1.76, -1.72) | 0.05 (-0.002, 0.10) | -0.01 (-0.05, 0.02) | 0.02 (-0.02, 0.06) |
| | | (0.06) | (0.43) | (037) |
| log ₁₀ RMSSD (s) | -1.93 (-1.95, -1.90) | 0.009 (-0.04, 0.07) | -0.017 (-0.05, 0.016) | 0.007 (-0.036, 0.051) |
| | | (0.73) | (0.31) | (0.74) |
| Breaths/min | 45.68 (44.65, 46.72) | -1.78 (-4.19, 0.63) | 0.16 (-1.31, 1.63) | 1.70 (-0.18, 3.58) |
| | | (0.14) | (0.82) | (0.07) |
| Systolic blood pressure (mmHg) | 82.75 (81.13, 84.38) | -0.29 (-4.01, 3.41) | 0.72 (-1.52, 2.97) | 0.06 (-2.75, 2.87) |
| | | (0.87) | (0.52) | (0.96) |
| Diastolic blood pressure (mmHg) | 50.17 (48.97, 51.37) | -0.70 (-3.45, 2.03) | -0.21 (-1.87, 1.45) | -1.32 (-3.40, 0.74) |
| | | (0.61) | (0.80) | (0.21) |
| Low-frequency power of HRV | -6.27 (-6.32, -6.21) | 0.12 (0.00074, 0.24) | -0.029 (-0.103, 0.043) | 0.056 (-0.036, 0.15) |
| | | (0.04) | (0.424) | (0.234) |
| High-frequency power of HRV | -6.84 (-6.89, | -0.04 (-0.149, 0.067) | -0.037 (-0.103, 0.028) | 0.028 (-0.055, 0.113) |
| | -6.80) | (0.45) | (0.26) | (0.506) |
| Ratio of low- and high-frequency | 0.58 (0.53, 0.63) | 0.117 (-0.003, 0.238) | 0.015 (-0.057, 0.087) | 0.039 (-0.054, 0.134) |
| power of HRV | | (0.056) | (0.68) | (0.40) |

Infants whose mothers did not smoke during pregnancy formed the reference group. Results with p values <0.01, <0.05, <0.10 are bolded.

Associations of smoking with baseline ANS parameters (Table 2 and Supplementary Tables S5 and S6)

In the site combined data analyses, compared to non-smokers, moderate-to-high-continuous smoking was associated with an increase in breathing rate in both sleep states (p < 0.001 and p = 0.07 respectively, Table 2). Similar results were seen for active sleep at each site (SA: 2.00 breaths/min, 95% CI: 0.43, 3.56; NP: 2.38 95% CI: -0.15, 4.91; Tables S-5 and S-6). However, this association was primarily driven by results for girls (Fig. 2a–c).

There were no significant associations of smoking with HR parameters in the analyses with sites combined. In SA, compared to infants of non-smoker mothers, SD-RRi in active sleep was 0.02 s (95% Cl: 0.002, 0.04) greater in the moderate-to-high-continuous smoking group (Supplementary Table S5), while rMSSD in active sleep (-0.08 s, 95% Cl: -0.14, -0.03) and quiet sleep (-0.11, 95% Cl: -0.22, -0.001) was decreased in the quit-early smoking group (Tables S-5 and S-6 respectively). In NP, SD-RRi in quiet sleep was increased in the quit-early group (p = 0.02; Supplementary Table S6).



Fig. 2 Marginal means in each exposure categories, stratified by study site and infant sex. a–c Median respiration rate for smoking exposure groups for both sites, South Africa, and Northern Plains, respectively. d–f Systolic BP for smoking exposure groups for both sites, South Africa and Northern Plains, respectively. g-i Diastolic BP for smoking exposure groups for both sites, South Africa and Northern Plains, respectively. j-l Ratio of low- to high-frequency heart rate variability for alcohol exposure groups for both sites, South Africa, and Northern Plains, respectively. *p < 0.1, **p < 0.05, ***p < 0.01 indicates significance of the difference in marginal mean in comparison to the reference group, non-smoker, or non-drinker.

There were no significant associations of smoking with diastolic BP in combined analyses. In sex stratified analyses, moderate-tohigh continuous smoking was associated with higher systolic (2.97 mmHg, Cl: 0.70, 5.24, p = 0.01) and diastolic BP (2.87 mmHg, Cl: 0.72, 5.02, p = 0.009) among girls only (Fig. 2d, g).

Associations of alcohol with Baseline ANS parameters (Table 3 and Supplementary Tables S7 and S8)

In site combined analyses, compared to infants of non-drinker mothers, infants in the low-continuous drinking group had 4.53 beats/min lower HR in quiet sleep (95% CI: -7.75, -1.32; Table 3). Similar results were seen in SA for the low-continuous and moderate-to-high continuous drinking group (Supplementary Table S7). Compared to no drinking, low-continuous drinking was associated with a reduced ratio of LF/HF in active sleep in combined analyses (-0.08 95% CI -0.15, -0.02; Table 3), as well as in site stratified analyses (Tables S-7 and S-8). Further

stratification by sex revealed that this association was significant among boys only (Fig. 2j-l).

In combined analyses, diastolic BP during quiet sleep was lower in low-continuous (-3.29 mmHg, 95% Cl: -6.59, -0.006) and quitearly (-1.36 mmHg, 95% Cl: 2.78, 0.05) drinking groups (Table 3) compared to non-drinkers. Low-continuous drinking was associated with reduced diastolic BP in the NP (p = 0.04; Supplementary Table S8) but not in SA.

Associations of smoking on tilt responses (Table 4 and Supplementary Tables S9 and S10)

The decrease in active sleep breathing rates following head-up tilt was less pronounced in low-continuous smoking group (0.81 breaths/min, 95% Cl: 0.07, 1.54, Table 4) compared to non-smokers. In stratified analyses, this association was significant in SA (Supplementary Table S9) but not in the NP (Supplementary Table S10).

Table 3. Association of prenatal alcohol with ANS measures at baseline in site combined analyses.

| | Non-mean (95%Cl) | Quit early | Low continuous | Moderate to high continuous |
|----------------------------------|-------------------------|-----------------------------|-----------------------------|-----------------------------|
| | | Mean difference (95% Cl) | Mean difference (95% Cl) | Mean difference (95% Cl) |
| | | P value | P value | P value |
| Active sleep | | | | |
| HR (bpm) | 126.36 (125.67, 127.05) | 0.46 (-0.37, 1.30) | -0.76 (-2.74, 1.22) | 0.56 (-0.54, 1.67) |
| | | (0.27) | (0.45) | (0.31) |
| log ₁₀ SD-RRi (s) | -1.61 (-1.61, -1.59) | -0.005 (-0.02, 0.009) | 0.004 (-0.03, 0.04) | -0.003 (-0.02, 0.015) |
| | | (0.51) | (0.80) | (0.73) |
| log ₁₀ RMSSD (s) | -1.96 (-1.97, -1.94) | -0.005 (-0.02, 0.01) | 0.022 (-0.01, 0.58) | 0.006 (-0.01, 0.03) |
| | | (0.49) | (0.23) | (0.53) |
| Breaths/min | 56.05 (55.28, 56.82) | -0.14 (-1.03, 0.75) | -1.61 (-3.72, 0.50) | -0.45 (-1.64, 0.73) |
| | | (0.76) | (0.13) | (0.45) |
| Systolic blood pressure (mmHg) | 84.22 (83.09, 85.35) | -0.21 (-1.55, 1.13) | -1.19 (-4.43, 2.05) | -0.43 (-2.19, 1.31) |
| | | (0.31) | (0.47) | (0.62) |
| Diastolic blood pressure (mmHg) | 52.04 (51.12, 52.97) | 0.22 (-0.87, 1.33) | -0.79 (-3.45, 1.87) | -0.48 (-1.92, 0.95) |
| | | (0.68) | (0.56) | (0.50) |
| Low-frequency power of HRV | -5.86 (-5.88, -5.84) | -0.015 (-0.043, 0.011) | -0.017 (-0.082, 0.048) | -0.013 (-0.05, 0.022) |
| | | (0.26) | (0.609) | (0.45) |
| High-frequency power of HRV | -6.86 (-6.89, -6.84) | -0.007 (-0.038, 0.022) | 0.089 (0.017, 0.16) | 0.036 (-0.004, 0.077) |
| | | (0.62) | (0.015) | (0.079) |
| Ratio of low- and high-frequency | 1.002 (0.98, 1.02) | -0.014 (-0.04, 0.01) | -0.08 (-0.15, -0.02) | -0.04 (-0.07, -0.006) |
| power of HRV | | (0.31) | (0.008) | (0.02) |
| Quiet sleep | | | | |
| HR (bpm) | 122.28 (121.16, 123.41) | 0.07 (-1.28, 1.42) | -4.53 (-7.75, -1.32) | -1.41 (-3.30, 0.46) |
| | | (0.10) | (0.006) | (0.13) |
| log ₁₀ SD-RRi (s) | -1.72 (-1.74, -1.69) | -0.001 (-0.03, 0.02) | -0.02 (-0.08, 0.05) | 0.002 (-0.03, 0.04) |
| | | (0.92) | (0.61) | (0.92) |
| log ₁₀ RMSSD (s) | -1.94 (-1.96, -1.91) | -0.006 (-0.04, 0.02) | 0.006 (-0.06, 0.08) | 0.017 (-0.022, 0.06) |
| | | (0.64) | (0.85) | (0.39) |
| Breaths/min | 45.84 (44.82, 46.86) | -0.51 (-1.73, 0.71) | -0.54 (-3.45, 2.36) | 0.48 (-1.22, 2.20) |
| | | (0.41) | (0.71) | (0.57) |
| Systolic blood pressure (mmHg) | 84.02 (82.48, 85.55) | -1.22 (-3.14, 0.70) | -3.46 (-7.92, 0.98) | 0.13 (-2.38, 2.65) |
| | | (0.21) | (0.12) | (0.10) |
| Diastolic blood pressure (mmHg) | 43.61 (40.65, 46.56) | -1.36 (-2.78. 0.05) | -3.29 (-6.59, 0.006) | 0.99 (-0.86. 2.84) |
| | | (0.06) | (0.05) | (0.29) |
| Low-frequency power of HRV | -6.23 (-6.28, | 0.015 (-0.04, 0.07) | -0.02 (-0.17, 0.11) | 0.003 (-0.08, 0.08) |
| | -6.18) | (0.62) | (0.72) | (0.93) |
| High-frequency power of HRV | -6.86 (-6.90, | -0.02 (-0.08, 0.02) | -0.014 (-0.14, 0.11) | 0.04 (-0.03, 0.11) |
| | -6.81) | (0.32) | (0.82) | (0.28) |
| Ratio of low- and high-frequency | 0.61 (0.56, 0.67) | 0.03 (-0.02, 0.09) | 0.02 (-0.12, 0.16) | -0.02 (-0.11, 0.05) |
| power of HRV | | (0.22) | (0.75) | (0.53) |

Infants whose mothers did not drink during pregnancy formed the reference group. Results with p values <0.01, <0.05, <0.10 are bolded.

In site combined analyses, the acute HR responses to tilt were increased during active sleep in the low-continuous (1.03 beats/min, 95% CI: 0.23, 1.83) and the moderate-to-high-continuous (1.22 beats/min, 95% CI: 0.23, 2.21) smoking groups compared to non-smokers (Table 4). The increases in the acute HR response in the moderate-to-high-continuous smoking group was significant in SA only (p = 0.02, SupplementaryTable S9). The decrease in RMSSD following head up tilt seen in the non-smoking (reference group) during active sleep was not seen in the low-continuous smoking group (0.35 s, 95% CI: 0.03, 0.67, Table 4). A less marked

reduction of RMSSD values in quiet sleep during head-up tilt was found in quit-early smoking group (p = 0.05, Table 4) in site combined analyses.

In combined sites analyses, there were smaller decreases in systolic (3.05 mmHg, 95% Cl: 1.22, 4.89) and diastolic BP (1.95 mmHg, 95% Cl: 0.31, 3.58) following head up tilts during active sleep in the quit-early smoking group compared to non-smoker group (Table 4). These differences remained significant in NP (Supplementary Table S10). Decreases in diastolic BP to tilt in active sleep were also diminished in the low-continuous smoking

Table 4. Association of prenatal smoking with mean changes in cardiorespiratory variables in response to 45° head-up tilts in site combined analyses.

| | Non-smoker | Quit early | Low continuous | Moderate to high continuous |
|---|-----------------------|-----------------------------|-----------------------------|-----------------------------|
| | Mean (95% Cl) | Mean difference (95% Cl) | Mean difference (95% Cl) | Mean difference (95% CI) |
| | | (P value) | (P value) | (P value) |
| Active sleep | | | | |
| Δ sustained HR (bpm) | 2.52 (2.08, 2.96) | 0.56 (-1.64, 0.52) | -035 (-0.95, 0.24) | -0.29 (-1.03, 0.45) |
| | | (0.31) | (0.24) | (0.44) |
| Δ acute HR (bpm) | 22. 98 (22.39, 23.57) | -0.69 (-2.13, 0.75) | 1.03 (0.23, 1.83) | 1.22 (0.23, 2.21) |
| | | (0.34) | (0.01) | (0.02) |
| Δ rMSSD-RRi (s) | -0.57 (-0.80, -0.33) | 0.46 (-0.12, 1.04) | 0.35 (0.03, 0.67) | -0.09 (-0.49, 0.30) |
| | | (0.12) | (0.03) | (0.64) |
| Δ change in breathing (breaths/ | -3.73 (-4.27, -3.19) | 0.35 (-0.98, 1.68) | 0.81 (0.07, 1.54) | -0.005 (-0.91, 0.90) |
| min) | | (0.60) | (0.03) | (0.99) |
| Δ systolic blood | -6.04 (-6.76, -5.30) | 3.05 (1.22, 4.89) | 0.68 (-0.29, 1.65) | -0.22 (-1.42, 0.98) |
| pressure (mmHg) | | (0.001) | (0.16) | (0.72) |
| Δ diastolic blood | -6.49 (-7.31, -5.83) | 1.95 (0.31, 3.58) | 0.82 (-0.03, 1.69) | 0.08 (-0.98, 1.14) |
| pressure (mmHg) | | (0.01) | (0.06) | (0.88) |
| Quiet sleep | | | | |
| Δ sustained HR (bpm) | 3.39 (2.82, 3.98) | 0.89 (-0.60, 2.38) | -0.38 (1.19, 0.0.41) | 0.34 (-0.65, 1.32) |
| | | (0.24) | (0.34) | (0.50) |
| Δ acute HR (bpm) | 22.39 (21.54, 23.24) | -1.89 (-4.11 <i>,</i> 0.32) | 0.08 (-1.08, 1.24) | 0.40 (-1.03, 1.83) |
| | | (0.09) | (0.89) | (0.85) |
| Δ rMSSD-RRi (s) | -1.06 (-1.41, -0.71) | -0.92 (-1.82, -0.015) | 0.12 (-0.36, 0.61) | 0.27 (-0.32, 0.86) |
| | | (0.05) | (0.61) | (0.36) |
| Δ change in breathing (breaths/ min) | -3.73 (-4.27, -3.19) | 0.30 (-1.19, 1.79) | -0.36 (-1.17, 0.46) | -0.23 (-1.22, 0.77) |
| | | (0.69) | (0.39) | (0.65) |
| Δ systolic blood | -5.07 (-5.89, -4.25) | -0.19 (-2.38, 2.00) | -0.02 (-1.12, 1.07) | 0.37 (-0.95, 1.69) |
| pressure (mmHg) | | (0.86) | (0.97) | (0.58) |
| Δ diastolic blood | -5.83 (-6.53, -5.12) | -0.24 (-2.08, 1.61) | 0.34 (-0.57, 1.26) | 0.32 (-0.78, 1.43) |
| pressure (mmHg) | | (0.80) | (0.46) | (0.57) |

Infants whose mothers did not smoke during pregnancy formed the reference group. Results with p values <0.01, <0.05, <0.10 are bolded.

group compared to non-smoking group, in site combined analyses (0.82 mmHg, 95% Cl: -0.03, 1.69, Table 4) and in NP (2.57, 95% Cl: 0.51, 4.63, Supplementary Table 510).

Associations of alcohol on tilt responses (Table 5 and Supplementary Tables S11 and S12)

The decrease in breathing rate in active sleep following head up tilting was less pronounced in the low-continuous drinking group (1.81 breaths/min, 95% Cl: 0.32, 3.29, Table 5) compared to nondrinking group. In stratified analyses this association remained significant in NP but not in SA. In the NP, decreases in breathing in quiet sleep to head up tilt were diminished among low-continuous drinking (4.55 breaths/min, 95% Cl: 0.19, 8.91, Supplementary Table S12),

Compared to non-drinking group, the decrease in RMSSD during quiet sleep was diminished in the quit-early drinking group (0.44 s, 95% CI: 0.03, 0.84, Table 5) but not in site stratified analyses (Tables S-11 and S-12).

Compared to non-drinking group, decreases in systolic BP in active sleep following head up tilts were less pronounced in the moderate-to-high-continuous drinking group in site combined analyses (1.22 mmHg, 95% CI: 0.11, 2.33, Table 5), which remained significant in SA (Supplementary Table S11), but not in the NP (Supplementary Table S12). In NP, decreases in diastolic BP in quiet sleep were diminished among moderate-to-high-continuous drinking group (4.36 mmHg, 95% Cl: 1.41, 7.30, Supplementary Table S12).

DISCUSSION

In this study, data from two large socio-culturally and ethnically diverse cohorts were used to evaluate the effects of prenatal drinking and smoking on ANS parameters extracted from physiological signals acquired from term newborns at birth. We found significant associations of prenatal smoking with newborn breathing and rMSSD during baseline, as well as greater acute changes in HR, and diminished changes in systolic and diastolic BP in response to head up tilts. There was a significant association between smoking and blood pressure among girls only. Prenatal drinking was associated with reductions in BP, reductions in the ratio of low- to high-frequency HRV, and diminished breathing rate and systolic BP responses to tilt. The associations of prenatal smoking and drinking were more pronounced in South African newborns, where women, on average, drank and smoked at higher levels, and were more likely to continue to drink and smoke throughout the pregnancy than women in the Northern Plains.

Moderate to high continuous smoking was associated with higher breathing rates. This extends the literature linking prenatal

| 2 | /. | o |
|---|----|---|
| 2 | 4 | 7 |

Table 5. Association of prenatal drinking with mean changes in cardiorespiratory variables in response to 45° head-up tilts in site combined analyses.

| | Non-drinker | Quit early | Low continuous | Moderate to high continuous |
|--|----------------------|-----------------------------|-----------------------------|-----------------------------|
| | Mean (95% CI) | Mean difference (95% CI) | Mean difference (95% Cl) | Mean difference (95% CI) |
| | | (P value) | (P value) | (P value) |
| Active sleep | | | | |
| Δ sustained HR (bpm) | 2.29 (1.86, 2.71) | -0.21 (-0.72, 0.28) | -0.22 (-1.45, 1.00) | 0.17 (-0.51, 0.86) |
| | | (0.40) | (0.72) | (0.62) |
| Δ acute HR (bpm) | 23.08 (22.51, 23.64) | -0.23 (-0.91, 0.44) | 1.33 (-0.29, 2.95) | 0.09 (-0.83, 1.01) |
| | | (0.50) | (0.10) | (0.85) |
| Δ rMSSD-RRi (ms) | -0.34 (-0.57, -0.12) | -0.11 (-0.38, 0.16) | 0.09 (-0.56, 0.75) | -0.15 (-0.52, 0.22) |
| | | (0.42) | (0.77) | (0.42) |
| Δ change in breathing (breaths/ | -3.95 (-4.46, -3.43) | 0.47 (-0.14, 1.09) | 1.81 (0.32, 3.29) | -0.29 (-1.14, 0.55) |
| min) | | (0.12) | (0.02) | (0.49) |
| Δ systolic blood | -5.60 (-6.29, -4.90) | 0.15 (0.69, 1.01) | 0.39 (-1.57, 2.37) | 1.22 (0.11, 2.33) |
| pressure (mmHg) | | (0.31) | (0.69) | (0.03) |
| Δ diastolic blood | -6.09 (-6.71, -5.47) | 0.32 (-0.43, 1.07) | 0.39 (-1.35, 2.13) | 0.55 (-0.43, 1.54) |
| pressure (mmHg) | | (0.40) | (0.66) | (0.26) |
| Quiet sleep | | | | |
| Δ sustained HR (bpm) | 3.81 (3.24, 4.38) | -0.37 (-1.04, 0.31) | -0.82 (-2.42, 0.78) | 0.38 (-0.51, 1.28) |
| | | (0.28) | (0.31) | (0.39) |
| Δ acute HR (bpm) | 22.17 (21.33, 23.01) | -0.68 (-1.66, 0.31) | 0.29 (-2.08, 2.66) | -0.12 (-1.43, 1.17) |
| | | (0.17) | (0.80) | (0.84) |
| Δ rMSSD-RRi (ms) | -1.09 (-1.44, -0.75) | 0.44 (0.03, 0.84) | -0.58 (-1.55 0.38) | -0.21 (-0.75, 0.32) |
| | | (0.04) | (0.23) | (0.43) |
| Δ change in breathing (breaths/ | -2.39 (-2.97, -1.81) | -0.08 (-0.76, 0.61) | -1.09 (-2.71, 0.52) | -0.34 (-1.24, 0.56) |
| min) | | (0.83) | (0.18) | (0.46) |
| Δ systolic blood | -4.85 (-5.65, -4.04) | -0.34 (-1.32, 0.64) | -0.35 (-2.53, 1.83) | -0.05 (-1.27, 1.17) |
| pressure (mmHg) | | (0.50) | (0.75) | (0.93) |
| Δ diastolic blood | -5.81 (-6.50, -5.13) | -0.03 (-0.85, 0.79) | 0.11 (-1.72, 1.94) | 0.27 (-0.75, 1.30) |
| pressure (mmHg) | | (0.94) | (0.90) | (0.59) |
| | | | | |

Infants whose mothers did not drink during pregnancy formed the reference group. Results with p values <0.01, <0.05, <0.10 are bolded.

smoking to alterations in regulation of infant respiration, specifically regarding the literature reporting increases in the frequency and length of obstructive apneas.⁴⁶ Evidence from animal studies indicates maternal smoking leads to nicotine-induced cholinergic sensitivity of respiratory neurons of neonatal mice,⁴⁷ a mechanism that could predispose newborns to SIDS. Prenatal smoking has also been shown to adversely impact airway development, leading to abnormalities associated with an increased tendency to wheeze and higher susceptibility to respiratory infections in early childhood.⁴⁸

Our results support the conclusion that prenatal smoking alters key aspects of cardiovascular regulation in newborn infants. The finding that prenatal smoking is associated with increases in baseline BP is consistent with a prior report by Beratis et al.⁴⁹ and Browne et al.⁵⁰ With regard to the responses to tilt, in a prior study in the NP, smaller increases in HR following head up tilting were found in tobacco-exposed infants.⁸ Consistent with this, in our study, infants of non-smokers had significant increases in HR following tilt whereas infants exposed to smoking did not (see Table 4). However, in contrast, Browne et al. found infants of smoking mothers had a larger decrease in blood pressure to tilt,⁴⁴ though the number of subjects assessed in the newborn period was small (26 non-smokers, 24 smokers).

Ours is the first study to report a significant reduction in the ratio of low-to-high frequency HRV among infants whose mothers

drank throughout the pregnancy in moderate-to-high amounts. This ratio is interpreted as a measure of sympathovagal balance, although this remains controversial⁵¹ has been utilized as potential marker for social arousal and engagement.⁵² The changes found suggest that prenatal drinking is associated with a greater relative parasympathetic control of HR. In contrast, an increase in the ratio of low-to-high frequency HRV associated with prenatal smoking was reported in other smaller studies.^{53,54} Greater effects of smoking on the ratio of low-to-high HRV among boys is potentially related to their greater vulnerability, given that male infants who are more likely to die of SIDS.⁵⁵ Similar sex differences in autonomic regulation are also reported in animal studies.⁵⁶ Taken together, these results indicate that the effects of smoking and drinking are not mediated through a common neurophysiological or psychological mechanism.

We also found moderate-to-high continuous smoking was associated with increases in systolic and diastolic BP among newborn girls. Similar effects were reported in prior studies among adolescent girls.^{57,58} Unlike previous studies in newborns and fetuses,^{8,9} we found no statistically significant associations of prenatal drinking on HR and HRV in active sleep. However, the average amount of alcohol consumed in our study populations was considerably lower than in previous studies. We also did not observe any significant associations of smoking on newborn baseline HR despite significant associations with fetal HR among the same cohort.⁹

The findings related to first trimester exposures followed by cessation in the guit early groups are potentially caused by the impact of withdrawal from smoking and drinking on maternal autonomic function and psychology, and, in turn, effects of these changes on fetal and infant physiology. Smoking cessation during pregnancy is known to elevate maternal blood pressure.59,60 In addition, women quitting smoking and drinking may experience depressed mood, anxiety, and stress symptoms.^{61,62} Although we adjusted for maternal depression in the analyses, residual confounding is plausible. Another limitation of our study is measurement error in the exposure assessment. Since smoking and drinking were based on self-reports, it is likely that there was underreporting by some pregnant women exposed to antismoking and anti-drinking information⁶³ and this could have contributed to less robust findings. In addition, data on medications related to smoking cessation or heart arrhythmias were not collected; therefore, we could not account for these potential contributions in the analyses. Given the low proportion of women guitting smoking in both sites and low socioeconomic condition of the participants, it was unlikely that many would have access to such interventions.

Strengths of our study include extensive prospectively collected alcohol and smoking data with rigorous quality controls, which allowed us to evaluate the role of both quantity and timing of the exposure. To date, ours is the largest epidemiological study to examine the role of PAE and PTE on newborn ANS in a populationbased study. We measured multiple aspects of autonomic activities including HRV, BP, and breathing rates, which allowed evaluation of their coordinated activities. We also had detailed information on several potential confounders that were not accounted for in previous studies. While studies conducted in laboratory settings have precise measurements of smoking and alcohol exposures and increased confounder control, such findings are not directly generalizable to human population health. Our findings are generalizable to pregnant women and their infants of similar racial, ethnic, and socioeconomic backgrounds.

CONCLUSION

We observed that prenatal smoking and drinking alters autonomic function in term newborns. Specifically, effects of smoking on breathing rate, beat-to-beat HRV, and effects of drinking on lowto high-frequency HRV highlight the pathways through which prenatal drinking and smoking could be linked to risk of SIDS and neurodevelopmental disorders like ADHD.^{24-26,64} While the changes in physiological measurement we presented might not meet thresholds for what may be termed clinical significance, they do indicate changes in ANS regulation profiles. Given that SIDS occurs when multiple factors intersect (vulnerable infant, critical period, and exogenous trigger) even small shifts in physiology can move infants into high-risk categories. Similar shifts in physiology have been reported in other at risk populations such as preterm infants.⁶⁵ The current study provides new evidence to support the conclusion that prenatal exposure to smoking and alcohol can alter newborn autonomic function. Given the high prevalence of prenatal smoking and drinking and their known associations with postnatal outcomes such as SIDS, and poor neurodevelopment, early examination of ANS parameters can serve as a marker for identification of newborns at risk of adverse outcomes.

REFERENCES

- Handmaker, N. S. et al. Impact of alcohol exposure after pregnancy recognition on ultrasonographic fetal growth measures. *Alcohol. Clin. Exp. Res.* **30**, 892–898 (2006).
- Scheffers-van Schayck, T., Tuithof, M., Otten, R., Engels, R. & Kleinjan, M. Smoking behavior of women before, during, and after pregnancy: indicators of smoking, quitting, and relapse. *Eur. Addict. Res.* 25, 132–144 (2019).

- Popova, S., Lange, S., Probst, C., Gmel, G. & Rehm, J. Estimation of national, regional, and global prevalence of alcohol use during pregnancy and fetal alcohol syndrome: a systematic review and meta-analysis. *Lancet Glob. Health* 5, e290–e299 (2017).
- Lange, S., Probst, C., Rehm, J. & Popova, S. National, regional, and global prevalence of smoking during pregnancy in the general population: a systematic review and meta-analysis. *Lancet Glob. Health* 6, e769–e776 (2018).
- Elliott, A. J. et al. Concurrent prenatal drinking and smoking increases risk for SIDS: safe passage study report. *EClinicalMedicine* 19, 100247 (2020).
- Hamulka, J., Zielinska, M. A. & Chadzynska, K. The combined effects of alcohol and tobacco use during pregnancy on birth outcomes. *Rocz. Panstw. Zakl. Hig.* 69, 45–54 (2018).
- Kesmodel, U., Wisborg, K., Olsen, S. F., Henriksen, T. B. & Secher, N. J. Moderate alcohol intake during pregnancy and the risk of stillbirth and death in the first year of life. Am. J. Epidemiol. 155, 305–312 (2002).
- Fifer, W. P., Fingers, S. T., Youngman, M., Gomez-Gribben, E. & Myers, M. M. Effects of alcohol and smoking during pregnancy on infant autonomic control. *Dev. Psychobiol.* 51, 234–242 (2009).
- 9. Lucchini, M. et al. Effects of Prenatal Exposure to Alcohol and Smoking on Fetal Heart Rate and Movement Regulation. *Front Physiol.* **12**, 594605 (2021).
- Shuffrey, L. C. et al. Association between prenatal exposure to alcohol and tobacco and neonatal brain activity: results from the safe passage study. JAMA Netw. Open 3, e204714 (2020).
- Flak, A. L. et al. The association of mild, moderate, and binge prenatal alcohol exposure and child neuropsychological outcomes: a meta-analysis. *Alcohol Clin. Exp. Res.* 38, 214–226 (2014).
- Julvez, J. et al. Maternal smoking habits and cognitive development of children at age 4 years in a population-based birth cohort. *Int. J. Epidemiol.* 36, 825–832 (2007).
- Manzo-Avalos, S. & Saavedra-Molina, A. Cellular and mitochondrial effects of alcohol consumption. Int. J. Environ. Res. Public Health 7, 4281–4304 (2010).
- Brocardo, P. S., Gil-Mohapel, J. & Christie, B. R. The role of oxidative stress in fetal alcohol spectrum disorders. *Brain Res. Rev.* 67, 209–225 (2011).
- Jarmasz, J. S., Basalah, D. A., Chudley, A. E. & Del Bigio, M. R. Human brain abnormalities associated with prenatal alcohol exposure and fetal alcohol spectrum disorder. J. Neuropathol. Exp. Neurol. 76, 813–833 (2017).
- Cohen, G. et al. Perinatal exposure to nicotine causes deficits associated with a loss of nicotinic receptor function. *Proc. Natl Acad. Sci. USA* **102**, 3817–3821 (2005).
- 17. Philips, E. M. et al. Changes in parental smoking during pregnancy and risks of adverse birth outcomes and childhood overweight in Europe and North America: an individual participant data meta-analysis of 229,000 singleton births. *PLoS Med.* **17**, e1003182–e1003182 (2020).
- Mamluk, L. et al. Evidence of detrimental effects of prenatal alcohol exposure on offspring birthweight and neurodevelopment from a systematic review of quasiexperimental studies. *Int. J. Epidemiol.* 49, 1972–1995 (2020).
- 19. Musser, E. D. et al. Emotion regulation via the autonomic nervous system in children with attention-deficit/hyperactivity disorder (ADHD). J. Abnorm. Child Psychol. **39**, 841–852 (2011).
- Grzadzinski, R. et al. Pre-Symptomatic Intervention for Autism Spectrum Disorder (Asd): Defining a Research Agenda. J. Neurodevelopmental Disord. 13, 49 (2021).
- Duncan, J. R. et al. The effect of maternal smoking and drinking during pregnancy upon (3)H-nicotine receptor brainstem binding in infants dying of the sudden infant death syndrome: initial observations in a high risk population. *Brain Pathol.* 18, 21–31 (2008).
- Kinney, H. C., Richerson, G. B., Dymecki, S. M., Darnall, R. A. & Nattie, E. E. The brainstem and serotonin in the sudden infant death syndrome. *Annu Rev. Pathol.* 4, 517–550 (2009).
- 23. Vivekanandarajah, A. et al. Nicotinic receptors in the brainstem ascending arousal system in sids with analysis of pre-natal exposures to maternal smoking and alcohol in high-risk populations of the safe passage study. *Front. Neurol.* **12**, 636668–636668 (2021).
- 24. Schechtman, V. L. et al. Dynamic analysis of cardiac R-R intervals in normal infants and in infants who subsequently succumbed to the sudden infant death syndrome. *Pediatr. Res.* **31**, 606–612 (1992).
- Schechtman, V. L. et al. Heart rate variation in normal infants and victims of the sudden infant death syndrome. *Early Hum. Dev.* 19, 167–181 (1989).
- Schechtman, V. L., Harper, R. M., Wilson, A. J. & Southall, D. P. Sleep apnea in infants who succumb to the sudden infant death syndrome. *Pediatrics* 87, 841–846 (1991).
- Kluge, K. A. et al. Spectral analysis assessment of respiratory sinus arrhythmia in normal infants and infants who subsequently died of sudden infant death syndrome. *Pediatr. Res.* 24, 677–682 (1988).
- Kahn, A. et al. Sleep and cardiorespiratory characteristics of infant victims of sudden death: a prospective case-control study. *Sleep* 15, 287–292 (1992).

- 29. Kahn, A. et al. Sudden infant deaths: from epidemiology to physiology. *Forensic Sci. Int.* **130**, S8–S20 (2002).
- Sahni, R., Fifer, W. P. & Myers, M. M. Identifying infants at risk for sudden infant death syndrome. *Curr. Opin. Pediatr.* 19, 145–149 (2007).
- Myers, M. M., Burtchen, N., Retamar, M. O., Lucchini, M. & Fifer, W. P. In SIDS Sudden Infant and Early Childhood Death: The Past, the Present and the Future (eds. Duncan, J. R. & Byard, R. W.) (Adelaide (AU): University of Adelaide Press, 2018).
- Horne, R. S. C. Cardiovascular autonomic dysfunction in sudden infant death syndrome. *Clin. Auton. Res* 28, 535–543 (2018).
- Zeskind, P. S. & Gingras, J. L. Maternal cigarette-smoking during pregnancy disrupts rhythms in fetal heart rate. J. Pediatr. Psychol. 31, 5–14 (2005).
- Pini, N., Lucchini, M., Fifer, W. P., Myers, M. M. & Signorini, M. G. In 2018 40th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC). 5874–5877.
- Dukes, K. A. et al. The safe passage study: design, methods, recruitment, and follow-up approach. *Paediatr. Perinat. Epidemiol.* 28, 455–465 (2014).
- Dukes, K. et al. A modified timeline followback assessment to capture alcohol exposure in pregnant women: application in the safe passage study. *Alcohol* 62, 17–27 (2017).
- 37. Sania, A. et al. The K nearest neighbor algorithm for imputation of missing longitudinal prenatal alcohol data. *Research Square* (2021).
- Pini, N. et al. Cluster analysis of alcohol consumption during pregnancy in the safe passage study. Annu Int Conf. IEEE Eng. Med Biol. Soc. 2019, 1338–1341 (2019).
- Myers, M. M. et al. Cardiorespiratory physiology in the safe passage study: protocol, methods and normative values in unexposed infants. *Acta Paediatr.* 106, 1260–1272 (2017).
- Isler, J. R., Thai, T., Myers, M. M. & Fifer, W. P. An automated method for coding sleep states in human infants based on respiratory rate variability. *Dev. Psychobiol.* 58, 1108–1115 (2016).
- Harper, R. M., Schechtman, V. L. & Kluge, K. A. Machine classification of infant sleep state using cardiorespiratory measures. *Electroencephalogr. Clin. Neurophysiol.* 67, 379–387 (1987).
- Malik, M. Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation* **93**, 1043–1065 (1996). B. J. a. C. J.
- Blakesley, R. E. et al. Comparisons of methods for multiple hypothesis testing in neuropsychological research. *Neuropsychology* 23, 255–264 (2009).
- 44. Greenland, S. Analysis goals, error-cost sensitivity, and analysis hacking: Essential considerations in hypothesis testing and multiple comparisons. *Paediatr Perinat Epidemiol.* **35**, 8–23 (2021).
- Rothman, K. J. No adjustments are needed for multiple comparisons. *Epidemiology* 1, 43–46 (1990).
- Kahn, A. et al. Prenatal exposure to cigarettes in infants with obstructive sleep apneas. *Pediatrics* 93, 778–783 (1994).
- Coddou, C., Bravo, E. & Eugenin, J. Alterations in cholinergic sensitivity of respiratory neurons induced by pre-natal nicotine: a mechanism for respiratory dysfunction in neonatal mice. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 364, 2527–2535 (2009).
- Milner, A. D., Rao, H. & Greenough, A. The effects of antenatal smoking on lung function and respiratory symptoms in infants and children. *Early Hum. Dev.* 83, 707–711 (2007).
- Beratis, N. G., Panagoulias, D. & Varvarigou, A. Increased blood pressure in neonates and infants whose mothers smoked during pregnancy. *J. Pediatr.* 128, 806–812 (1996).
- Browne, C. A., Colditz, P. B. & Dunster, K. R. Infant autonomic function is altered by maternal smoking during pregnancy. *Early Hum. Dev.* 59, 209–218 (2000).
- Eckberg, D. L. Sympathovagal balance: a critical appraisal. *Circulation* 96, 3224–3232 (1997).
- Sheinkopf, S. J. et al. Developmental trajectories of autonomic functioning in autism from birth to early childhood. *Biol. Psychol.* 142, 13–18 (2019).
- Nordenstam, F. et al. Prenatal exposure to snus alters heart rate variability in the infant. *Nicotine Tob. Res.* 19, 797–803 (2017).
- Franco, P., Chabanski, S., Szliwowski, H., Dramaix, M. & Kahn, A. Influence of Maternal smoking on autonomic nervous system in healthy infants. *Pediatr. Res.* 47, 215–220 (2000).
- Moscovis, S. M., Hall, S. T., Burns, C. J., Scott, R. J. & Blackwell, C. C. The male excess in sudden infant deaths. *Innate Immun.* 20, 24–29 (2014).
- Boychuk, C. R., Fuller, D. D. & Hayward, L. F. Sex differences in heart rate variability during sleep following prenatal nicotine exposure in rat pups. *Behav Brain Res.* 219, 82–91 (2011).
- 57. Zhang, Z. et al. Parental smoking and blood pressure in children and adolescents: a national cross-sectional study in China. *BMC Pediatrics* **19**, 116 (2019).
- Geerts, C. C. et al. Tobacco smoke exposure of pregnant mothers and blood pressure in their newborns. 50, 572–578 (2007).

- Bakker, R., Steegers, E. A., Mackenbach, J. P., Hofman, A. & Jaddoe, V. W. Maternal smoking and blood pressure in different trimesters of pregnancy: the generation R study. J. Hypertens. 28, 2210–2218 (2010).
- Lewandowska, M. & Więckowska, B. The influence of various smoking categories on the risk of gestational hypertension and pre-eclampsia. J. Clin. Med 9, 1743 (2020).
- Killen, J. D., Fortmann, S. P., Schatzberg, A., Hayward, C. & Varady, A. Onset of major depression during treatment for nicotine dependence. *Addictive Behav.* 28, 461–470 (2003).
- Jesse, S. et al. Alcohol withdrawal syndrome: mechanisms, manifestations, and management. Acta Neurol. Scand. 135, 4–16 (2017).
- Graham, H. & Owen, L. Are there socioeconomic differentials in under-reporting of smoking in pregnancy? *Tob. Control* 12, 434–434 (2003).
- Rukmani, M. R., Seshadri, S. P., Thennarasu, K., Raju, T. R. & Sathyaprabha, T. N. Heart rate variability in children with attention-deficit/hyperactivity disorder: a pilot study. *Ann. Neurosci.* 23, 81–88 (2016).
- Lucchini, M., Fifer, W. P., Sahni, R. & Signorini, M. G. Novel heart rate parameters for the assessment of autonomic nervous system function in premature infants. *Physiol. Meas.* 37, 1436–1446 (2016).

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the cooperation of the study participants, PASS investigators, the PASS Steering Committee Chairman Gary D.V. Hankins, MD, and members of the National Institute of Child Health and Human Development advisory safety monitoring board: Elizabeth Thom, Ph.D. (Chair); Reverend Phillip Cato, Ph.D.; James W. Collins Jr, MD, MPH; Terry Dwyer, MD, MPH; George Macones, MD; Philip A. May, Ph.D.; Jeff Murray, MD; Richard M. Pauli, MD, Ph.D.; Raymond W. Redline, MD; and Michael Varner, MD. Furthermore, the following individuals made significant contributions to the research and warrant recognition: DCAC: Idania Ramirez, MPH; Jamie Collins, MA; Laura Spurchise, MPH; DBPC: Richard A. Belliveau, BA; Kristin McMillan, BA; Megan Minter, MS; PAC: Johnston T. Grier, BA; Emilia F. Vignola, BA; Joseph J. Violaris, BA. All contributors were compensated by one or more of the listed funding sources.

The following institutions and researchers comprise the PASS Network (additional network members other than authors listed): Data Coordinating and Analysis Center (DCAC): Director: Kimberly Dukes, Ph.D.; Codirector: Lisa M. Sullivan, PhD; Biostatistics: Tara Tripp, MA; Project Management and Regulatory Affairs: Julie M. Petersen, BA; Rebecca A. Young, MPH; Data Management and Information Technology: Travis Baker, BS; Derek Petersen, BS; Gregory Toland, MS. Developmental Biology and Pathology Center (DBPC): Director: Hannah C. Kinney, MD; Assistant Director: Robin L. Haynes, Ph.D.; Coinvestigators: David S. Paterson, Ph.D.; Kevin G. Broadbelt, PhD; Kyriacos Markianos, Ph.D.; Ingrid A. Holm, MD; Theonia Boyd, MD: Drucilla Roberts, MD: Richard G. Goldstein, MD: Hanno Stein, Ph.D.: Technicians: Claire Maggiotto, BS; Catherine Hassett, BS. Comprehensive Clinical Site Northern Plains: Coinvestigators: Donald Habbe, MD; H. Eugene Hoyme, MD; William Massello III, MD: Bradley Randall, MD: Mary Ann Sens, MD, Ph.D.; Catherine Stoos, MD, Peter Van Eerden, MD; Project Management: Whitney Adler, BA; Elizabeth Berg, RN; Jessica Gromer, RN: Bethany Norton, MA: Liz Swenson, RN: Deb Tobacco, MA. Comprehensive Clinical Site South Africa: Project Management: Erna Carstens, RN; Mandy Potter, RN; Lucy Brink, MSc; Sylvia Abrahams, RN; Allison Borchardt, RN; Noorjahan Bosman, RN; Ciska Botha, BSocSc; Lehana Breytenbach, RN; Lucille Burger, RN; Carina Carstens, BCur (Nursing); Erna Carstens, RN; Weslin Charles, Hons BA (Psychology); Milly de Jager, RN: Carlie du Plessis, RN: Sandra Felix, RN: Nicola James (Nel), MCur (Nursing); Ronnie Jurgens, RN; Belda Klukowski, RN; Mugeleigh Lakay, MCur (Nursing); Samacia Maloy, RN; Marilyn Mentor, RN; Petra Opperman, RN; Berenice Pillay, RN; Mandy Potter, RN; Earl Prinsloo, BA Hons (Sociology); Sonja Schell, RN; Marisa van der Spuy, BCur (Nursing); Mariëtta Warwick, RN. Physiology Assessment Center (PAC): Project Management: Daianna Rodriguez, BA; Albany Perez, BA; Data Processing: Margaret C. Shair, BA; Tracy Thai, BA. National Institutes of Health: Project Scientists: Marian Willinger, PhD (National Institute of Child Health and Human Development); Dale Herald, Ph.D. (National Institute on Alcohol Abuse and Alcoholism); Howard J. Hoffman, Ph.D. (National Institute on Deafness and Other Communication Disorders); Chuan-Ming Li, MD, Ph.D. (National Institute on Deafness and Other Communication Disorders); Program Officers: Bill Dunty, Ph.D. (National Institute on Alcohol Abuse and Alcoholism); Tonse Raju, MD, DCH (National Institute of Child Health and Human Development); Gordon B. Hughes, MD (National Institute on Deafness and Other Communication Disorders).

AUTHOR CONTRIBUTIONS

A.S. conceptualized and conducted statistical analysis, drafted the initial manuscript, and reviewed and revised the manuscript. M.M.M. was a principal investigator of the study; he conceptualized and designed the study, supervised data collection and data processing, drafted the manuscript, reviewed, and revised it. A.J.E., W.P.F., and H.J.O. were principal investigators of the study, they acquired funding, contributed to data analyses and interpretation of the results, and writing of the manuscript. N.P. M.L., and L.C.S. contributed to the interpretation of results and drafting the manuscript. J.D.N., J.B., and J.A. participated in study supervision, data collection and processing, drafting and critical revision of the manuscript. S.R. participated in data analysis and drafting of the manuscript. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

FUNDING

This research was supported by grants UH3OD023279, U01HD055154, U01HD045935, U01HD055155, and U01AA016501, issued by the Office of the Director, National Institutes of Health of the National Institutes of Health, National Institute on Alcohol Abuse and Alcoholism, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, and the National Institute on Deafness and Other Communication Disorders. Ayesha Sania is supported by UH3OD023279-0551, re-entry supplement from Office of the Director, NIH, and Office of Research on Women Health (ORWH). The opinions expressed in this paper are those of the authors and do not necessarily represent the official views of the National Institutes of Health, the Eunice Kennedy Shriver National Institute on Alcohol Abuse and Alcoholism, or the National Institute on Deafness and Other Communication Disorders.

COMPETING INTERESTS

The authors declare no competing interests.

CONSENT STATEMENT

Written informed consent was obtained from all participants. Ethical approval was obtained from Stellenbosch University, Avera Health, Sanford Health, the Indian Health Service and from participating Tribal Nations.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at https://doi.org/10.1038/s41390-022-02060-5.

Correspondence and requests for materials should be addressed to Ayesha Sania.

Reprints and permission information is available at http://www.nature.com/ reprints

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.