CORRECTION Correction: Hormonal treatment for newly diagnosed metastatic prostate cancer: a population-based study from the California cancer registry

David J. Benjamin 💿, Anshu Shrestha, Dimitra Fellman, Rosemary D. Cress, Mark P. Lythgoe 💿 and Arash Rezazadeh Kalebasty

© The Author(s), under exclusive licence to Springer Nature Limited 2023

Prostate Cancer and Prostatic Diseases; https://doi.org/10.1038/s41391-023-00749-0

Correction to: *Prostate Cancer and Prostatic Diseases* https:// doi.org/10.1038/s41391-023-00732-9, published online 05 October 2023

The following sentence in the second paragraph of the Results section should read:

"Men who were 85 years or older were least likely to receive ADT (76.0%) as opposed to men between the ages of 20–54 (78.9%). Men between ages 75–84 had the highest proportion of receipt of ADT (81.6%)."

Instead of:

"Men between ages 75–84 were least likely to receive ADT (81.6%) as opposed to men between the ages of 20–54 (78.9%). Men between ages 75–84 had the highest proportion of receipt of ADT among all age groups."

There is also a typographical error in Table 2. Please find the correct table below:

| Characteristics | Adjusted OR* | Lower Cl | Upper Cl | p value |
|-----------------|-----------------|-------------|-------------|---------|
| Age (vs 20-54) | | | | |
| 55–64 | 0.831 | 0.693 | 0.996 | 0.0447 |
| 65–74 | 0.738 | 0.614 | 0.885 | 0.0011 |
| 75–84 | 0.820 | 0.671 | 1.002 | 0.0525 |
| 85+ | 0.558 | 0.442 | 0.705 | <0.0001 |

*The regression model included all the variables included in table 1 and year of diagnosis.

Additionally, the following sentence should read:

"Specifically, men who were 85 years or older had 44% lower odds of receiving ADT (95% CI: 0.44, 0.71) and men from the lowest nSES areas had 21% lower odds of receiving ADT as the initial treatment compared to men of ages 20–54 and those from the highest nSES neighborhood, respectively."

Instead of:

"Specifically, men who were 85 years or older had 44% lower odds of receiving ADT (95% CI: 0.44, 0.71) and men from the lowest nSES areas had 19% lower odds of receiving ADT as the initial treatment compared to men of ages 20–54 and those from the highest nSES neighborhood, respectively."