



Author Correction: Is body-weight supported treadmill training or robotic-assisted gait training superior to overground gait training and other forms of physiotherapy in people with spinal cord injury? A systematic review

J. Mehrholz^{1,2} · L. A. Harvey³ · S. Thomas¹ · B. Elsner²

Published online: 24 January 2018
© International Spinal Cord Society 2018

Correction to: Spinal Cord advance online publication 11 April 2017; <https://doi.org/10.1038/sc.2017.31>.

Since publication of this paper, the authors have become aware of two minor errors in the labelling of Figures 4 and 5. They are:

1. Figure 4: the title of column 2 is incorrect. It should read “Robotic” (not BWSTT).

2. Figure 5: the titles of columns 2 and 3 are the wrong way around. So column 2 should be “Robotic” (not “Overground”) and column 3 should read “Overground” (not “Robotic”).

The authors apologise for these errors and any inconvenience caused.

The original article can be found online at <https://doi.org/10.1038/sc.2017.31>.

✉ J. Mehrholz
jan.mehrholz@klinik-bavaria.de

¹ Institute of Science, Private European Medical Academy of the Klinik Bavaria Kreischa, Kreischa, Germany

² Department of Public Health, Dresden Medical School, Technical University Dresden, Dresden, Germany

³ John Walsh Centre for Rehabilitation Research, Kolling Institute, Sydney Medical School/Northern, University of Sydney, Sydney, New South Wales, Australia