CORRECTION **OPEN**

Correction: Importance of test–retest reliability for promoting fMRI based screening and interventions in major depressive disorder

Laurie Compère , Greg J. Siegle and Kymberly Young

© The Author(s) 2021

Translational Psychiatry (2021)11:429; <https://doi.org/10.1038/s41398-021-01544-y>

Correction to: *Translational Psychiatry* <https://doi.org/10.1038/s41398-021-01507-3>, published online 10 July 2021

The original version of this article unfortunately contained mistakes in Tables 1 and 4. The correct tables are given below. The original article has been corrected.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021

Table 1. Table of mean, standard deviation and median values of ICCs for each sample, reactivity model, and ROI.

Population	Reactivity model	Amygdala	DLPFC	rACC	sgACC liberally thresholded	sgACC conservatively thresholded
Controls & patients	Canonical amplitude	0.11 (± 0.09); 0.11	0.24 (± 0.16); 0.26	0.09 (± 0.10); 0.09	0.15 (± 0.08); 0.13	0.17 (± 0.09); 0.18
	Amplitude	0.23 (± 0.14); 0.22	0.12 (± 0.11); 0.12	0.11 (± 0.10); 0.12	-0.01 (± 0.13); -0.04	-0.04 (± 0.14); -0.08
	Area under the curve	0.13 (± 0.14); 0.12	0.08 (± 0.10); 0.07	0.03 (± 0.11); 0.03	-0.03 (± 0.09); -0.04	-0.06 (± 0.10); -0.07
	Onset delay	0 (± 0.09); -0.01	0.01 (± 0.09); 0	0 (± 0.10); 0	0 (± 0.08); 0.01	-0.01 (± 0.10); 0
	Rise decay	0 (± 0); 0	0 (± 0); 0	0 (± 0); 0	0 (± 0); 0	0 (± 0); 0
Patients	Height	0.08 (± 0.10); 0.09	0.21 (± 0.15); 0.23	0.13 (± 0.12); 0.14	0.16 (± 0.12); 0.17	0.18 (± 0.12); 0.23
	Canonical amplitude	0.09 (± 0.11); 0.11	0.22 (± 0.16); 0.23	0.08 (± 0.14); 0.08	0.10 (± 0.12); 0.07	0.14 (± 0.15); 0.12
	Amplitude	0.22 (± 0.15); 0.22	0.11 (± 0.13); 0.11	0.10 (± 0.13); 0.11	-0.06 (± 0.15); -0.07	-0.08 (± 0.14); -0.08
	Area under the curve	0.13 (± 0.14); 0.12	0.6 (± 0.12); 0.03	0.03 (± 0.13); 0.04	-0.08 (± 0.13); -0.08	-0.10 (± 0.13); -0.09
	Onset delay	-0.01 (± 0.12); -0.01	0.01 (± 0.12); 0	-0.01 (± 0.13); -0.01	0.02 (± 0.11); 0.02	0.01 (± 0.12); 0.05
	Rise decay	0 (± 0); 0	0 (± 0); 0	0 (± 0); 0	0 (± 0); 0	0 (± 0); 0
	Height	0.09 (± 0.12); 0.08	0.22 (± 0.16); 0.23	0.12 (± 0.15); 0.13	0.16 (± 0.17); 0.18	0.17 (± 0.17); 0.21

Mean ($\pm$ standard deviation); median.**Table 4.** Table of mean, standard deviation and median values of ICCs for each sample, preprocessing, and first level parameter with cluster correction applied.

Preprocessing	TBV style	Standard
Without training - Control - Baseline	Canonical amplitude	-0.07 (± 0.21); -0.09
	Amplitude	0.29 (± 0.2); 0.3
	Area under the curve	0.02 (± 0.21); 0.01
	Onset-delay	-0.03 (± 0.23); -0.05
	Rise-decay	NA ($\pm$ NA); NA
With training - Experimental - Transfer	Height	0.36 (± 0.23); 0.33
	Canonical amplitude	-0.11 (± 0.21); -0.12
	Amplitude	0.3 (± 0.18); 0.31
	Area under the curve	0.06 (± 0.20); 0.07
	Onset-delay	0.02 (± 0.24); -0.02
	Rise-decay	NA ($\pm$ NA); NA
	Height	0.52 (± 0.19); 0.56

Mean ($\pm$ standard deviation); median.