




## ARTICLE OPEN



# Pramipexole restores behavioral inhibition in highly impulsive rats through a paradoxical modulation of frontostriatal networks

Robin Magnard<sup>1</sup> <sup>✉</sup>, Maxime Fouyssac<sup>2</sup>, Yvan M. Vachez<sup>1</sup>, Yifeng Cheng<sup>3</sup>, Thibault Dufourd<sup>1</sup>, Carole Carcenac<sup>1</sup>, Sabrina Boulet<sup>1</sup>, Patricia H. Janak<sup>3,4</sup>, Marc Savasta<sup>1</sup>, David Belin<sup>1,5</sup>  and Sebastien Carnicella<sup>1,5</sup> 

© The Author(s) 2024

Impulse control disorders (ICDs), a wide spectrum of maladaptive behaviors which includes pathological gambling, hypersexuality and compulsive buying, have been recently suggested to be triggered or aggravated by treatments with dopamine  $D_{2/3}$  receptor agonists, such as pramipexole (PPX). Despite evidence showing that impulsivity is associated with functional alterations in corticostriatal networks, the neural basis of the exacerbation of impulsivity by PPX has not been elucidated. Here we used a hotspot analysis to assess the functional recruitment of several corticostriatal structures by PPX in male rats identified as highly (HI), moderately impulsive (MI) or with low levels of impulsivity (LI) in the 5-choice serial reaction time task (5-CSRTT). PPX dramatically reduced impulsivity in HI rats. Assessment of the expression pattern of the two immediate early genes *C-fos* and *Zif268* by in situ hybridization subsequently revealed that PPX resulted in a decrease in *Zif268* mRNA levels in different striatal regions of both LI and HI rats accompanied by a high impulsivity specific reduction of *Zif268* mRNA levels in prelimbic and cingulate cortices. PPX also decreased *C-fos* mRNA levels in all striatal regions of LI rats, but only in the dorsolateral striatum and nucleus accumbens core (NAc Core) of HI rats. Structural equation modeling further suggested that the anti-impulsive effect of PPX was mainly attributable to the specific downregulation of *Zif268* mRNA in the NAc Core. Altogether, our results show that PPX restores impulse control in highly impulsive rats by modulation of limbic frontostriatal circuits.

*Translational Psychiatry* (2024)14:86; <https://doi.org/10.1038/s41398-024-02804-3>

## INTRODUCTION

Impulse control disorders (ICDs) represent a set of heterogeneous maladaptive behaviors characterized by a deficit in urge regulation and inhibition that are core symptoms of several neuropsychiatric conditions, that include, for instance, intermittent explosive disorder, pathological gambling and hypersexuality [1].

Despite the profound consequences ICDs have on the quality of life of patients, their neurobiological basis has not yet been elucidated. However ICDs show a relatively high prevalence in Parkinson's disease (PD) [2, 3], restless leg syndrome [4–6], fibromyalgia [7] and hyperprolactinemia [8–10], the treatment of which involves dopamine  $D_{2/3}$  receptors agonists such as pramipexole (PPX) or ropinirole. This implicates the iatrogenic effect of dopaminergic drugs and a potential abnormal dopaminergic function in the development of ICDs.

Alongside their apparent reliance on aberrant dopaminergic mechanisms, the compulsive nature of ICDs, which is not without similarities with that of substance use disorders [11–18], has led to their conceptualization as a form of behavioral addiction [19–21]. Similarly to psychostimulant use disorder, the vulnerability to develop ICDs has been shown to be associated with a high

impulsivity trait [14, 17, 21–24], which is characterized by a tendency to act prematurely, without forethought or concerns for adverse upcoming consequences [25]. Impulsivity is a multifaceted construct [26] that encompasses the inability to tolerate delays to reinforcement and adapt to risks, referred to as cognitive impulsivity, on one hand, and the inability to withhold prepotent responses, resulting in poor or adverse outcomes, so-called motor/waiting impulsivity [25], on the other hand. Cognitive impulsivity is principally assessed in delayed discounting [27–29] or risk taking [30–32] tasks whereas waiting impulsivity is canonically assessed as the rate of premature responses in the 5-choice serial reaction time task (5-CSRTT) [33, 34].

The respective contribution of each of these dimensions of impulsivity, which are behaviorally and neurally dissociable [17, 35–37], to the emergence of ICDs following exposure to dopaminergic drugs such as PPX has not yet been elucidated. While preclinical studies in rodents and non-human primates have demonstrated that PPX exacerbates cognitive impulsivity, as assessed in delay discounting [27–29] and risk taking [30–32] tasks, the influence of PPX on waiting impulsivity is less clear. Indeed, PPX has been shown to exacerbate waiting impulsivity as

<sup>1</sup>Univ. Grenoble Alpes, Inserm, U1216, Grenoble Institut Neurosciences, 38000 Grenoble, France. <sup>2</sup>Department of Psychology, University of Cambridge, Downing Street, CB2 3EB Cambridge, United Kingdom. <sup>3</sup>Department of Psychological and Brain Sciences, Johns Hopkins University, Baltimore, MD 21218, USA. <sup>4</sup>Solomon H. Snyder Department of Neuroscience, Johns Hopkins School of Medicine, Johns Hopkins University, Baltimore, MD 21205, USA. <sup>5</sup>These authors jointly supervised this work: David Belin, Sebastien Carnicella. ✉email: robin.magnard@jhu.edu

Received: 5 August 2023 Revised: 18 January 2024 Accepted: 23 January 2024

Published online: 09 February 2024

assessed in differential reinforcement of low rate of responding and fixed consecutive number tasks [38]. In the 5-CSRTT, one study [39] found PPX to have a detrimental effect mainly on accuracy, the index of attention, and to bring omissions rate up to almost 50%, suggesting that the animals were no longer engaged in the task in that study, precluding any assessment of the effect of the drug on waiting impulsivity per se. In addition, the interaction between PPX treatment and impulsivity trait was not investigated.

Considering the heuristic and translational value of high impulsivity trait, characteristic of individuals belonging to the upper quartile of the population ranked on the rate of premature responses in the 5-CSRTT, with regards to the vulnerability to develop compulsive behaviors [40–43], in the present study we sought to test the hypothesis that PPX may exert an impulsivity trait specific effect on impulse control associated with functional alterations in underlying corticostriatal circuits [25]. A pro-impulsive action of PPX in highly impulsive (HI) rats characterized in the 5-CSRTT would provide experimental evidence towards a facilitation of the development of compulsivity by dopaminergic drugs in this vulnerable population.

At the neural systems level, impulsivity has been shown to depend on the functional engagement of a distributed corticostriatal network [25] that involves the insular cortex, the nucleus accumbens core (NAc Core) and shell (NAc Shell) and the dorsomedial striatum (DMS) [25, 37, 44, 45]. In contrast, compulsive behaviors that result from the interaction between a pre-existing impulsive trait and dopaminergic drugs have repeatedly been shown to be associated with, if not mediated by, dopaminergic mechanisms in the dorsolateral striatum (DLS) [46, 47]. Thus, using *in situ* hybridization we quantified the mRNA levels of the cellular activity and plasticity markers *C-fos* [48–50] and *Zif268* [51–55], respectively, in corticostriatal structures and used a mediation analysis that aimed to identify the functional signature of the effect of PPX on impulsivity in vulnerable individuals.

## MATERIALS AND METHODS

### Subjects

Experiments were performed on 48 male Sprague Dawley rats (Janvier, France) that were 6 weeks old (weighing 200 g) when they were habituated to the animal facility in which they were housed 1 per cage under a 12 h/light/dark cycle (lights ON at 7 am). Rats were food restricted to 90% of their theoretical free feeding weight for 5-CSRTT training, but had *ad libitum* access to water throughout the experiment. Protocols complied with the European Union 2010 Animal Welfare Act and the new

French directive 2010/63 and were approved by the French national ethics committee n°004.


### 5-CSRTT training

The 5-CSRTT procedure has been adapted for Sprague-Dawley from previous studies [33, 56–59]. Briefly, the apparatus consisted of eight 25 × 25 × 25 cm operant chambers (Med associates, St Albans, VT) each equipped with a curved rear wall. Set into the curved walls were five 2.5 × 2.5 cm square holes, 4 cm deep and 2 cm above the floor. Each hole was equipped with an infra-red beam crossing the entrance horizontally and a cue light at its rear that provided illumination. A houselight was located at the top of this wall. 45 mg sucrose pellets (TestDiet, VA) were delivered from a pellet dispenser to a tray at the front of the cage, also equipped with a cue light and infra-red beam.

**Phase 1: Learning.** The procedure began one week after food restriction was initiated. To prevent neophobia, rats were exposed to 10 sucrose pellets in their home cage for two consecutive days. The following two days, rats were placed in their operant box for 15 min habituation sessions, during which ten pellets were placed in the magazine and two pellets were placed in each hole, all cue- and house-lights were turned ON.

**Phase 2: Task acquisition.** As detailed in Fig. 1A each daily session consisted of 100 discrete trials with stable performance being achieved after about 40 sessions. In order to facilitate brief stimulus detection by Sprague-Dawley rats, the house-light remained OFF during the length of the session and was turned on during time out periods [60]. Rats were trained to enter the food magazine to initiate a trial. After a 5 s intertrial interval (ITI), a brief light stimulus was pseudo-randomly presented in one of five holes to indicate the individual the location of the correct response for that trial. Following a nosepoke in this hole ('correct response'), rats were rewarded with the delivery of one sucrose pellet in the food tray. A nosepoke response in any of the adjacent holes ('incorrect response'), or a failure to respond within 5 s after the onset of the stimulus ('omission'), resulted in no pellet delivery and a 5 s time-out period signaled by the house light being turned ON. Additional nosepokes performed after a correct or incorrect response (referred as 'perseverative response') were recorded but had no consequence. Nosepokes made during the ITI, that is, before the onset of the stimulus (or 'premature responses') were recorded as a measure of waiting impulsivity, and resulted in a 5 s time-out (house light turned ON) and reward omission. Nosepokes made during this time out period ('time-out response') were recorded, but had no programmed consequence. The stimulus duration was progressively reduced from 20 s to 1 s across training stages, as previously described [58, 59]. Progress across the different stages was determined by at least 80% accuracy and less than 20% omissions on any particular stage. Rats which failed to fulfill these criteria were excluded from the study ( $n = 4$ ).

**Phase 3: Challenge with long intertrial interval.** Following acquisition of the task, rats were challenged with three 7 s-long intertrial interval (LITI)



**Fig. 1 Structure of the 5-CSRTT and identification of low, moderate and high impulsivity rats. A** 5-CSRTT trial structure. **B** Segregation of low impulsive (LI,  $n = 11$ ), moderately impulsive (MI,  $n = 22$ ) and highly impulsive (HI,  $n = 11$ ) rats. BL baseline, LITI long intertrial interval. Data are shown as means  $\pm$  SEM. HI vs. LI  $^{\circ\circ\circ}p < 0.001$ ; HI vs. MI  $^{\wedge\wedge}p < 0.001$ ; MI vs. LI  $^{++}p < 0.01$ ;  $^{+++}p < 0.001$ .

sessions separated by two baseline sessions with the regular 5 s ITI. Subjects were ranked according to their level of premature responses during second and third LITI [45] and those in the upper and lower quartile ( $n = 11$  each) [45] were considered as having high (HI) or low impulsivity (LI), respectively. The rest of the population was considered to be moderately impulsive (MI,  $n = 22$ ).

### Pharmacological procedures

Individuals of the HI, MI and LI groups were randomly allocated to a PPX or a vehicle treatment group so that the PPX-treated HI and LI rats ( $n = 6$  and  $5$  respectively), as well as MI rats ( $n = 12$ ) showed a similar level of premature responses before initiation of treatment as that shown by Veh-treated HI, LI and MI rats ( $n = 6$ ,  $n = 5$  and  $n = 10$ , respectively). The experimenter was not blinded to group allocation to facilitate treatment administration and ensure equitable distribution of operant cages. Rats received an intraperitoneal administration of 0.2 mg/kg/day PPX (Sigma–Aldrich, St. Louis, MO), diluted in 0.9% NaCl, or vehicle (Veh) 30 min prior to each daily behavioral session, with treatment starting one day before to the first of twelve sessions (11 baseline 5 s ITI sessions and one final 7 s LITI session) over which the effect PPX on impulsivity in HI and LI rats was measured. The dose of pramipexole used here was carefully chosen because it exacerbates impulsivity in a delay discounting task (Magnard & Carnicella, unpublished) and improve motivational function in dopamine-deficient rats [61] (see for review [62]), while it does not influence locomotion [63, 64].

### Tissue collection

15 min after the final LITI session, rats were deeply anesthetized by isoflurane and killed by decapitation. Brains were quickly removed and snap frozen in liquid nitrogen, then stored at  $-80^{\circ}\text{C}$  until they were processed into 14  $\mu\text{m}$ -thick coronal brain sections with a cryostat (Microm HM 500, Microm, Francheville, France) and collected on permafrost gelatin-coated slides (Colorfrost Plus, Fisher scientific, Pittsburg, PA) that were stored at  $-80^{\circ}\text{C}$ .

### In situ hybridization

The in situ hybridization (ISH) procedure was carried out as previously described [65, 66] with oligonucleotide probes specifically complementary to the sequence of the mRNA of the immediate early genes C-Fos (nucleotides 159-203 of the NCBI Reference Sequence NM\_022197.2) or Zif268 (nucleotides 1680-1724 of the NCBI Reference Sequence NM\_012551.3), tailed by 3'OH incorporation of  $^{35}\text{S}$ -dATP (1250 mCi/mmol, Perkin Elmer, UK) by a terminal Deoxynucleotidyl transferase (Promega, M1875) with a specificity of  $2.5 \times 10^6$  cpm/ml (C-Fos) or  $3 \times 10^6$  cpm/ml (Zif268).

Following fixation and pre-hybridization treatments aiming at reducing non-specific hybridization, slides were incubated overnight at  $42^{\circ}\text{C}$  in the hybridization buffer (50% deionized formamide, 10% dextran sulfate, 50 ng/ml denaturated salmon sperm DNA, 5% Sarcosyl, 0.2% SDS, 1 mM EDTA, 300 nM NaCl, 5X Denhardt's in 2X standard sodium citrate (SSC)) with the probes diluted at a concentration of 8.75 ng/ml (C-Fos) or 6.25 ng/ml (Zif268). Slides were then washed in decreasing concentration of SSC and dehydrated in increased ethanol concentration baths. Sections were exposed to Biomax MR films (Kodak, Rochester, USA) for four weeks (Zif268-labeled sections) or 6 weeks (C-Fos-labeled sections) at room temperature. Films were revealed in a dark room. Pictures of each brain section were taken on a Northern light (Imaging Res Inc.) light table with a Qicam (QImaging) camera equipped with a SIGMA 50 mm 1:2.8 DG MacroD Fast 1394 (Nikon) objective and subsequently analyzed with ImageJ software [67]. A region of interest was drawn for each striatal territory in which the optical density reflective of the mRNA level was measured (according to the rat brain atlas [68]). As shown in Supplementary Fig. 1A, B, the optical density in an mRNA-free part of the brain (i.e., corpus callosum) was defined as background, and this value was subtracted to that obtained from the area of interest to compute the relative optical density used as the dependent variable in subsequent analyses.

### Data and statistical analyses

Data, presented as mean  $\pm$  SEM with or without superimposed individual data points, were analyzed using SigmaStat (Systat software Inc., San Jose, USA) and SPSS (IBM, Amarak, NY). Sample size was estimated based on previous studies using the same procedures [42, 45]. Three-way repeated measure analyses of variance (RM ANOVAs) were used to compare groups

across sessions, with sessions as within-subject factor, impulsivity (LI, MI or HI) and treatment (Veh or PPX) as between-subject factors. Two-way RM ANOVAs were used to compare levels of impulsivity between LI, MI and HI rats during the screening period, with sessions as within-subject factor and impulsivity as between-subject factor. Two-way ANOVAs were also used to compare LI, MI and HI rats for BL11 and LITI sessions with impulsivity and treatment as between-factors. Two-way ANCOVA was used to control for the omission and BL11 premature response performances on premature responses during LITI, with impulsivity and treatment as the between-subject variables, LITI omissions and BL11 premature responses as covariants.

Differences between LI, MI and HI rats in Zif268 and C-fos mRNA were analyzed using three-way ANOVA with impulsivity and treatment as between-subject factors and structures as within-subject factor. When indicated, *post hoc* analyses were carried-out using the Student-Newman-Keuls test.

Dimensional inter-relationships were analyzed with nonparametric Spearman correlation coefficient  $\rho$  and Spearman rank order tests. Assumptions for the normality of the distributions and the homogeneity of variance were verified using the Shapiro–Wilk and Levene test, respectively. Significant violations of homogeneity of variances and normality were corrected using square root transformations. Significance for  $p$  values was set at  $\alpha = 0.05$ . Effect sizes for the ANOVAs are also reported using partial  $\eta^2$  values ( $\eta_p^2$ ) [69, 70].

In order better to delineate the neural basis of the influence of PPX on impulsivity, we employed mediation analysis [71, 72] that helped decipher the direct effect of treatment and the indirect effect mediated by C-fos and Zif268 expression on impulsivity. For this, we applied:

- a. dependent regression (direct effect)

$$\text{Impulsivity} = \sum_{i=1}^7 \beta_{\text{fos}}^i \text{Fos}(i) + \sum_{i=1}^7 \beta_{\text{zif}}^i \text{Zif}(i) + \beta_T T + \beta_L L$$

- b. mediator regression

$$\text{Fos}(i) = \alpha_{\text{fos}}^i * T$$

$$\text{Zif}(i) = \alpha_{\text{zif}}^i * T$$

- c. effect decomposition (indirect effect)

$$Y_{\text{fos}}^i = \beta_{\text{fos}}^i * \alpha_{\text{fos}}^i$$

$$Y_{\text{zif}}^i = \beta_{\text{zif}}^i * \alpha_{\text{zif}}^i$$

where  $\text{Fos}(i)$  and  $\text{Zif}(i)$  represent the mRNA levels of the immediate early genes (IEG) C-fos and Zif268, respectively, from each brain region.  $i$  represents the arbitrary index of the seven brain regions we stained (IL, PrL, Cg, DLS, DMS, NAc Core, NAc Shell).  $T$  is a binary variable (1 for PPX, 0 for Veh).  $L$  is impulsivity measured under LITI prior to pharmacological treatment.  $\beta$  represents the direct effect from each variable.  $\alpha$  represents the mediation effect from treatment (T) to cellular activity/plasticity.  $\gamma$  represents the indirect effect of treatment on impulsivity mediated by the two IEGs. To strengthen the analysis and prevent any bias by analyzing only a subpopulation of this study, we included LI, MI and HI -Veh and PPX-treated animals to this analysis which was carried out using the JASP statistical software with bootstrap method, 1000 replications and bias-corrected percentile [71, 72].

## RESULTS

### 5-CSRTT screening for low impulsive and high impulsive rats

As previously described [37, 45] when Sprague Dawley rats well-trained in a 5-CSRTT are challenged with longer ITIs, thereby requiring individuals to refrain from expressing prepotent responses for a slightly longer period of time than they usually do (Fig. 1A), marked individual differences in waiting impulsivity are revealed that allow identification of HI and LI rats in the upper and lower quartiles of the population, respectively, and MI in the two middle quartiles. In accordance, HI rats displayed a greater

increase in premature responses than LI rats during the LITI sessions (Fig. 1B) [impulsivity  $\times$  session interaction:  $F_{22,451} = 11.74$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.36$ ].

**PPX influences impulse control differentially in LI and HI rats**  
LI, MI and HI rats were then split into three treatment groups that did not differ in their baseline impulsivity level [no impulsivity  $\times$  pretreatment  $\times$  session interaction:  $F_{22,418} < 1$ ,  $p = 0.914$ ,  $\eta_p^2 = 0.03$ ]. Treatment groups received daily IP injections of either PPX (LI, MI or HI PPX-treated rats) or vehicle (LI, MI or HI Veh-treated rats) for 12 sessions.

PPX treatment was revealed to influence differentially baseline impulsivity and its exacerbation by LITI mostly in HI rats (Fig. 2A) [sessions  $\times$  impulsivity  $\times$  treatment interaction:  $F_{22,418} = 2.72$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.125$ ]. While HI-Veh rats did not differ from LI-Veh rats during baseline sessions, PPX treatment resulted in a progressive increase in baseline premature responses only in HI rats (Fig. 2A, B, BL white panels). In marked contrast, PPX treatment prevented the exacerbation of premature responses otherwise shown by HI-Veh and MI-Veh rats upon introduction of LITI (Fig. 2A, B, LITI gray panels) [impulsivity  $\times$  treatment interaction:  $F_{2,44} = 3.75$ ,  $p = 0.03$ ,  $\eta_p^2 = 0.16$ ].

This effect was not due to a difference in baseline levels of impulsivity as it persisted when the premature responses expressed during the baseline session that preceded the LITI (BL11) was used as a covariate in an ANCOVA [impulsivity  $\times$  treatment interaction:  $F_{2,44} = 6.4$ ,  $p = 0.004$ ,  $\eta_p^2 = 0.257$ ]. Within-subject analyses across LITIs confirmed that PPX decreased the level of premature responding under LITI as compared to pre-treatment performance in HI and to a lesser extent in MI rats (Fig. 2C) [period  $\times$  impulsivity  $\times$  treatment interaction:  $F_{2,38} = 4.11$ ,  $p = 0.024$ ,  $\eta_p^2 = 0.178$ ]. Thus, while PPX treatment moderately increased impulsivity in HI rats in baseline conditions, it abolished its exacerbation during LITI sessions, when inhibitory control is most challenged.

#### The effects of PPX were specific to impulsivity

PPX treated rats showed a progressive decline in accuracy over the course of treatment (Fig. 2D) [treatment:  $F_{1,38} = 10.27$ ,  $p = 0.003$ ,  $\eta_p^2 = 0.21$ ], which may reflect an impairment in attention [34]. However, this effect was not dependent on impulsive trait [no sessions  $\times$  impulsivity interaction  $F_{22,418} < 1$ ,  $p = 0.57$ ,  $\eta_p^2 = 0.05$ ; no sessions  $\times$  impulsivity  $\times$  treatment interaction  $F_{22,418} = 1.02$ ,  $p = 0.44$ ,  $\eta_p^2 = 0.05$ ]. This result was however not supported by the analysis of performance during the LITI session (Fig. 2E) [treatment:  $F_{2,44} = 1.9$ ,  $p = 0.17$ ,  $\eta_p^2 = 0.05$ ; no treatment  $\times$  impulsivity interaction:  $F_{2,44} = 0.27$ ,  $p = 0.76$ ,  $\eta_p^2 = 0.014$ ]. Together these results demonstrate that under the present experimental conditions, a moderate daily dose of PPX, had a restricted effect on attention, that was not dependent on impulsive phenotype.

In addition, we assessed response omissions and collection latencies, which jointly not only provide an insight into task engagement [34] but could also potentially influence the rate of premature responses. PPX increased the percentage of omitted trials (Fig. 2F) [sessions  $\times$  treatment interaction  $F_{11,418} = 2.5$ ,  $p = 0.005$ ,  $\eta_p^2 = 0.06$ ; no sessions  $\times$  impulsivity  $\times$  treatment interaction  $F_{22,418} = 0.67$ ,  $p = 0.86$ ,  $\eta_p^2 = 0.03$ ]. Because this effect was exacerbated in MI and HI rats irrespective of ITI duration [impulsivity  $\times$  treatment interaction  $F_{2,38} = 3.26$ ,  $p = 0.049$ ,  $\eta_p^2 = 0.147$ ], even though only an effect of the treatment was observed during the LITI session (Fig. 2G) [treatment:  $F_{1,44} = 10$ ,  $p = 0.003$ ,  $\eta_p^2 = 0.20$ ; no impulsivity  $\times$  treatment interaction  $F_{2,44} < 1$ ,  $p = 0.43$ ,  $\eta_p^2 = 0.043$ ] an ANCOVA was carried out to determine a potential influence of this PPX-induced increase in omissions on impulsivity using omissions expressed during the LITI session as a covariate. This analysis revealed that the effect of PPX on omissions during the LITI session cannot account for its profound

effect on premature responses [impulsivity  $\times$  treatment interaction:  $F_{2,44} = 4.15$ ,  $p = 0.024$ ,  $\eta_p^2 = 0.187$ ]. Whereas PPX also increased the latencies to correct responses irrespective of the impulsive phenotype, (Fig. 2H) [treatment  $F_{1,38} = 45.5$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.54$ ; no sessions  $\times$  impulsivity  $\times$  treatment interaction:  $F_{22,418} = 1.37$ ,  $p = 0.18$ ,  $\eta_p^2 = 0.06$ ; no impulsivity  $\times$  treatment interaction  $F_{2,38} = 2.14$ ,  $p = 0.13$ ,  $\eta_p^2 = 0.10$ ] and (Fig. 2I) [no impulsivity  $\times$  treatment interaction:  $F_{2,44} = 1.78$ ,  $p = 0.18$ ,  $\eta_p^2 = 0.08$ ; treatment  $F_{1,44} = 15.42$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.28$ ], it did not influence reward collection latencies (Fig. 2J) [no sessions  $\times$  impulsivity  $\times$  treatment interaction,  $F_{22,418} < 1$ ,  $p = 0.85$ ,  $\eta_p^2 = 0.035$ ; treatment  $F_{1,38} < 1$ ,  $p = 0.42$ ,  $\eta_p^2 = 0.017$ ] and (Fig. 2K) [treatment:  $F_{1,44} = 1.7$ ,  $p = 0.19$ ,  $\eta_p^2 = 0.04$ ; no impulsivity  $\times$  treatment interaction:  $F_{2,44} < 1$ ,  $p = 0.79$ ,  $\eta_p^2 = 0.01$ ].

Thus, the slight decrease in response readiness caused by PPX especially in HI rats, is unlikely to account for the effect of this dopaminergic drug on impulsivity. This was further supported by the absence of correlation between premature responses performed during the final LITI session, under differential treatment, and the response rate before treatment during either baseline (Supplementary Fig. 2A) [ $\rho = -0.133$ ,  $p = 0.391$ ] (see also Supplementary Fig. 2B–D for LI, MI and HI rats, respectively) or LITI sessions (Supplementary Fig. 2E) [ $\rho = 0.184$ ,  $p = 0.233$ ] (see also Supplementary Fig. 2F–H for LI, MI and HI rats, respectively), thus confirming that the effect of PPX on response readiness did not influence premature responses.

Finally, PPX did not influence perseverative responses (Supplementary Fig. 3A) [no sessions  $\times$  impulsivity  $\times$  treatment interaction,  $F_{22,418} < 1$ ,  $p = 0.92$ ,  $\eta_p^2 = 0.03$ ; no impulsivity  $\times$  treatment interaction  $F_{3,38} < 1$ ,  $p = 0.63$ ,  $\eta_p^2 = 0.02$ ], but reduced magazine entries reminiscent of reduced response readiness noticed in PPX treated rats (Supplementary Fig. 3B) [session  $\times$  treatment interaction:  $F_{11,418} = 2.9$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.07$ ; no sessions  $\times$  impulsivity  $\times$  treatment interaction,  $F_{22,418} < 1$ ,  $p = 0.82$ ,  $\eta_p^2 = 0.03$ ; no impulsivity  $\times$  treatment interaction:  $F_{2,38} = 1.3$ ,  $p = 0.26$ ,  $\eta_p^2 = 0.06$ ]. Therefore, PPX treatment did not induce repetitive or stereotyped behaviors that may influence performance in the 5-CSRTT.

#### The dampening effect of PPX on the exacerbated impulsivity during LITI sessions in HI rats was associated with a functional disengagement of corticostriatal circuits


In order to identify the neural basis of the effect of PPX on impulsivity trait, we carried out a hot-spot analysis in structures of the corticostriatal circuitry using in situ hybridization for the markers of cellular activity and plasticity C-fos and Zif268, respectively.

A first quality control analysis of the pattern of expression of the two IEGs, based on a covariance analysis for each marker, revealed a convergence of RNA levels in line with the functional organization of the corticostriatal circuitry, with similar trends in expression in structures inter-connected to each other (Supplementary Fig. 4). However, as anticipated, the two IEGs were shown here not to reflect the same functional process as their mRNA levels did not covary (Supplementary Fig. 4).

An impulsive trait dependent reduction of Zif268 was observed in the dorsal structures of the mPFC, namely the prelimbic and cingulate cortexes of PPX treated rats (Fig. 3A) [structure  $\times$  impulsivity  $\times$  treatment interaction:  $F_{4,54} = 3.18$ ,  $p = 0.02$ ,  $\eta_p^2 = 0.19$ ].

This decreased recruitment of a marker of cellular plasticity in the prefrontal cortical areas was accompanied by a decrease in the level of expression of Zif268 across the ventral and dorsal domains of the striatum in PPX-treated rats (Fig. 3B) [structure  $\times$  treatment interaction:  $F_{3,84} = 11.12$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.28$ ; no structure  $\times$  impulsivity  $\times$  treatment interaction:  $F_{6,84} = 0.53$ ,  $p = 0.78$ ,  $\eta_p^2 = 0.037$ ]. While the decrease in Zif268 mRNA levels in the DLS was quantitatively similar to that observed in the DMS, differences were observed between the NAc Core and NAc Shell in that Zif268 mRNA levels were much more reduced by PPX in the Core than the Shell.






**Fig. 2 PPX influences premature responses differentially in LI, MI and HI rats in baseline and LITI conditions.** **A** Percentage of premature responses across baseline (BL) and long intertrial interval (LITI) sessions. **B** Differential effect of PPX on premature responses between the last BL and the LITI session. **C** Comparison of the level of premature responding during LITI before and during differential PPX vs. Veh treatment. **D** Response accuracy across BL and LITI sessions. **E** Response accuracy during the LITI session under differential PPX vs. Veh treatment. **F** Response omissions across BL and LITI sessions. **G** Response omission during the LITI session under differential PPX vs. Veh treatment. **H** Correct response latencies across BL and LITI sessions. **I** Correct response latencies during the LITI session under differential PPX vs. Veh treatment. **J** Reward collection latencies across BL and LITI sessions. **K** Reward collection latencies during the LITI session under differential PPX vs. Veh treatment. LI-Veh  $n = 5$ ; LI-PPX  $n = 6$ ; MI-Veh  $n = 10$ ; MI-PPX  $n = 12$ ; HI-Veh  $n = 6$ ; HI-PPX  $n = 5$ . LI low impulsive, MI Moderately impulsive, HI highly impulsive. Data are shown as means  $\pm$  SEM. HI-Veh vs. LI-Veh  $^{\circ}p < 0.05$ ;  $^{\circ\circ}p < 0.01$ . HI-Veh vs. MI-Veh  $^{\wedge}p < 0.05$ ;  $^{\wedge\wedge}p < 0.01$ . MI-Veh vs. LI-Veh  $^{+}p < 0.05$ . LI-Veh vs. LI-PPX  $^{\S}p < 0.05$ ;  $^{\S\S\S}p < 0.001$ . MI-Veh vs. MI-PPX  $^{\text{f}}p < 0.05$ ;  $^{\text{ff}}p < 0.01$ ;  $^{\text{fff}}p < 0.001$ . HI-Veh vs. HI-PPX  $^{*}p < 0.05$ ;  $^{**}p < 0.01$ ;  $^{***}p < 0.001$ . HI-PPX vs. MI-PPX  $^{\%}p < 0.05$ ;  $^{\%\%}p < 0.01$ . HI-PPX vs. LI-PPX  $^{\#}p < 0.05$ ;  $^{\#\#}p < 0.01$ ;  $^{\#\#\#}p < 0.001$ . LI-PPX vs. MI-PPX  $^{\text{¶}}p < 0.05$ ;  $^{\text{¶¶}}p < 0.01$ . LI-PPX BL11 vs. LI-PPX LITI  $^{\&}p < 0.05$ . HI-Veh BL11 vs. HI-Veh LITI  $^{\text{sss}}p < 0.001$ . MI-Veh BL11 vs. MI-Veh LITI  $^{\text{yy}}p < 0.01$ . MI-PPX BL11 vs. MI-PPX LITI  $^{\text{xx}}p < 0.01$ . HI-PPX prior treatment vs. HI-PPX under treatment  $^{\text{cc}}p < 0.01$ . MI-PPX prior treatment vs. MI-PPX under treatment  $^{\text{b}}p < 0.05$ .

The effect of PPX on the levels of C-fos mRNA in the prefrontal regions of LI, MI and HI rats followed the same trend as that observed for Zif268, albeit of a lesser magnitude (Fig. 3C) [structure  $\times$  impulsivity  $\times$  treatment interaction:  $F_{4,54} = 2.77$ ,  $p = 0.036$ ,  $\eta_p^2 = 0.17$ ]. However, structure level analysis failed to

isolate significant differences between the different treatment or impulsivity conditions.


In the striatum, PPX treatment resulted in a global decrease in C-fos mRNA levels in the DLS, DMS, NAc Core, but no difference was observed in the NAc Shell (Fig. 3D) [structure  $\times$  treatment interaction:



**Fig. 3 Zif268- and C-fos-based functional signature of the effect of PPX on impulsivity in the prefrontal corticostriatal circuitry of LI, MI and HI rats.** **A** Zif268 mRNA levels quantified in cortical areas. Upper panel: representative high-resolution photos of the signal obtained following in situ hybridization with a probe against Zif268 in Veh- and PPX-treated LI, MI and HI rats. Lower panel: Optical densities measured in the IL, the PrL and the Cg cortex. **B** Zif268 mRNA levels quantified in the striatum. Upper panel: representative high-resolution photos of the signal obtained following in situ hybridization with a probe against Zif268 in Veh- and PPX-treated LI, MI and HI rats. Lower panel: Optical density measured in the DLS, the DMS, the NAc Core and the NAc Shell. **C** C-fos mRNA levels quantified in cortical areas. Upper panel: representative high-resolution photos of the signal obtained following in situ hybridization with a probe against C-fos in Veh- and PPX-treated LI, MI and HI rats. Lower panel: Optical densities measured in the IL, the PrL and the Cg cortex. **D** C-fos mRNA levels quantified in the striatum. Upper panel: representative high-resolution photos of the signal obtained following in situ hybridization with a probe against C-fos in Veh- and PPX-treated LI, MI and HI rats. Lower panel: Optical density measured in the DLS, the DMS, the NAc Core and the NAc Shell. Photos per different condition represent the same animal stained with different in situ hybridization probe. LI-Veh  $n = 4-5$ ; LI-PPX  $n = 5-6$ ; MI-Veh  $n = 4-8$ ; MI-PPX  $n = 7-9$ ; HI-Veh  $n = 5-6$ ; HI-PPX  $n = 5$ . LI low impulsive, MI moderately impulsive, HI highly impulsive, IL infralimbic cortex, PrL prelimbic cortex, Cg cingulate cortex, DLS dorsolateral striatum, DMS dorsomedial striatum, NAc Core nucleus accumbens core, NAc Shell nucleus accumbens shell. Data are shown as means  $\pm$  SEM. HI-PPX vs. HI-Veh  $^*p < 0.05$ ;  $^{**}p < 0.01$ . HI-PPX vs. LI-PPX  $^{\#}p < 0.05$ . LI-Veh vs. LI-PPX  $^{\$}p < 0.05$ ;  $^{SS}p < 0.01$ ;  $^{SSS}p < 0.001$ . MI-Veh vs. MI-PPX  $^{\xi}p < 0.05$ ;  $^{\xi\xi}p < 0.01$ . LI-Veh vs. MI-Veh  $^{+}p < 0.05$ ;  $^{++}p < 0.01$ .

$F_{3,90} = 4.24$ ,  $p = 0.007$ ,  $\eta_p^2 = 0.12$ ). This effect of the treatment was found to be independent of the impulsive phenotype of these rats [no structure  $\times$  impulsivity  $\times$  treatment interaction:  $F_{6,90} < 1$ ,  $p = 0.71$ ,

$\eta_p^2 = 0.04$ ]. Together these results suggest that a high impulsivity trait dampens the inhibition exerted by PPX on NAc Shell and DMS activity, as assessed by C-fos mRNA levels.



**Fig. 4** The PPX-induced alterations of Zif268 mRNA levels in a network involving the PrL, the NAc Core and the DLS predicted the anti-impulsive effect of the drug. **A, B** Simplified diagrams of the mediation analysis. **C** Statistical results of the mediation analysis. Upper panel: direct effect of the treatment on premature responses (i.e., impulsivity). Lower panel: indirect effects of the treatment on premature responses mediated by IEGs mRNA levels. LI-Veh  $n = 5$ ; LI-PPX  $n = 6$ ; MI-Veh  $n = 8$ ; MI-PPX  $n = 9$ ; HI-Veh  $n = 6$ ; HI-PPX  $n = 5$ . LI low impulsive, MI moderately impulsive, HI highly impulsive.

#### PPX-induced alterations of Zif268 mRNA levels in a network involving the PrL, the NAc Core and the DLS predicted the anti-impulsive effect of the drug

We then carried-out a mediation analysis in order to identify which of the functional changes in the prefrontal corticostriatal circuitry described above predicted best the anti-impulsive effect of PPX. Mediation analysis tests the validity of a hypothetical causal chain in which one variable, here PPX treatment, affects a second variable, here Zif268 and C-fos expression, which, in turn, affects a third variable, impulsivity, in a so-called indirect effect. Thus, in our statistical model, mRNA levels of Zif268 or C-fos in the different corticostriatal structures would “mediate” the relationship between the predictor, treatment (PPX), and the behavioral outcome, impulsivity (premature responses during the last LITI under treatment) (Fig. 4A, B). The first outcome of the analysis was a negative estimate c between treatment and impulsivity (Fig. 4C upper panel) which confirmed the reduction of LITI-exacerbated impulsivity by PPX, thereby representing a first validation of the model. The indirect effect analysis also confirmed that PPX decreased both C-fos and Zif268 mRNA expression in the seven structures investigated (i.e negative estimate a; Fig. 4C lower panel). While no significant relationship with C-fos expression was identified, the mediation analysis revealed that a reduction of Zif268 mRNA level in the NAc Core, PrL and DLS was correlated with PPX-induced changes in premature responding. More

specifically, at the cohort level, a reduction of Zif268 expression in the NAc Core was positively correlated to a reduction in impulsivity (i.e positive estimate b) whereas in the PrL and the DLS it was associated with increased impulsivity (i.e negative estimate b). Overall, these results enabled the identification of a network involving the PrL, NAc Core and DLS as the neural substrate of the anti-impulsive effects of PPX.

#### DISCUSSION

This study provides a detailed behavioral characterization of the effect of PPX on the expression of a high impulsivity trait either at baseline or when challenged under long inter-trial interval conditions. Repeated PPX administration was shown to exacerbate baseline impulsivity, assessed across several 5 s ITI sessions, only in HI rats. In marked contrast, PPX prevented the exacerbation of premature responding characteristic of HI and MI rats, to a lesser extent, upon introduction of a longer 7 s ITI. In addition, the gene expression hotspot analyses carried out on the entire population revealed that the trait-dependent effect of PPX on impulse control is mediated by functional alterations of a corticostriatal network involving the PrL, the NAc Core and the DLS.

These results together challenge an established wisdom about the pro-impulsivity effects of D<sub>2</sub>-like agonists since they are demonstrated here to be dependent on pre-existing differences in

impulsivity. Impulsivity trait-dependent effects have already been described for the noradrenergic reuptake inhibitor, atomoxetine, and the psychostimulant drugs, amphetamine and methylphenidate, which all decrease premature responding under LITI in HI rats but potentiate that of LI rats [42, 73–75] (but see [76]). NAc Core deep brain stimulation (DBS) has been shown to exert the same impulsivity trait-dependent effect on impulse control in the 5-CSRTT by reducing premature responses in HI rats and tending to increase them in LI rats. Together these data highlight profound neurobiological differences between HI and LI individuals [77], which are in part associated with a reduced expression of  $D_{2/3}R$  in the ventral striatum of HI individuals, as shown by two PET scan and ISH studies [45, 74, 78]. HI rats have also been shown to have lower  $D_2$  mRNA levels than LI rats both in the nucleus accumbens and the ventral tegmental area [45], which suggests that their lower accumbal  $D_2$  dopamine receptor level, as assessed with PET, is attributable to both pre and post-synaptic decreases. It is therefore unclear whether and how, depending on the receptors dynamic and PPX doses, these pre- and post-synaptic striatal  $D_{2/3}$  dopamine receptors may account for the impulsivity trait-dependent effect of PPX.

The effect of PPX on baseline impulsivity in HI rats observed in the present study is in agreement with the previous demonstration that PPX exacerbates waiting impulsivity in non-parkinsonian animals in a DRL task [38]. In another study that used a 5-CSRTT similar to the one used here, PPX was shown to promote premature responses only in animals with a virus-mediated  $\alpha$ -synuclein-induced nigrostriatal lesion. This effect that was exacerbated during LITI, but it was not restricted to HI rats, and was instead accompanied with attentional and motivational deficits which precluded any interpretation with regards to the specific effects of PPX on impulsivity [39]. In contrast, the only side effect of the PPX treatment observed in the present study was a reduction in general activity (see [61]) or motor readiness [79] since deficits in attention or disengagement from the task were not observed in PPX-treated animals.

In marked contrast to its effect on baseline impulsivity, PPX completely blocked the exacerbation of impulsivity by an acute increase in waiting time, as operationalized under LITI, in HI rats. This observation suggests that the engagement of negative urgency in highly impulsive individuals may be a contributing factor in the transition from a pro-impulsive to an anti-impulsive effect of these drugs. Indeed, the increase in waiting time during LITI profoundly challenges the inhibitory control of highly impulsive individuals [80, 81] which experience more often the negative consequences of their premature responses, a loss in reinforcement that has been suggested to generate heightened negative urgency [82]. In a recent study, such negative urgency, the trait of making a harsh decision under stress, has been shown to mediate compulsive goal-directed relapse in individuals having developed with a DLS dopamine-dependent incentive habit for cocaine that had lost the opportunity to express it [66]. In line with this interpretation, the effects of PPX on impulsive control in HI rats may be linked to the differential impact of dopaminergic drugs on corticostriatal networks involved in the control of habitual and goal-directed behavior. Under baseline conditions, as rats become overtrained, daily stimulation of  $D_{2/3}$  receptors by PPX may strengthen habitual behavior by a time-stamped DLS-dependent stimulus-response strategy [83–85], in line with previous causal evidence based on photostimulation of the DLS indirect pathway [86]. The PPX-induced disruption of the underlying corticostriatal circuitry may result in an alteration of time-based response initiation, thereby resulting in a facilitation of responding before the end of the inter-trial interval. In contrast, under LITI conditions, the perturbation of the time-based stimulus-response strategy alongside with a potential heightened negative urgency produced by the introduction of significantly longer inter-trial intervals may result in the recruitment of a goal-directed

strategy in order to perform in the task at a time the less cognitively demanding habits are no longer adaptive. The functional engagement of the underlying PrL/Cg to DMS circuitry, which is also important for inhibitory control [87, 88], may be facilitated in HI rats by a PPX-induced decrease in the functional engagement of striatal components of the habit system. This strongly suggests that PPX may facilitate the transition from a time-stamped habitual strategy to one that depends on the goal-directed system [25, 87] when the duration of the ITI is increased, thereby enhancing inhibitory control under LITI in HI rats.

In the present study, we conducted a thorough hotspot analysis to investigate the expression patterns of C-fos and Zif268. These IEGs exhibited contrasting expression profiles due to their unique roles and regulatory mechanisms in neuronal activity and plasticity. C-fos, stands as a reliable indicator of neuronal activation, its heightened expression consistently links to various facets of neural activity, such as learning and memory retrieval, underscoring its significance as an activity marker [89, 90]. It codes for the fos protein which dimerizes to form the AP-1 complex, involved in the regulation of the transcription of a broad range of target genes [48, 91]. Zif268 is involved in related yet distinctive neural processes. While studies support its involvement in learning and memory formation, its specific contribution to plasticity involves a wide array of genes associated with vesicular transport, neurotransmitter release, clathrin-dependent pathways, and actin regulation [92, 93]. This complex network of target genes highlights Zif268's role in synaptic plasticity, thereby contributing to learning and memory. The close but yet different role in cellular processes may therefore underlie the differences in expression profile of C-fos and Zif268 observed in the present study and others [94], with potential complementary roles and information.

At a structure level, prior research has underscored the distinct contribution of the sub-regions of the mPFC in regulating inhibitory control and attention, with its ventral part (IL/PrL) being involved in preparatory attention and cue processing, while the dorsal mPFC (PrL/Cg) is necessary for suppressing irrelevant behavior [87, 95–99] (but see [100]). At the cellular level, the activation of  $D_2/D_3$  dopamine receptors by PPX results in an inhibition of the activity of the PKA and its downstream transduction signaling [101] which leads to a reduction in synaptic strength and eventually promotes long term synaptic depression [102, 103]. In the present study, PPX treatment resulted in a decrease in the expression of Zif268, a marker critical for synaptic plasticity, particularly the maintenance of long-term potentiation (LTP) [55, 104]. The reduced Zif268 expression detected in the prefrontal (PrL and Cg cortex) and striatal regions of HI-PPX rats may indicate a shift towards a more balanced functioning of the goal-directed system in HI rats. In addition, considering the functional antagonism in the control over waiting impulsivity between the NAc Shell and the NAc Core, the latter promoting impulsive action and the former inhibiting it [105–109], the PPX-induced decrease in the mRNA levels of both C-fos and Zif268 in the NAc Core associated with sustained C-fos expression in the Shell of HI rats suggests a PPX-induced ventral striatum-mediated normalization of impulse control at a time HI rats have re-engaged their goal-directed system. While the present results highlight the role of the NAc Core in the modulatory effects of PPX on impulsivity, other studies have also previously demonstrated using a similar 5-CSRTT paradigm that the NAc Shell [107, 108], and its innervation by the VTA [110, 111] also contribute to the control of impulsivity.

The decreased NAc Core Zif268 mRNA levels associated with optimal impulse control observed here in HI-PPX rats is similar to the profile of LI rats previously established by Besson and colleagues [45]. Together these observations suggest that a decrease in synaptic plasticity in the NAc Core may be a key determinant of behavioral inhibition. In contrast with the Besson study however, no differences were observed here between HI-



Veh and LI-Veh rats in the mRNA levels of Zif268 and C-fos in the NAc Core. This apparent discrepancy may be attributable to the difference in the rat strain used in the two studies, namely Sprague Dawley in the present one and Lister-Hooded in the earlier, a difference in housing conditions (standard versus reversed cycle respectively), a difference in cue-light duration (0.5 vs. 1 s in the Besson and the present study, respectively), which may result in a lower cognitive load in the present study [59, 112], as well as differences in training history. Indeed, as compared to the Besson study, rats in the present one were also trained under differential treatment for an additional 12 sessions, an extension of their training history that may have resulted in a lesser recruitment of cellular plasticity processes overall [113].

Taken together, the results of the present study demonstrate a selective anti-impulsive effect of PPX in HI rats under conditions of heightened negative urgency, which is correlated with a specific diminution of the expression of the plasticity marker Zif268 in the NAc Core. These results also identified a specific expression pattern of the cell activity marker, C-fos, maintained by PPX in HI rats in pro-inhibition structures, namely the NAc Shell and DMS and reduced in the pro-impulsive structure, the NAc Core, suggestive of an increased functional recruitment of the mPFC-DMS goal directed system at the expense of the DLS-dependent habit system. These results thereby shed new light on the hitherto accepted pro-impulsive effect of D<sub>2</sub>-like agonists on waiting impulsivity, by revealing a state dependent effect modulated by different corticostriatal circuits. Although further studies are needed to determine whether similar effects are observed in females, these results suggest that targeting D<sub>2</sub>-like receptors may represent a valuable strategy to improve, or normalize, waiting impulsivity in psychiatric disorders in which this dimension of impulsivity is aberrantly exacerbated, such as ADHD or addictions.

## DATA AVAILABILITY

Data are available upon request.

## REFERENCES

- Houeto J-L, Magnard R, Dalley JW, Belin D, Carnicella S. Trait Impulsivity and Anhedonia: Two Gateways for the Development of Impulse Control Disorders in Parkinson's Disease? *Front Psychiatry*. 2016;7:91 <https://doi.org/10.3389/fpsyt.2016.00091>.
- Weintraub D, Koester J, Potenza MN, Siderowf AD, Stacy M, Voon V, et al. Impulse Control Disorders in Parkinson Disease: A Cross-Sectional Study of 3090 Patients. *Arch Neurol*. 2010;67:589–95. <https://doi.org/10.1001/archneurol.2010.65>.
- Voon V, Sohr M, Lang AE, Potenza MN, Siderowf AD, Whetteckey J, et al. Impulse control disorders in parkinson disease: A multicenter case-control study. *Ann Neurol*. 2011;69:986–96.
- Cornelius JR, Tippmann-Peikert M, Slocumb NL, Frerichs CF, Silber MH. Impulse Control Disorders with the use of Dopaminergic Agents in Restless Legs Syndrome: a Case-Control Study. *Sleep*. 2010;33:10.
- Lipford MC, Silber MH. Long-term use of pramipexole in the management of restless legs syndrome. *Sleep Med*. 2012;13:1280–5.
- Voon V, Schoerling A, Wenzel S, Ekanayake V, Reiff J, Trenkwalder C, et al. Frequency of impulse control behaviours associated with dopaminergic therapy in restless legs syndrome. *BMC Neurol*. 2011;11:117.
- Holman AJ. Impulse Control Disorder Behaviors Associated with Pramipexole Used to Treat Fibromyalgia. *J Gambl Stud*. 2009;25:425–31.
- Bancos I, Nannenga MR, Bostwick JM, Silber MH, Erickson D, Nippoldt TB. Impulse control disorders in patients with dopamine agonist-treated prolactinomas and nonfunctioning pituitary adenomas: a case-control study. *Clin Endocrinol*. 2014;80:863–8.
- Celik E, Ozkaya HM, Poyraz BC, Saglam T, Kadioglu P. Impulse control disorders in patients with prolactinoma receiving dopamine agonist therapy: a prospective study with 1 year follow-up. *Endocrine*. 2018;62:692–700.
- Dogansen SC, Cikrikcili U, Oruk G, Kutbay NO, Tanrikulu S, Hekimsoy Z, et al. Dopamine Agonist-Induced Impulse Control Disorders in Patients With Prolactinoma: A Cross-Sectional Multicenter Study. *J Clin Endocrinol Metab*. 2019;104:2527–34.
- Evans AH, Lees AJ. Dopamine dysregulation syndrome in Parkinson's disease. *Curr Opin Neurol*. 2004;17:393–8.
- Aarsland D, Bronnick K, Alves G, Tysnes OB, Pedersen KF, Ehrst U, et al. The spectrum of neuropsychiatric symptoms in patients with early untreated Parkinson's disease. *J Neurol Neurosurg Psychiatry*. 2009;80:928–30.
- Rabinak CA, Nirenberg MJ. Dopamine Agonist Withdrawal Syndrome in Parkinson Disease. *Arch Neurol*. 2010;67:58–63. <https://doi.org/10.1001/archneurol.2009.294>.
- Leeman RF, Potenza MN. Similarities and differences between pathological gambling and substance use disorders: a focus on impulsivity and compulsivity. *Psychopharmacology*. 2012;219:469–90.
- Aarts E, Helmich RC, Janssen MJR, Oyen WJG, Bloem BR, Cools R. Aberrant reward processing in Parkinson's disease is associated with dopamine cell loss. *NeuroImage*. 2012;59:3339–46.
- American Psychiatric Association, American Psychiatric Association (eds.). *Diagnostic and statistical manual of mental disorders: DSM-5*. 5th ed. Washington, D.C: American Psychiatric Association; 2013.
- Jentsch JD, Ashenurst JR, Cervantes MC, Groman SM, James AS, Pennington ZT. Dissecting impulsivity and its relationships to drug addictions: Impulsivity subtypes and drug addiction. *Ann N Y Acad Sci*. 2014;1327:1–26.
- Laque A, Wagner GE, Matzeu A, De Ness GL, Kerr TM, Carroll AM, et al. Linking drug and food addiction via compulsive appetite. *Br J Pharm*. 2022;179:2589–609.
- Karim R, Chaudhri P. Behavioral Addictions: An Overview. *J Psychoact Drugs*. 2012;44:5–17.
- Yau YHC, Potenza MN. Gambling Disorder and Other Behavioral Addictions: Recognition and Treatment. *Harv Rev Psychiatry*. 2015;23:134–46.
- Robbins T, Clark L. Behavioral addictions. *Curr Opin Neurobiol*. 2015;30:66–72.
- Perry JL, Carroll ME. The role of impulsive behavior in drug abuse. *Psychopharmacology*. 2008;200:1–26.
- Potenza MN, Balodis IM, Derevensky J, Grant JE, Petry NM, Verdejo-Garcia A, et al. Gambling disorder. *Nat Rev Dis Prim*. 2019;5:51.
- Verdejo-Garcia A, Albein-Urios N. Impulsivity traits and neurocognitive mechanisms conferring vulnerability to substance use disorders. *Neuropharmacology*. 2021;183:108402.
- Dalley JW, Everitt BJ, Robbins TW. Impulsivity, Compulsivity, and Top-Down Cognitive Control. *Neuron*. 2011;69:680–94.
- Evenden JL. Varieties of impulsivity. *Psychopharmacology*. 1999;146:348–61.
- Madden GJ, Johnson PS, Brewer AT, Pinkston JW, Fowler SC. Effects of pramipexole on impulsive choice in male wistar rats. *Exp Clin Psychopharmacol*. 2010;18:267–76.
- Koffarnus MN, Newman AH, Grundt P, Rice KC, Woods JH. Effects of selective dopaminergic compounds on a delay-discounting task. *Behav Pharm*. 2011;22:300–11.
- Martinez E, Pasquereau B, Saga Y, Météreau É, Tremblay L. The Anterior Caudate Nucleus Supports Impulsive Choices Triggered by Pramipexole. *Mov Disord*. 2020;35:296–305.
- Johnson PS, Madden GJ, Brewer AT, Pinkston JW, Fowler SC. Effects of acute pramipexole on preference for gambling-like schedules of reinforcement in rats. *Psychopharmacology*. 2011;213:11–18.
- Holtz NA, Tedford SE, Persons AL, Grasso SA, Napier TC. Pharmacologically distinct pramipexole-mediated akinesia vs. risk-taking in a rat model of Parkinson's disease. *Prog Neuropsychopharmacol Biol Psychiatry*. 2016;70:77–84.
- Pes R, Godar SC, Fox AT, Burgeno LM, Strathman HJ, Jarmolowicz DP, et al. Pramipexole enhances disadvantageous decision-making: Lack of relation to changes in phasic dopamine release. *Neuropharmacology*. 2017;114:77–87.
- Cole BJ, Robbins TW. Effects of 6-hydroxydopamine lesions of the nucleus accumbens septi on performance of a 5-choice serial reaction time task in rats: Implications for theories of selective attention and arousal. *Behav Brain Res*. 1989;33:165–79.
- Robbins T. The 5-choice serial reaction time task: behavioural pharmacology and functional neurochemistry. *Psychopharmacology*. 2002;163:362–80.
- Molander AC, Mar A, Norbury A, Steventon S, Moreno M, Caprioli D, et al. High impulsivity predicting vulnerability to cocaine addiction in rats: some relationship with novelty preference but not novelty reactivity, anxiety or stress. *Psychopharmacology*. 2011;215:721–31.
- Robinson ESJ, Eagle DM, Mar AC, Bari A, Banerjee G, Jiang X, et al. Similar Effects of the Selective Noradrenaline Reuptake Inhibitor Atomoxetine on Three Distinct Forms of Impulsivity in the Rat. *Neuropsychopharmacology*. 2008;33:1028–37.
- Belin-Rauscent A, Daniel M-L, Puaud M, Jupp B, Sawiak S, Howett D, et al. From impulses to maladaptive actions: the insula is a neurobiological gate for the development of compulsive behavior. *Mol Psychiatry*. 2016;21:491–9.

38. Engeln M, Ansquer S, Dugast E, Bezard E, Belin D, Fernagut P-O. Multi-faceted impulsivity following nigral degeneration and dopamine replacement therapy. *Neuropharmacology*. 2016;109:69–77.
39. Jiménez-Urbieta H, Gago B, Quiroga-Varela A, Rodríguez-Chinchilla T, Merino-Galán L, Oregi A, et al. Pramipexole-induced impulsivity in mildparkinsonian rats: a model of impulse control disorders in Parkinson's disease. *Neurobiol Aging*. 2019;75:126–35.
40. Belin D, Mar AC, Dalley JW, Robbins TW, Everitt BJ. High Impulsivity Predicts the Switch to Compulsive Cocaine-Taking. *Science*. 2008;320:1352–5.
41. Ersche KD, Barnes A, Jones PS, Morein-Zamir S, Robbins TW, Bullmore ET. Abnormal structure of frontostriatal brain systems is associated with aspects of impulsivity and compulsivity in cocaine dependence. *Brain*. 2011;134:2013–24.
42. Ansquer S, Belin-Rauscent A, Dugast E, Duran T, Benatru I, Mar AC, et al. Atomoxetine Decreases Vulnerability to Develop Compulsivity in High Impulsive Rats. *Biol Psychiatry*. 2014;75:825–32.
43. Jones JA, Belin-Rauscent A, Jupp B, Fouyssac M, Sawiak SJ, Zuhlsdorff K et al. Neurobehavioral precursors of compulsive cocaine-seeking in dual frontostriatal circuits. *Neuroscience*. 2022. <https://doi.org/10.1101/2022.11.09.515779>.
44. Eagle DM, Baunez C. Is there an inhibitory-response-control system in the rat? Evidence from anatomical and pharmacological studies of behavioral inhibition. *Neurosci Biobehav Rev*. 2010;34:50–72.
45. Besson M, Pelloux Y, Dilleen R, Theobald DE, Lyon A, Belin-Rauscent A, et al. Cocaine Modulation of Frontostriatal Expression of Zif268, D2, and 5-HT2c Receptors in High and Low Impulsive Rats. *Neuropsychopharmacology*. 2013;38:1963–73.
46. Murray JE, Dilleen R, Pelloux Y, Economidou D, Dalley JW, Belin D, et al. Increased Impulsivity Retards the Transition to Dorsolateral Striatal Dopamine Control of Cocaine Seeking. *Biol Psychiatry*. 2014;76:15–22.
47. Giuliano C, Marti-Prats L, Domi A, Puaud M, Pena-Oliver Y, McKenzie C et al. The development of compulsive coping behaviors depends on the engagement of dorsolateral striatum dopamine-dependent mechanisms. *Neuroscience*. 2022. <https://doi.org/10.1101/2022.09.27.509680>.
48. Hope B, Kosofsky B, Hyman SE, Nestler EJ. Regulation of immediate early gene expression and AP-1 binding in the rat nucleus accumbens by chronic cocaine. *Proc Natl Acad Sci*. 1992;89:5764–8.
49. Crombag HS, Jedynek JP, Redmond K, Robinson TE, Hope BT. Locomotor sensitization to cocaine is associated with increased Fos expression in the accumbens, but not in the caudate. *Behav Brain Res*. 2002;136:455–62.
50. Cruz FC, Javier Rubio F, Hope BT. Using c-fos to study neuronal ensembles in corticostriatal circuitry of addiction. *Brain Res*. 2015;1628:157–73.
51. Thomas KL, Arroyo M, Everitt BJ. Induction of the learning and plasticity-associated gene *Zif268* following exposure to a discrete cocaine-associated stimulus: Cocaine CS induced *Zif268* expression. *Eur J Neurosci*. 2003;17:1964–72.
52. Valjent E. Plasticity-Associated Gene *Krox24/Zif268* Is Required for Long-Lasting Behavioral Effects of Cocaine. *J Neurosci*. 2006;26:4956–60.
53. Hearing MC, See RE, McGinty JF. Relapse to cocaine-seeking increases activity-regulated gene expression differentially in the striatum and cerebral cortex of rats following short or long periods of abstinence. *Brain Struct Funct*. 2008;213:215–27.
54. Unal CT, Beverley JA, Willuhn I, Steiner H. Long-lasting dysregulation of gene expression in corticostriatal circuits after repeated cocaine treatment in adult rats: effects on *zif 268* and *homer 1a*. *Eur J Neurosci*. 2009;29:1615–26.
55. Veyrac A, Besnard A, Caboche J, Davis S, Laroche S. The Transcription Factor *Zif268/Egr1*, Brain Plasticity, and Memory. *Prog Mol Biol Transl Sci*. 2014;122:89–129.
56. Carli M, Robbins TW, Evenden JL, Everitt BJ. Effects of lesions to ascending noradrenergic neurones on performance of a 5-choice serial reaction task in rats; implications for theories of dorsal noradrenergic bundle function based on selective attention and arousal. *Behav Brain Res*. 1983;9:361–80.
57. Bari A, Dalley JW, Robbins TW. The application of the 5-choice serial reaction time task for the assessment of visual attentional processes and impulse control in rats. *Nat Protoc*. 2008;3:759–67.
58. Auclair AL, Besnard J, Newman-Tancredi A, Depoortère R. The five choice serial reaction time task: Comparison between Sprague–Dawley and Long–Evans rats on acquisition of task, and sensitivity to phencyclidine. *Pharm Biochem Behav*. 2009;92:363–9.
59. Turner KM, Peak J, Burne THJ. Measuring Attention in Rodents: Comparison of a Modified Signal Detection Task and the 5-Choice Serial Reaction Time Task. *Front Behav Neurosci*. 2016;9:370 <https://doi.org/10.3389/fnbeh.2015.00370>.
60. Turner KM, Young JW, McGrath JJ, Eyles DW, Burne THJ. Cognitive performance and response inhibition in developmentally vitamin D (DVD)-deficient rats. *Behav Brain Res*. 2013;242:47–53.
61. Favier M, Duran T, Carcenac C, Drui G, Savasta M, Carnicella S. Pramipexole reverses Parkinson's disease-related motivational deficits in rats: Pramipexole and Motivation. *Mov Disord*. 2014;29:912–20.
62. Magnard R, Vachez Y, Carcenac C, Krack P, David O, Savasta M, et al. What can rodent models tell us about apathy and associated neuropsychiatric symptoms in Parkinson's disease? *Transl Psychiatry*. 2016;6:e753.
63. Chang W, Breier MR, Yang A, Swerdlow NR. Disparate effects of pramipexole on locomotor activity and sensorimotor gating in Sprague–Dawley rats. *Pharm Biochem Behav*. 2011;99:634–8.
64. Maj J, Rogóż Z, Skuza G, Kołodziejczyk K. The behavioural effects of pramipexole, a novel dopamine receptor agonist. *Eur J Pharm*. 1997;324:31–7.
65. Belin D, Deroche-Gamonet V, Jaber M. Cocaine-induced sensitization is associated with altered dynamics of transcriptional responses of the dopamine transporter, tyrosine hydroxylase, and dopamine D2 receptors in C57Bl/6J mice. *Psychopharmacology*. 2007;193:567–78.
66. Fouyssac M, Peña-Oliver Y, Puaud M, Lim NTY, Giuliano C, Everitt BJ, et al. Negative Urgency Exacerbates Relapse to Cocaine Seeking After Abstinence. *Biol Psychiatry*. 2022;91:1051–60.
67. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*. 2012;9:671–5.
68. Paxinos G, Watson C. The rat brain in stereotaxic coordinates. San diego: Elsevier Academic Press; 1998.
69. Levine TR, Hulleit CR. Eta Squared, Partial Eta Squared, and Misreporting of Effect Size in Communication Research. *Hum Commun Res*. 2002;28:612–25.
70. Magnard R, Vachez Y, Carcenac C, Boulet S, Houeto J-L, Savasta M, et al. Nigrostriatal Dopaminergic Denervation Does Not Promote Impulsive Choice in the Rat: Implication for Impulse Control Disorders in Parkinson's Disease. *Front Behav Neurosci*. 2018;12:312.
71. Baron RM, Kenny DA. The Moderator-Mediator Variable Distinction in Social Psychological Research: Conceptual, Strategic, and Statistical Considerations. *J Pers Soc Psychol*. 1986;51:1173–82.
72. Ditlevsen S, Christensen U, Lynch J, Damsgaard MT, Keiding N. The Mediation Proportion: A Structural Equation Approach for Estimating the Proportion of Exposure Effect on Outcome Explained by an Intermediate Variable. *Epidemiology*. 2005;16:114–20.
73. Tomlinson A, Grayson B, Marsh S, Harte MK, Barnes SA, Marshall KM, et al. Pay attention to impulsivity: Modelling low attentive and high impulsive subtypes of adult ADHD in the 5-choice continuous performance task (5C-CPT) in female rats. *Eur Neuropsychopharmacol*. 2014;24:1371–80.
74. Caprioli D, Jupp B, Hong YT, Sawiak SJ, Ferrari V, Wharton L, et al. Dissociable Rate-Dependent Effects of Oral Methylphenidate on Impulsivity and D2/3 Receptor Availability in the Striatum. *J Neurosci*. 2015;35:3747–55.
75. Higgins GA, Silenieks LB, MacMillan C, Thevarkunnel S, Parachikova AI, Mombereau C, et al. Characterization of Amphetamine, Methylphenidate, Nicotine, and Atomoxetine on Measures of Attention, Impulsive Action, and Motivation in the Rat: Implications for Translational Research. *Front Pharm*. 2020;11:427.
76. Fernando ABP, Economidou D, Theobald DE, Zou M-F, Newman AH, Spoelder M, et al. Modulation of high impulsivity and attentional performance in rats by selective direct and indirect dopaminergic and noradrenergic receptor agonists. *Psychopharmacology*. 2012;219:341–52.
77. Schippers MC, Bruinsma B, Gaastra M, Mesman TI, Denys D, De Vries TJ, et al. Deep Brain Stimulation of the Nucleus Accumbens Core Affects Trait Impulsivity in a Baseline-Dependent Manner. *Front Behav Neurosci*. 2017;11:52 <https://doi.org/10.3389/fnbeh.2017.00052>.
78. Dalley JW, Fryer TD, Brichard L, Robinson ESJ, Theobald DEH, Lääne K et al. Nucleus Accumbens D2/3 Receptors Predict Trait Impulsivity and Cocaine Reinforcement. *Science*. 2007;315:1267–70.
79. Brown VJ, Robbins TW. Simple and choice reaction time performance following unilateral striatal dopamine depletion in the rat: impaired motor readiness but preserved response preparation. *Brain*. 1991;114:513–25.
80. Zeeb FD, Soko AD, Ji X, Fletcher PJ. Low Impulsive Action, but not Impulsive Choice, Predicts Greater Conditioned Reinforcer Salience and Augmented Nucleus Accumbens Dopamine Release. *Neuropsychopharmacology*. 2016;41:2091–2100.
81. Fitzpatrick CM, Maric VS, Bate ST, Andreasen JT. Influence of intertrial interval on basal and drug-induced impulsive action in the 5-choice serial reaction time task: Effects of d-amphetamine and (±)-2,5-dimethoxy-4-iodoamphetamine (DOI). *Neurosci Lett*. 2018;662:351–5.
82. Toschi C, El-Sayed Hervig M, Burghi T, Sell T, Lycas MD, Moazen P, et al. Dissociating reward sensitivity and negative urgency effects on impulsivity in the five-choice serial reaction time task. *Brain Neurosci Adv*. 2022;6:239821282211022.
83. Yin HH, Knowlton BJ, Balleine BW. Lesions of dorsolateral striatum preserve outcome expectancy but disrupt habit formation in instrumental learning. *Eur J Neurosci*. 2004;19:181–9.

84. Faure A. Lesion to the Nigrostriatal Dopamine System Disrupts Stimulus-Response Habit Formation. *J Neurosci*. 2005;25:2771–80.
85. Nelson A, Killcross S. Amphetamine Exposure Enhances Habit Formation. *J Neurosci*. 2006;26:3805–12.
86. Vicente AM, Galvão-Ferreira P, Tecuapetla F, Costa RM. Direct and indirect dorsolateral striatum pathways reinforce different action strategies. *Curr Biol*. 2016;26:R267–9.
87. Terra H, Bruinsma B, de Kloet SF, van der Roest M, Pattij T, Mansvellder HD. Prefrontal Cortical Projection Neurons Targeting Dorsomedial Striatum Control Behavioral Inhibition. *Curr Biol*. 2020;30:4188–4200.e5.
88. Brockett AT, Tennyson SS, deBettencourt CA, Gaye F, Roesch MR. Anterior cingulate cortex is necessary for adaptation of action plans. *Proc Natl Acad Sci*. 2020;117:6196–204.
89. Maviel T, Durkin TP, Menzaghi F, Bontempi B. Sites of Neocortical Reorganization Critical for Remote Spatial Memory. *Science*. 2004;305:96–9.
90. Lopez J, Herbeaux K, Cosquer B, Engeln M, Muller C, Lazarus C, et al. Context-dependent modulation of hippocampal and cortical recruitment during remote spatial memory retrieval. *Hippocampus*. 2012;22:827–41.
91. Chiu R, Boyle WJ, Meek J, Smeal T, Hunter T, Karin M. The c-fos protein interacts with c-Jun/AP-1 to stimulate transcription of AP-1 responsive genes. *Cell*. 1988;54:541–52.
92. Koldamova R, Schug J, Lefterova M, Cronican AA, Fitz NF, Davenport FA, et al. Genome-wide approaches reveal EGR1-controlled regulatory networks associated with neurodegeneration. *Neurobiol Dis*. 2014;63:107–14.
93. Duclot F, Kabbaj M. The Role of Early Growth Response 1 (EGR1) in Brain Plasticity and Neuropsychiatric Disorders. *Front Behav Neurosci*. 2017;11:35 <https://doi.org/10.3389/fnbeh.2017.00035>.
94. Guzowski JF, Setlow B, Wagner EK, McGaugh JL. Experience-Dependent Gene Expression in the Rat Hippocampus after Spatial Learning: A Comparison of the Immediate-Early Genes *Arc*, *c-fos*, and *zif268*. *J Neurosci*. 2001;21:5089–98.
95. Muir JL, Everitt BJ, Robbins TW. The Cerebral Cortex of the Rat and Visual Attentional Function: Dissociable Effects of Medial Frontal, Cingulate, Anterior Dorsolateral, and Parietal Cortex Lesions on a Five-Choice Serial Reaction Time Task. *Cereb Cortex*. 1996;6:470–81.
96. Narayanan NS, Laubach M. Top-Down Control of Motor Cortex Ensembles by Dorsomedial Prefrontal Cortex. *Neuron*. 2006;52:921–31.
97. Pezze M, McGarrity S, Mason R, Fone KC, Bast T. Too Little and Too Much: Hypoactivation and Disinhibition of Medial Prefrontal Cortex Cause Attentional Deficits. *J Neurosci*. 2014;34:7931–46.
98. Luchicchi A, Mnie-Filali O, Terra H, Bruinsma B, de Kloet SF, Obermayer J, et al. Sustained Attentional States Require Distinct Temporal Involvement of the Dorsal and Ventral Medial Prefrontal Cortex. *Front Neural Circuits*. 2016;10:70 <https://doi.org/10.3389/fncir.2016.00070>.
99. Hardung S, Epple R, Jäckel Z, Eriksson D, Uran C, Senn V, et al. A Functional Gradient in the Rodent Prefrontal Cortex Supports Behavioral Inhibition. *Curr Biol*. 2017;27:549–55.
100. Chudasama Y, Passetti F, Rhodes SEV, Lopian D, Desai A, Robbins TW. Dissociable aspects of performance on the 5-choice serial reaction time task following lesions of the dorsal anterior cingulate, infralimbic and orbitofrontal cortex in the rat: differential effects on selectivity, impulsivity and compulsivity. *Behav Brain Res*. 2003;146:105–19.
101. Neve KA, Seamans JK, Trantham-Davidson H. Dopamine Receptor Signaling. *J Recept Signal Transduct*. 2004;24:165–205.
102. Sun X, Zhao Y, Wolf ME. Dopamine Receptor Stimulation Modulates AMPA Receptor Synaptic Insertion in Prefrontal Cortex Neurons. *J Neurosci*. 2005;25:7342–51.
103. Shen W, Flajolet M, Greengard P, Surmeier DJ. Dichotomous Dopaminergic Control of Striatal Synaptic Plasticity. *Science*. 2008;321:848–51.
104. Jones MW, Errington ML, French PJ, Fine A, Bliss TVP, Garell S, et al. A requirement for the immediate early gene *Zif268* in the expression of late LTP and long-term memories. *Nat Neurosci*. 2001;4:289–96.
105. Sesia T, Temel Y, Lim LW, Blokland A, Steinbusch HWM, Visser-Vandewalle V. Deep brain stimulation of the nucleus accumbens core and shell: Opposite effects on impulsive action. *Exp Neurol*. 2008;214:135–9.
106. Sesia T, Bulthuis V, Tan S, Lim LW, Vlaming R, Blokland A, et al. Deep brain stimulation of the nucleus accumbens shell increases impulsive behavior and tissue levels of dopamine and serotonin. *Exp Neurol*. 2010;225:302–9.
107. Moreno M, Economidou D, Mar AC, López-Granero C, Caprioli D, Theobald DE, et al. Divergent effects of D2/3 receptor activation in the nucleus accumbens core and shell on impulsivity and locomotor activity in high and low impulsive rats. *Psychopharmacology*. 2013;228:19–30.
108. Feja M, Hayn L, Koch M. Nucleus accumbens core and shell inactivation differentially affects impulsive behaviours in rats. *Prog Neuropsychopharmacol Biol Psychiatry*. 2014;54:31–42.
109. Feja M, Koch M. Frontostriatal systems comprising connections between ventral medial prefrontal cortex and nucleus accumbens subregions differentially regulate motor impulse control in rats. *Psychopharmacology*. 2015;232:1291–302.
110. Flores-Dourojeanni JP, van Rijt C, van den Munkhof MH, Boekhoudt L, Luijendijk MCM, Vanderschuren LJM, et al. Temporally Specific Roles of Ventral Tegmental Area Projections to the Nucleus Accumbens and Prefrontal Cortex in Attention and Impulse Control. *J Neurosci*. 2021;41:4293–304.
111. Flores-Dourojeanni JP, Van Den Munkhof MH, Luijendijk MCM, Vanderschuren LJM, Adan RAH. Inhibition of ventral tegmental area projections to the nucleus accumbens shell increases premature responding in the five-choice serial reaction time task in rats. *Brain Struct Funct*. 2023;228:787–98.
112. Caballero-Puntiverio M, Fitzpatrick CM, Woldbye DP, Andreasen JT. Effects of amphetamine and methylphenidate on attentional performance and impulsivity in the mouse 5-Choice Serial Reaction Time Task. *J Psychopharmacology*. 2017;31:272–83.
113. Hernandez PJ, Schiltz CA, Kelley AE. Dynamic shifts in corticostriatal expression patterns of the immediate early genes *Homer 1a* and *Zif268* during early and late phases of instrumental training. *Learn Mem*. 2006;13:599–608.

## ACKNOWLEDGEMENTS

The authors want to thank Dr Bizhu He for help with structural equation modeling analysis.

## AUTHOR CONTRIBUTIONS

RM, DB and SC designed experiments with inputs from YMV and SB. RM, MF, YV, TD, and CC performed experiments with help from SB, DB and SC. RM performed the analysis, with help from DB, SC and YC. RM, DB and SC wrote the manuscript with inputs from MF, YMV, YC, TD, CC, SB, MS and PHJ.

## FUNDING

This work was supported by the Institut National de la Santé et de la Recherche Médicale (Inserm), Association France Parkinson, Agence Nationale de la Recherche (ANR13 SAMA001401) and Grenoble Alpes University.

## COMPETING INTERESTS

The authors declare no competing interests.

## ADDITIONAL INFORMATION

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41398-024-02804-3>.

**Correspondence** and requests for materials should be addressed to Robin Magnard.

**Reprints and permission information** is available at <http://www.nature.com/reprints>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024