sulfiram-like reaction with metronidazole and ethanol. *Ann Pharmacother* 2002; **36**: 971-974.

5. Stockley's Drug Interactions. Alcohol + metronidazole and related drugs. 2016. Available at: <https://www.medicinescomplete.com/#/content/stockley/x02-0033#00000166> (accessed April 2020).
6. Mergenhagen K A, Wattengel B A, Skelly M K et al. Fact vs. fiction: the evidence behind alcohol and antibiotic interactions, a review. *Antimicrob Agents Chemother* 2020; **64**: pii: e02167-19.

<https://doi.org/10.1038/s41415-020-2012-x>

Oral surgery

New MRONJ guidance

Sir, there has been guidance for dentists on the prevention of medication-related osteonecrosis of the jaws (MRONJ) for several years.¹ Although this guidance also included information for prescribers and dispensers, it does not appear to have been widely disseminated. Indeed, only 4% of general medical practitioners in a Birmingham study were aware of MRONJ guidelines.² I therefore welcome the publication by the Royal College of Physicians for the multi-disciplinary team.³

MRONJ is defined as exposed bone, or bone that can be probed through a fistula, in the maxilla or mandible, that has been present for more than eight weeks.⁴ It is an adverse side effect of anti-angiogenic or anti-resorptive medication.^{1,3} The incidence of MRONJ in patients who take these medications is reported as 1% in cancer patients and 0.1% in patients with metabolic bone disease.¹

There have been instances of MRONJ occurring spontaneously, but it is most common following a dental extraction.^{1,3,4} Prevention of MRONJ involves pre-treatment screening, extraction of teeth with poor prognosis, adjustment of prostheses and education, including oral hygiene instruction and controlling risk factors such as smoking and alcohol. This has been shown to reduce the incidence of MRONJ by up to 50%.⁵

Dentists have reported poor communication with other healthcare professionals and often rely on patient recall for their medical history.⁶ The new guidance introduces dental alert cards which are to be carried by patients and shown to their dentists.³ Referral and reply letters between oncology, dentists and maxillofacial/oral surgeons have also been produced.

We should welcome and familiarise ourselves with these new methods of communication and remember to report all cases of suspected MRONJ to the MHRA.⁷ Reducing the incidence of MRONJ will have

financial benefit for the NHS, as this is a difficult condition to manage, and for severe cases may require surgery for debridement or resection.⁴

C. Frazer-Cox, North Wales, UK

References

1. Scottish Dental Clinical Effectiveness Programme. Oral health management of patients at risk of medication-related osteonecrosis of the jaws. 2017. Available at: <http://www.sdcep.org.uk/published-guidance/medication-related-osteonecrosis-of-the-jaw/> (accessed July 2020).
2. Rahman Z, Nayani S, Anstey H, Murphy M J. A survey evaluating the awareness of MRONJ within the Birmingham GMP community. *Oral Surg* 2018. Available at: <https://doi.org/10.1111/ors.12372> (accessed July 2020).
3. Royal College of Physicians. Medication-related osteonecrosis of the jaw: guidance for the oncology multidisciplinary team. 2019. Available at: <https://www.rcplondon.ac.uk/guidelines-policy/medication-related-osteonecrosis-jaw-guidance-oncology-multidisciplinary-team> (accessed July 2020).
4. Ruggiero S L, Dodson T B, Fantasia J et al. American Association of Oral and Maxillofacial Surgeons position paper on medication-related osteonecrosis of the jaw - 2014 update. *J Oral Maxillofac Surg* 2014; **72**: 1938-1956.
5. Vandone A M, Donadio M, Mozzati M et al. Impact of dental care in the prevention of bisphosphonate-associated osteonecrosis of the jaw: a single-center clinical experience. *Ann Oncol* 2012; **23**: 193-200.
6. Sturrock A, Preshaw P M, Hayes C, Wilkes S. General dental practitioners' perceptions of, and attitudes towards, improving patient safety through a multidisciplinary approach to the prevention of medication-related osteonecrosis of the jaw (MRONJ): a qualitative study in the North East of England. *BMJ Open* 2019; **9**: e029951. doi:10.1136/bmjopen-2019-029951.
7. Yellow Card. Report online: www.mhra.gov.uk/yellowcard; via the Yellow Card App or by emailing yellowcard@mhra.gov.uk.

<https://doi.org/10.1038/s41415-020-2013-9>

Dental education

Undergraduate research

Sir, I read with interest a colleague's letter on the subject of getting published.¹ I believe that research experience is an area in which many dentists including dental core trainees (DCTs) do not have any substantial experience. Although the importance of research is touched upon at undergraduate level, there are few opportunities and most of the research activity in dental schools is carried out by postgraduate students. Many dental students would benefit from a further insight into research or opportunity to carry out small research projects at an undergraduate level supervised by senior staff. This would also give them a good foundation for future projects at a postgraduate level.

At a DCT level academic clinical fellow posts address this ability for trainees to be able to carry out research; however, these are few and far between. Some DCT posts may provide an insight into research such

as critical appraisal, however, I feel there should be more scope to incorporate further research opportunities into these posts. Many trainees would be willing to assist senior colleagues in established projects or like the chance to complete small projects of their own with guidance and assistance to provide good learning opportunities as well as supporting portfolios.

G. K. Randhawa, Bristol, UK

Reference

1. Jackson M. How to get published? *Br Dent J* 2020; **228**: 317-318.

<https://doi.org/10.1038/s41415-020-2014-8>

Periodontology

Periodontal therapy and cell adhesion

Sir, periodontitis is a chronic infection and inflammation of the periodontium.¹ A relationship between periodontal therapy (PT) and reduction in risk of cardiovascular diseases has been reported.² One hypothesis is that PT reduces the expression of cell adhesion molecules that are pivotal in the pathogenesis of vascular inflammation and atherosclerosis.³ This first meta-analysis included eight studies (three randomised control trials, one prospective non-randomised study, and four retrospective). The main outcome was change in the levels of cell adhesion molecules (soluble endothelial selectin s-E-selectin-1, soluble intercellular adhesion molecule-1 s-ICAM-1, and soluble vascular cell adhesion molecule-1 s-VCAM-1) following PT. Standardised difference in means (SDM) was used for effect size (ES) measurement. An ES of 0.2-0.5 is considered small, 0.5-0.8 is medium, and more than 0.8 is large.

Of the 797 patients (445 in PT, 352 in control), men formed 62.3% (497) with 69.8% (311) and 52.8% (186) in the respective groups; the mean age of patients was 51.9 and 55.89 years respectively. PT included oral hygiene instructions, scaling and root planing, antibiotics, chlorhexidine mouthwashes, tooth extractions and surgical PT. The control group received no treatment. Patients had moderate to severe periodontitis. Follow up varied between 3-6 months. PT showed a positive and medium effect size (SDM 0.52, 95% CI 0.10-0.94) on decreasing s-E-selectin values at follow up. PT did not show a statistically significant effect size on lowering s-ICAM-1 (SDM 0.41,

95% CI – 0.002 to 0.827) and s-VCAM-1 (SDM – 0.047, 95% CI – 0.27 to 0.18).

Thus, PT has a moderate effect in lowering s-E-selectin in patients with moderate to severe periodontitis. E-selectin is expressed by the endothelium and helps in migration of leucocytes to arterial sub-endothelium in an important mechanism by which atherosclerosis starts and progresses. Lowering of E-selectin may be a mechanism by which PT improves endothelial function and cardiovascular outcomes.⁴ However, our study has limitations. None of the studies checked if the change in the biochemical markers translated into clinical outcomes.

R. Bashir, K. Changal, Ohio, USA

References

1. Akram Z, Abduljabbar T, Hassan M I A, Javed F, Vohra F. Cytokine profile in chronic periodontitis patients with and without obesity: a systematic review and meta-analysis. *Dis Markers* 2016; **2016**: 4801418.
2. Li C, Lv Z, Shi Z, Zhu Y, Wu Y, Li L, Iheozor-Ejiofor Z. Periodontal therapy for the management of cardiovascular disease in patients with chronic periodontitis. *Cochrane Database Syst Rev* 2014; CD009197.
3. Dietrich T, Webb I, Stenhouse L *et al*. Evidence summary: the relationship between oral and cardiovascular disease. *Br Dent J* 2017; **222**: 381.
4. Tonetti M S, Van Dyke T E; Working group 1 of the joint EFP/AAP workshop. Periodontitis and atherosclerotic cardiovascular disease: consensus report of the Joint EFP/AAP Workshop on Periodontitis and Systemic Diseases. *J Clin Periodontol* 2013; **40 Suppl 14**: S24-S29. <https://doi.org/10.1038/s41415-020-2015-7>

Dental products

Dietary restrictions

Sir, while we may well be aware of what medication is necessary to manage a clinical presentation, we may not always appreciate that a given preparation may be unsuitable for certain patients due to cultural and religious backgrounds with a prescription liable to cause distress, delay, and possibly offence.

For example, some observant Jews make specific dietary changes at Passover and some medications may contain ingredients which make their ingestion impermissible at this time, but permissible during other times. Online resources exist which outline these medications and the dental professional is well-advised to consult these.¹

Muslims do not consume food or water during daylight hours in the month of Ramadhan. The dosing regimen of prescribed medication may need to be altered to account for this dietary restriction. It is important to note that a discrepancy between the Lunar and Gregorian calendar means Ramadhan

falls at different times of the year each year when considered from the perspective of the Gregorian calendar.

Readers may be surprised to see the list of commonly prescribed medications and products not considered Kosher ie permitted for use.¹ These include denture and oral hygiene products, as well as certain antibiotics and analgesics. Simply having an awareness of the possible suitability of prescribed products, and that this may change at different times of the year, has the potential to save both patient and practitioner distress and delay. In addition, the dental professional may consult the local pharmacist who often has experience in the unique needs of members of the locality they serve.

However, it is imperative to state that not all patients from a given community may adhere to any dietary restrictions, and as such every patient should be treated on an individual basis with questions asked regarding any restrictions to diet before a prescription is provided.

S. Adam, N. Adam, E. Walshaw, Leeds, UK

Reference

1. Kosher.org.uk. Medicines Approved for Passover 2019. Available at: https://www.kosher.org.uk/sites/default/files/2019%20Medicines_0.pdf (accessed 4 May 2020). <https://doi.org/10.1038/s41415-020-2016-6>

Dental trauma

Milk allergy and tooth reimplantation

Sir, traumatic dental injuries (TDI) are very frequent and comprise 85% of patients presenting with injuries to the oral region.¹ In the UK, one in five children have experienced TDI to their permanent anterior teeth before leaving school, and dental avulsion is common within this group.²

One such six-year-old attended our emergency department with avulsion of the upper right permanent central incisor. The parents had stored the tooth in milk for the two hours since the injury. The history revealed that the child had asthma and an intolerance to multiple allergens including dairy products with a previous mild reaction. Taking into account the history, the tooth was thoroughly rinsed with saline and reinserted. Within minutes, the child displayed signs of anaphylaxis and was transferred to resus for treatment. He was appropriately resuscitated, the offending

tooth was removed and he made a full recovery. He was discharged the next day and follow-up was arranged for the child to attend the dental institute after discharge for dental rehabilitation.

Cow's milk protein allergies (CMPA) are a range of adverse reactions to cow's milk that can be reproduced and are immune-mediated. They are immunoglobulin E (IgE)-mediated, cell-mediated (non-IgE), or a combination of the two. It is estimated that up to 4.9% of children suffer from CMPA and although it will resolve in 85-90% by three years of age, the natural history is actively changing, showing a definite trend to persist longer.^{3,4}

Considering the increase in CMPA, and the high number of traumatic dental injuries including avulsion, we should always be mindful when reimplanting teeth stored in milk to decrease possible serious life threatening reactions.

O. Marland, E. Swinnerton, A. Power, York, UK

References

1. Locker D. Self-reported dental and oral injuries in a population of adults aged 18-50 years. *Dent Traumatol* 2007; **23**: 291-296.
2. Alkhadra T, Presling W, El-Bialy T. Prevalence of traumatic dental injuries in patients attending University of Alberta Emergency Clinic. *Open Dent J* 2016; **10**: 315-321.
3. Fiochi A, Brozek J, Schunemann H *et al*. World Allergy Organisation (WAO) diagnosis and rationale for action against Cow's milk allergy (DRACMA) guidelines. *Pediatr Allergy Immunol* 2010; **21 Suppl 21**: 1-125.
4. NHS Fife. Diagnosis and Management of Infants with Suspected Cow's Milk Protein Allergy. A guide for health-care professionals working in primary care. Available at: <https://www.fifeadtc.scot.nhs.uk/media/2330/cmpa-pathway-sept-2018-v5.pdf> (accessed 20 April 2020). <https://doi.org/10.1038/s41415-020-2017-5>

Titles and terminology

Call me doctor

Sir, it was a delight to be reminded by Douglas Pike about the copious work he undertook in the early 1990s, to enable dentists to use the courtesy title of 'Doctor' if they wish.¹

In the article, he refers to the organisations that did and do continue to decline to address dentists as Doctor. However, upon my election as Chairman of the BDA's General Dental Services Committee in January 1997, I received a congratulatory letter from the then Secretary of State for Health: it was addressed to Mr Anthony Kravitz. In my reply I politely requested that I was referred to as Doctor in future correspondence, and from that time all correspondence from the