Ludwig's angina

Sir, a 25-year-old male attended a Tier 2 primary care MOS service referred for extraction of the 36 molar under IV sedation. The referral stated the patient was suffering pain and had had multiple courses of antibiotics for repeated swellings. There was nothing of clinical importance in the medical history. A periapical radiograph was included (Fig. 1).

A large swelling was visible under his chin and he reported a three-day history of worsening swelling with increased pain and difficulty swallowing. Examination revealed a bilateral submental and submandibular enlargement which was firm, tender and warm to touch. Mouth opening was limited to less than one finger. A degree of hoarseness was noted in the voice but he was able to complete full sentences. Baseline observations were unremarkable. Due to the presence of red-flag signs a provisional diagnosis of Ludwig's angina was reached and a prompt referral to the local OMFS unit arranged. The patient was blue-lighted to resuscitation. He underwent emergency surgery involving incision and drainage with extraction of the infected 36 under GA. Post-operatively he remained in hospital for several days due to significant swelling and active extra-oral drains. Fortunately, escalation to intensive care was avoided.

True Ludwig's angina is uncommon. It is named after German physician Karl Friedrich Wilhelm von Ludwig, who first described it in 1836. The word angina is derived from the Latin word for choke (*angere*).¹ Sepsis is not uncommon with large swellings caused by a nidus of infection and can quickly result in multiple organ failure especially in those

Fig. 1 Pre-operative periapical radiograph of the lower left first molar included in referral. Deep secondary caries and associated periapical pathology noted

with underlying health conditions that cause immunosuppression such as diabetes. This case serves to remind clinicians of the signs and symptoms when assessing and managing dental abscesses. It is crucial to consider these signs and symptoms especially in this COVID-19 pandemic where remote consultations and prescribing have become more commonplace. Providing a safety net for patients will reduce the risk of serious complications and this can be done by discussing the red flags and the actions to take:

- Significant trismus
- Bilateral submental and submandibular swelling
- Muffled (hot potato) voice
- Fever
- Firm/raised floor of mouth
- Limited/altered tongue mobility
- Difficulty swallowing or drooling.

These signs can indicate a significantly worsening swelling requiring emergency management.

M. R. Miah, A. S. Ali, Manchester, UK

Reference

1. Costain N, Marrie T J. Ludwig's angina. *Am J Med* 2011; **124**: 115-117.
<https://doi.org/10.1038/s41415-020-2132-3>

Foreign body retrieval

Sir, a 38-year-old gentleman presented to A&E with a right sided focal fluctuant swelling to the submandibular and submental area which crossed the midline. The neck was diffusely swollen. He had dysphonia, odynophagia and dyspnoea. He was tachycardic and tachypnoeic and gave a history of intermittent dental abscess from the lower right quadrant further to a dental extraction many years ago. On previous occasions this had usually resolved

with courses of antibiotics. Medically the patient had a history of recreational drug use and was a heavy smoker. A provisional diagnosis of impending airway loss secondary to spreading dental sepsis from the 46 region was made.

Routine blood showed raised WBC (21.2), neutrophils (19.7), CRP (456), lactate (2.1), and urea 8.3 mmol/L, demonstrating significant cervico-facial infection. An OPG revealed a foreign body (FB) in the 46 region (Fig. 1). Urgent theatre arrangements were made in order to stabilise the airway and achieve immediate abscess drainage and attempt foreign body removal. The abscess was drained via an extraoral and intraoral approach. The FB was not retrievable with blunt dissection.

Fig. 1 Orthopantomogram showing a radiopaque foreign body in the 46 region

Post-operatively, a CT scan was carried out to locate the FB with plans for elective retrieval. With the patient's airway now safe, on questioning, he admitted attempting to clear the lower right molar socket of debris with an interdental brush and we hypothesised the FB to be its metallic core. Whilst the patient initially improved, worsening clinical signs and neck swelling necessitated a repeat CT. This demonstrated a spreading fascial space infection with further collections extending to the right anterior neck, retrosternal and antero-superior mediastinum. The position of the FB remained unchanged.

Fig. 2 Interdental brush retrieved measuring 15 mm

◀ In spite of having pre-op 3D scans, locating the FB proved to be challenging intraoperatively. Ultimately, a marker needle, in conjunction with an intra-operative C-arm and parallax technique, allowed FB position triangulation. Blunt dissection was carried out around the needle to successfully retrieve the interdental brush (Fig. 2).

C. Frazer-Cox, D. Kumar, North Wales, UK
<https://doi.org/10.1038/s41415-020-2133-2>

Go with your gut

Sir, the dental and maxillofacial team were asked to assess a 12-year-old male following the discovery of primary teeth in his ileostomy bag (Figs 1 and 2). The loss of teeth had not been noted prior to discovery. The patient has a medical history consisting of cerebral palsy, scoliosis and epilepsy, as well as being tracheostomy dependent.

The mechanism of ingestion is thought to have resulted from an epileptic seizure dislodging mobile primary teeth that were close to exfoliation. A dental assessment was conducted revealing a partial mixed primary-adult dentition. Chest and plain abdomen film radiographs were conducted to rule out ingestion of any further teeth.

Fig. 1 Exfoliated teeth found in ileostomy bag

Fig. 2 Exfoliated teeth found in ileostomy bag

As far as the authors are aware there are no recorded cases in the literature of teeth being recovered from a stoma bag. In this case there were no adverse effects; however, ingestion of teeth could have resulted in occlusion of the stoma. This patient has a tracheostomy, and as such his airway is relatively protected, though it is not a cuffed tube. However, in patients who do not have a tracheostomy, the exfoliated teeth could have been inhaled rather than swallowed, with the potential sequelae of pneumonia or a lung abscess.

For children with reduced neuromuscular control in the mixed dentition (age 6–14 years), particularly those with vulnerable airways, consideration should be given to dental assessment and vigilance of mobile teeth, allowing for planned extraction of teeth posing an aspiration or ingestion risk.

A. Davies, J. Indoe, D. Laraway, Liverpool, UK
<https://doi.org/10.1038/s41415-020-2134-1>

Histopathology Reconsidering syphilis

Sir, although broadly considered a predominantly historical disease, syphilis is on the ascendancy and dentists may be well-placed to aid in its diagnosis.

A 31-year-old male attended the oral and maxillofacial out-patient clinic for oral ulceration. Due to the COVID-19 pandemic this referral had been made following a virtual consultation only with his GP. The patient expressed a high level of concern, reporting ulcers which appeared and resolved, in different oral mucosal sites for the previous two weeks. Medical history revealed HIV, with full virological suppression. He reported a current attempt to quit smoking and feeling additional stress due to the pandemic. Recent blood tests demonstrated low serum folate, which he had commenced supplementation for in the previous days; haematinics were otherwise normal.

Oral examination revealed multiple aphthous ulcers but no suspicious mucosal abnormalities otherwise. The patient requested diagnostic biopsy stating he had 'health anxieties', despite reassurance against overt clinical malignancy. Due to COVID, his results were to be discussed by telephone but the patient was subsequently non-contactable. A letter was therefore dictated,

explaining that the histopathology demonstrated non-specific inflammatory changes.

Interestingly, a colleague in sexual health saw the patient, who had subsequently self-presented to their clinic. Examination revealed multiple mouth ulcers, as noted on OMFS examination, but also a single ulcerated mucosal abnormality with a rolled margin on his left oropharynx, in keeping with primary syphilitic chancre. Syphilis serology was positive and he was treated for primary syphilis with IM benzathine penicillin. The histopathology slides were then reviewed and demonstrated a lymphoplasmacytic infiltrate.

Given the history and examination findings the OMFS differential diagnosis was recurrent aphthous stomatitis. Tissue was therefore sent for histopathological investigation and PCR was not requested. In view of the definitive diagnosis, immunohistochemistry was requested and was positive for *Treponema pallidum*, which is a highly invasive spirochaete bacterium causing syphilis.¹

This demonstrates some of the issues which arise due to patient non-disclosure of important history. The patient disclosed to the sexual health clinic that he had a number of new unprotected male partners in recent months and had his own suspicions about syphilis but did not volunteer this during his OMFS appointment. It is usual for enquiries about sexual history to be made in the sexual health clinic but this may be an area outside of the 'comfort zone' in OMFS or general dental practice.

The sexual health consultant advised syphilis has had a rise again, particularly in male homosexual groups, but that there has also been a rise in heterosexual syphilis. In many cases, clinical manifestations of syphilis are highly characteristic and given there are oral manifestations, dentists are well placed to aid in diagnosis.

A. Cameron, Bath, UK

Reference

1. Radolf J D. *Treponema*. In: Baron S (ed). *Medical microbiology*. 4th edition. Chapter 36. Galveston (TX): University of Texas Medical Branch at Galveston, 1996. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK7716/> (accessed August 2020).

<https://doi.org/10.1038/s41415-020-2135-0>