



Correction: Global deletion of optineurin results in altered type I IFN signaling and abnormal bone remodeling in a model of Paget's disease

Sing-Wai Wong · Bo-Wen Huang · Xiangxiang Hu · Eui Ho Kim · Joseph P. Kolb · Ricardo J. Padilla · Peng Xue · Lufei Wang · Thomas H. Oguin III · Patricia A. Miguez · Henry C. Tseng · Ching-Chang Ko · Jennifer Martinez

Published online: 16 July 2020

© The Author(s), under exclusive licence to ADMC Associazione Differenziamento e Morte Cellulare 2020, corrected publication 2022

Correction to: *Cell Death & Differentiation*

<https://doi.org/10.1038/s41418-019-0341-6>

Following publication of this article, the authors noticed that there was an error in the legend for Fig. 3. The relevant and corrected section is provided below:

(D, G) *Optn*^{+/+} and *Optn*^{-/-} bone-marrow-derived precursors were cultured under osteoclastogenic conditions, and autophagic activity (d) and noncanonical NF-κB signaling (g) was assessed by western blot on days 0–3 post RANKL treatment from the sample protein lysates. In d, cells were probed for OPTN, LC3, p62, ATG5, ATG7, and β-actin. In g, the western blots were reprobbed for p100/p52

with the same OPTN and β-actin bands shown for reference. e, f *Optn*^{+/+} and *Optn*^{-/-} bone-marrow-derived precursors were cultured with RANKL for 0–60 min or 0–3 days, to assess the activation of canonical NF-κB signaling by western blot. Cells were probed for OPTN (e, f), phos-p65 (e, f), Total p65 (f), Total IκB (e), and β-actin (e, f).

In addition, in the 100 μm TRAP panel of Fig. 2b, an adjacent serial histologic section from the same sample was inadvertently used during image compilation. The correct image is provided below, the correct 100 μm TRAP image is from the same histologic section as that used for the 1 mm TRAP image.

The authors apologise for these errors.

Fig. 2




Fig. 3

