Oxidative stress in critically ill ventilated adults: effects of vitamin D₃ and associations with alveolar macrophage function

Jenny E. Han^{1,2} · Jessica A. Alvarez³ · Bashar Staitieh^{1,2} · Vin Tangpricha^{3,4} · Li Hao³ · Thomas R. Ziegler^{3,4} · Greg S. Martin^{1,2} · Lou Ann S. Brown⁵

Received: 28 March 2017 / Revised: 16 August 2017 / Accepted: 21 September 2017 / Published online: 29 December 2017
© Macmillan Publishers Limited, part of Springer Nature 2018

Abstract

Background/objectives Disruptions in redox balance lead to oxidative stress, a promoter of morbidity in critical illness. This study aimed to: (1) characterize the plasma and alveolar thiol/disulfide redox pools, (2) examine their associations with alveolar macrophage phagocytosis, and (3) determine the effect of high dose vitamin D₃ on plasma thiol/disulfide redox.

Subjects/methods Subjects were 30 critically ill, ventilated adults in a double-blind randomized trial of high-dose (250 000 or 500 000 IU) vitamin D₃ or placebo. Baseline bronchoalveolar lavage fluid (BALF) samples were analyzed for determination of alveolar phagocytosis index (PI) and for concentrations of glutathione (GSH), glutathione disulfide (GSSG), cysteine (Cys), cystine (CySS), and their respective redox potentials (EhGSSG and EhCySS). Plasma redox outcomes were assessed at baseline and days 7 and 14.

Results Baseline plasma Cys was inversely associated with alveolar PI ($\rho = -0.69$, $P = 0.003$), and E_hCySS was positively associated with PI ($\rho = 0.61$, $P = 0.01$). Over time, among all subjects there was an increase in plasma GSH levels and a decrease in E_hGSSG ($P < 0.01$ for both), with no difference by treatment group. Vitamin D₃ decreased oxidized plasma GSSG to a more normal state (P for group $\times$ time = 0.009).

Conclusions Oxidative stress indicators were positively associated with alveolar macrophage phagocytic function in acutely ill ventilated adults. High-dose vitamin D₃ decreased plasma GSSG concentrations, which suggests that vitamin D can possibly improve the oxidative stress environment.

Background

In critically ill patients, oxidative stress is a key mechanism of injury that results from sepsis and multi-system organ

failure and is associated with increased morbidity and mortality [1]. The balance in complex biochemical oxidation-reduction (redox) states can be assessed by measuring thiol/disulfide couples, glutathione (GSH) and glutathione disulfide (GSSG)—a major intracellular thiol redox system and cysteine (Cys) and cystine (CySS)—the predominant extracellular thiol redox system—(Fig. 1) within biofluids or tissues [2].

Normally the thiol redox systems are tightly controlled; however extracellular redox imbalance of thiol pairs can impact key cellular functions, such as proliferation, differentiation and apoptosis [2]. In acute critically ill pediatric patients, Grunwell et al [3] reported higher systemic oxidative stress compared to healthy controls, as indicated by a more oxidized plasma GSH/GSSG redox system. In adults with acute respiratory distress syndrome, GSH levels in the bronchoalveolar lavage fluid (BALF) have been reported to be depleted [4–6]. However, the thiol/disulfide redox systems within the plasma or alveolar space of adults with critical illness have not been well-characterized.

Jenny E. Han and Jessica A. Alvarez are Contributed equally to this work

✉ Jenny E. Han
jehan2@emory.edu

¹ Division of Pulmonary, Allergy, Critical Care and Sleep Medicine, Department of Medicine, Emory University, Atlanta, GA, USA

² Emory Critical Care Center, Emory University, Atlanta, GA, USA

³ Division of Endocrinology, Metabolism and Lipids, Department of Medicine, Emory University, Atlanta, GA, USA

⁴ Atlanta VA Medical Center, Decatur, GA, USA

⁵ Department of Pediatrics, Division of Neonatal-Perinatal Medicine, Emory University, Atlanta, GA, USA

Fig. 1 Glutathione and Cysteine Redox Equipoise. Both amino acids undergo reversible oxidation to form disulfides

The lung has an important role of providing effective barriers against oxidants and microorganisms, and GSH in extracellular lining fluid plays an important role in this host defense and are in high concentrations compared to other extracellular environments [2]. Macrophages, key mediators in the response to critical illness, are highly influenced by oxidative stress. Oxidative stress controls a number of important macrophage signaling pathways [6] and a wide range of functions, including protein translation, wound healing [7], apoptosis [8], and phagocytosis [9]. Liang et al have shown that GSH depletion and the resulting oxidative stress are associated with impaired alveolar macrophage phagocytosis, microbe clearance and increased risk for apoptosis [10, 11]. The relationship between systemic and alveolar oxidative stress and macrophage phagocytosis has not previously been investigated in critically ill ventilated patients.

Experimental and some clinical studies in non-ICU clinical populations suggest that vitamin D can reduce oxidative stress [11–16]. Approximately 60% of critically ill patients are vitamin D deficient [17–20]. We have previously shown that high-dose vitamin D₃ supplementation in ventilated, critically-ill patients' decreases hospital length of stay [21]. Amrein et al demonstrated a reduction in hospital mortality in a subgroup of patients with the highest degree of vitamin D deficiency [22]. Also vitamin D status has been shown to correlate with plasma redox biomarkers in both healthy subjects [23] and in a pediatric intensive care unit population [24]. Whether the improved outcome in our previous study with critically-ill patients treated with high-dose vitamin D₃ was linked to improved reduced/oxidized redox states remains to be determined.

The aims of this study were to (1) examine the relationship between the plasma and alveolar thiol/disulfide redox pools, (2) determine if previous studies identifying a link between oxidative stress and impaired alveolar macrophage phagocytosis is true for mechanically ventilated critically ill patients, and (3) evaluate the effect of high-dose vitamin D₃ on systemic oxidative stress. The primary and secondary clinical outcomes of this pilot, randomized control trial (e.g., plasma 25(OH) D concentrations, hospital length of stay, and mortality) have been previously published [21]. Here, we discuss secondary translational

outcomes to provide more mechanistic data to supplement the clinical outcomes.

Methods

Study design and participants

This was a pilot double blind randomized control trial and was approved by Emory University Institutional Review Board and registered at www.clinicaltrials.gov (NCT01372995). The enrollment goal was 36 patients from two Atlanta, Georgia hospitals; Emory University Hospital (EUH) and Emory University Midtown (EUH-M). The study was conducted from July 2011–March 2014 and enrolled critically ill patients expected to remain ventilated for at least 72 h after entry. Full details of trial design, inclusion and exclusion criteria, safety criteria and other methodological details are provided in the previous publication [21].

Participant selection

In summary, major inclusion criteria were (1) age > 18 years; (2) respiratory failure requiring mechanical ventilation for at least 72 h after study entry; and (3) anticipated stay in the ICU for at least 96 h after entry. Major exclusion criteria were (1) use of high-dose vitamin D₃ supplementation ($\geq 50,000$ IU a week) to treat vitamin D deficiency within the prior 6 months; (2) history of medical disorders associated with hypercalcemia, chronic renal failure requiring dialysis, cirrhosis or HIV infection; and (3) hypercalcemia (albumin-corrected serum calcium > 10.8 mg/dL or ionized calcium > 5.2 mg/dL) [21].

Intervention

Following informed consent, patients were assigned to either a daily dose of oral 50 000 IU vitamin D₃ for 5 days [total of 250 000 IU] or daily dose of 100 000 IU vitamin D₃ for 5 days [total 500 000 IU] or placebo in a 1:1:1 ratio using a blinded block randomization schedule, overseen by biostatisticians of the Atlanta Clinical and Translational Science Institute (ACTSI) biostatistics core [21]. Randomization was stratified based on clinical site and APACHE II score > 15 or ≤ 15 . The medications were dissolved in sterile water and administered through an enteral feeding tube. Cholecalciferol 50 000 IU tablets were manufactured from Tischon (Westbury, NY) and Biotech (Fayetteville, AR) [21]. EUH and EUH-M Investigational Drug. With the exception of the pharmacists, all study staff were blinded to the group allocation. Full details of trial design, participant selection and intervention were published previously [21].

Data collection and laboratory analysis

Blood sampling

Twenty milliliter of venous blood was collected at baseline and every 7 days while the subject remained in the hospital.

Total 25(OH)D

Plasma total 25(OH)D was measured using a chemiluminescent-based automated machine (IDS-iSYS; Immunodiagnostic Systems, Scottsdale, AZ). The Emory Vitamin D laboratory participates in the Vitamin D External Quality Assessment Scheme and NIST/NIH Vitamin D Metabolites Quality Assurance Program. Insufficient vitamin D status was defined as plasma total 25(OH)D concentrations < 30 ng/mL, with values ≥ 30 ng/mL considered sufficient.

BALF collection: A bronchoalveolar lavage was performed on study day 0 and day 7, if subjects were still intubated. If the subject was expected to be taken off the ventilator before day 7, the sample could be collected as early as day 5. The BALF procedure was performed by serial instillation of 30 mL aliquots of normal saline into a pulmonary sub segment and BALF was collected by suction.

Alveolar macrophage phagocytosis index and percent positive analysis: BALF was centrifuged at 1200 rpm for 7 min for recovery of cell pellet. Manual cell counts were performed with a hemocytometer and differentials were obtained from 300 consecutive cells after Diff-Quik staining (Andwin Scientific, Addison, IL). The cell pellet was resuspended in 10 mL of 1:1 Dulbecco's modified Eagle's medium/Ham's F-12 solution containing 2% fetal bovine serum, L-glutamine, 15 mmol/L HEPES, penicillin (10,000 U), streptomycin (10,000 mg/mL), amphotericin (25 mg/mL), and gentamicin (4 μ g/mL). Alveolar macrophages (100,000 cells) were added to glass-chamber slides containing 100 μ L of medium and 20 μ L of phosphate-buffered saline. Alveolar macrophages were incubated at 37°C with 10% CO₂ for 15 h, after which 10×10^5 particles of pH-sensitive pHrodo fluorescein isothiocyanate-conjugated inactivated *Staphylococcus aureus* were added (10:1 ratio of *S. aureus*/alveolar macrophages) to the cell cultures. After a 2 h incubation, the cells were washed, fixed with 4% paraformaldehyde, and stored at 4°C until analysis. Fluorescence of phagocytized *S. aureus* was determined by quantitative computer analysis of images taken with a Zeiss inverted microscope using 10 fields per experimental condition with the same pinhole, detector gain, and amplifier offset. The basis of this assay is that *S. aureus* is internalized and the pHrodo component fluoresces once it reaches the lower pH (pH = 4) present in the

phagolysosome. Macrophages with any internalized bacteria were considered positive for phagocytosis. Phagocytosis was quantified by phagocytic index, which was calculated as the percentage of cells positive for phagocytosis (percent positive) multiplied by the mean relative fluorescence units of *S. aureus* per cell [25].

Determination of plasma and BALF oxidative stress: Details for redox determination have been previously described [26]. Briefly, plasma and BALF samples were transferred to a microcentrifuge tube containing a preservative solution consisting of borate buffer, γ -glutamyl-glutamate (internal standard), serine, heparin, bathophenanthroline disulfonate, and iodoacetic acid. Samples were centrifuged, the supernatant transferred to a second solution containing perchloric acid and boric acid, and immediately frozen at -80°C . For analysis, samples were thawed, protein precipitated, and supernatant treated with a dansyl chloride solution for derivatization. Plasma and BALF samples were subsequently analyzed using high-performance liquid chromatography (Waters 2690, Waters Corp., Milford, MA) with fluorescence detection (Waters 474 and Gilson 121 detectors). Plasma and BALF GSH, GSSG, Cys, and CySS were quantified based on the integral areas of the peak for each respective analyte relative to the integral area of the internal standard. A dilution factor for urea ($[\text{urea}]_{\text{plasma}} / [\text{urea}]_{\text{BALF}}$) was applied to BALF redox quantifications [26]. The redox potential for the thiol pairs (E_h GSH/GSSG, E_h Cys/CySS) was calculated using the Nernst equation ([expressed in millivolts (mV)]); redox potential data for the two redox pools in BALF were corrected for pH. In this system, a less negative E_h redox potential value indicates a more oxidized state [26]. The percentage of CySS in relation to the total Cys pool [% CySS = (CySS / (Cys + CySS)) * 100] and percentage of GSSG in relation to the total GSH pool [% GSSG = (GSSG / (GSH + GSSG))] were also calculated.

Statistical analysis

Descriptive statistics were conducted on demographic data and study outcomes. Data are reported as mean $\pm$ SD for normally distributed variables and median (IQR) for non-normally distributed variables. Variables deviating from a normal distribution were log-transformed for analyses, as necessary, or non-parametric methods were used. Baseline group differences (treatment group or vitamin D status group) were assessed with a *t*-test or Wilcoxon test. Paired *t*-tests were used to compare plasma and BALF redox status in subjects with both measurements available. Correlation analyses were assessed using a Pearson or Spearman method. Repeated measures were assessed with a linear mixed-effect method with Tukey's post-hoc analyses. Study outcomes did not differ between the two vitamin D groups

Table 1 Baseline demographic and plasma redox variables by treatment group

Variable	All (n = 30)*	Placebo (n = 10)	Vitamin D (n = 20)	P
Age (yr)	63.5 ± 17.3	64.8 ± 17.5	64.0 ± 17.7	0.87
Male		6 (60)	13 (65)	0.78
African American	12 (40)	4 (40)	10 (50)	0.10
BMI (kg/m ²)	30.6 ± 7.59	28.2 ± 9.9	31.5 ± 6.4	0.39
APACHE II	21 ± 9	23.2 ± 8.8	20 ± 9	0.37
Infection on Admission	13 (43)	6 (60)	7 (35)	0.17
Plasma 25(OH)D (ng/mL)	21.4 ± 9.1	21.5 ± 12.2	21.1 ± 7.9	0.75
25(OH)D < 20 ng/mL	13 (43)	5 (50)	8 (40)	0.59
25(OH)D < 30 ng/mL	25 (83)	8 (80)	17 (85)	0.73
Cys (μM)	1.11 (0.60, 2.61)	1.36 (0.69, 8.34)	0.98 (0.57, 1.80)	0.40
CySS (μM)	7.05 (4.60, 9.07)	6.62 (4.26, 12.34)	7.25 (4.24, 8.87)	0.77
%CySS	83.42 (73.70, 93.20)	82.32 (50.42, 94.03)	85.74 (72.47, 90.16)	0.88
E _h CySS (mV)	−51.16 ± 31.21	−54.79 ± 46.39	−49.34 ± 21.59	0.68
GSH (μM)	0.21 (0.07, 0.35)	0.21 (0.08, 0.72)	0.21 (0.07, 0.32)	0.63
GSSG (μM)	0.04 (0.008, 0.24)	0.04 (0.007, 0.22)	0.09 (0.009, 0.30)	0.46
%GSSG	34.18 (3.19, 66.05)	11.53 (2.00, 59.90)	40.31 (7.84, 67.63)	0.37
E _h GSSG (mV)	−75.71 ± 40.94	−88.20 ± 42.13	−69.47 ± 40.05	0.27

Data reported as *N* (%), mean ± SD for normal distributions, or median (IQR) for non-normal distributions

**N* = 27 for plasma redox outcomes (*n* = 9 for placebo group, *n* = 18 for vitamin D group)

Table 2 Comparison of baseline plasma and alveolar oxidative stress*

g	Plasma	BALF	P-value
Cys (μM)	0.94 (0.66, 2.95)	0.06 (0.01, 0.11)	0.05 (0.01, 0.21)
< 0.001	CySS (μM)	7.05 (3.86, 10.12)	
< 0.001			
%CySS	83.42 (60.68, 93.21)	56.03 (33.44, 62.21)	0.007
E _h CySS (mV)	−42.77 (−77.85, −34.00)	−23.96 (−38.38, −8.05)	0.01
GSH (μM)	0.26 (0.08, 0.51)	0.08 (0.02, 0.19)	0.22
GSSG (μM)	0.13 (0.02, 0.64)	0.03 (0.02, 0.05)	0.006
%GSSG	60.89 (3.76, 77.73)	37.07 (20.30, 69.58)	0.85
E _h GSSG (mV)	−62.80 (−116.44, −32.99)	−39.07 (−70.05, −13.76)	0.10

**N* = 17 subjects with paired plasma and bronchoalveolar lavage fluid (BALF) samples. Bold signifies statistical significance

(250,000 IU vs 500,000 IU); therefore, these two groups were merged for reported analyses. This is secondary analysis from our previously published study [21] in which the primary outcome was the change in plasma 25(OH)D from baseline to 7 days; 12 subjects in each group were required to test the hypothesis that high-dose vitamin D₃ would achieve plasma 25(OH)D levels ≥ 30 ng/mL with power of 0.94 and α of 0.05. Data were analyzed using JMP

statistical software (version 12, SAS Institute, Cary, NC). All analyses were two-sided with a significance level of α = 0.05.

Results

The study design, CONSORT diagram, enrollment and randomization are reported in a previous publication [21]. Baseline demographic data by treatment group is shown in Table 1.

Baseline systemic and alveolar oxidative stress

Baseline systemic and alveolar macrophage oxidative stress data is shown in Table 2. A comparison of baseline plasma thiol/disulfide redox measures revealed significantly higher Cys, CySS, %CYSS and GSSG and more reduced E_hCYSS compared to BALF-derived redox measures.

Correlation of systemic oxidative stress and phagocytosis index (PI) and percent of cells positive for phagocytosis (percent positive)

Correlation analyses of plasma baseline redox measures against phagocytosis outcomes are shown in Table 3. Plasma Cys was inversely associated with both PI (ρ = −0.69, p = 0.003) and percent positive (ρ = −0.51, p = 0.04). Both plasma %CySS and E_hCySS were strongly

Table 3 Spearman correlations between systemic oxidation stress and alveolar macrophage phagocytosis

	Phagocytosis Index	Percent of cells phagocytosis-positive
Plasma Cys	−0.69 (0.003)	−0.51 (0.04)
Plasma CySS	0.23 (0.39)	0.16 (0.55)
Plasma % CySS	0.56 (0.02)	0.44 (0.09)
Plasma E _h CySS	0.61 (0.01)	0.46 (0.07)
Plasma GSH	−0.06 (0.81)	−0.30 (0.25)
Plasma GSSG	−0.26 (0.32)	−0.39 (0.13)
Plasma % GSSG	−0.06 (0.83)	−0.13 (0.63)
Plasma E _h GSSG	0.13 (0.64)	0.11 (0.70)

Spearman Correlations. $N = 17$. Bold signifies statistical significance

positively correlated with PI ($\rho = 0.56$ and 0.61 , respectively; $p \leq 0.02$ for both). There were no significant correlations between the alveolar Cys/CySS or GSH/GSSG redox indexes and the macrophage phagocytosis indexes.

Effect of vitamin D on systemic oxidative stress

There were no differences in oxidative stress indices at baseline between placebo and vitamin D-treated subjects (data not shown). Over time, both groups demonstrated an increase in plasma GSH levels (P for time = 0.001) but the difference between treatment groups and treatment-by-time effect were not statistically significant (Fig. 2a). Plasma GSSG increased in the placebo group, while there was a decrease in plasma GSSG in the vitamin D group (P for treatment $\times$ time = 0.009 , Fig. 2b). Plasma E_hGSSG significantly decreased (i.e., became more reduced) over time (P for time = 0.009), although there was no significance between treatment group effect or treatment-by-time effect (Fig. 2c). There were no statistically significant changes in the plasma Cys/CySS redox system (P for time, group, and group $\times$ time > 0.05 for all).

Discussion

GSH and GSSG represent a major intracellular thiol/disulfide redox system, while Cys and CySS represent a major extracellular redox system [27]. Assessment of these redox systems can provide useful data on oxidative stress and redox balance within critically ill patient populations. In this pilot study of mechanically-ventilated adults in the ICU, we showed that oxidative stress, as assessed by thiol/disulfide redox systems, is higher in the alveolar space compared to

the plasma and is associated with phagocytosis. Furthermore, reductions in plasma oxidative stress are apparent over time and can be influenced by high-dose vitamin D₃ supplementation.

We evaluated the baseline thiol/disulfide redox environment in both the plasma and the lung of adult ICU patients. The lower E_hCySS within the plasma compartment indicates a more reduced (less oxidized) environment relative to the alveolar compartment. A previous study showed that, in patients with acute respiratory distress syndrome the alveolar environment is more oxidized than healthy controls [4]. The alveolar space in our patient population is far more oxidized compared to BALF of previous critically ill patients [4, 5]. One possible explanation is that mechanical ventilation contributed to a more oxidized environment in the alveolar space. A reduction in oxidative stress during the course of recovery in critically-ill populations is expected. However, the novel evaluation of plasma thiol/disulfide redox measures over time in our population showed a significant increase in plasma GSH and a more reduced E_hGSSG, with no changes in the plasma Cys/CySS redox system. Our data are, therefore, consistent with pediatric cross-sectional data by Grunwell [3] indicating that plasma Cys/CySS redox pools are preserved in critical illness and possibly maintained by changes in the supply of GSH from tissue to plasma.

We further investigated the relationship between macrophage phagocytosis function and systemic and alveolar oxidative stress. Experiments with alcohol-fed rodents by Liang et al [11] showed that GSH depletion within alveolar macrophages and related chronic oxidative stress impairs alveolar macrophage function. We found that alveolar macrophage phagocytosis of *S. aureus* was strongly positively correlated with the plasma Cys/CySS redox potential (E_hCySS) and %CySS, both indicative of higher oxidative stress. This data suggest that the extracellular redox environment is closely linked to innate immune functions of alveolar macrophages. It is possible that increased phagocytosis may create more reactive oxygen species and apoptosis resulting in a more oxidized environment. Increased oxidation may lead to further dysregulation of immune function contributing to more deleterious clinical effects over time in this patient population. Alternatively, a more oxidized plasma Cys/CySS redox pool may be a homeostatic response to critical illness required to promote phagocytosis by macrophages [9]. Additional detailed studies are required to establish causal relationships between the plasma redox environment and alveolar macrophage function in critical illness. Furthermore, additional studies are needed to determine if phagocytosis is upregulated in the resident alveolar macrophage pool or reflective of macrophages newly recruited from the oxidized systemic environment.

Cross-sectional studies by members of our group have previously demonstrated significant associations between vitamin D status and plasma thiol/disulfide redox biomarkers in both a generally healthy adult population [23] and in critically ill pediatric patients [24]. In healthy adults, plasma 25(OH)D was positively associated with plasma GSH and inversely associated with $E_h\text{GSSG}$ [23]; whereas in critically ill children, a more sufficient vitamin D status

Fig. 2 Plasma GSH/GSSG redox outcomes over time in critically ill adults. Data are shown as least squares (LS) mean $\pm$ SEM on back-transformed data. Data were analyzed using linear mixed-effect models with Tukey's post-hoc analyses. Points not connected by the same letter indicate statistically significant differences. $N = 20$ on day 0 and 7, $n = 15$ on day 14. Figure 2a Plasma GSH concentrations. The time effect was statistically significant ($P = 0.001$). The treatment and treatment-by-time effects were not statistically significant ($P = 0.22$ and 0.98 , respectively). Figure 2b Plasma GSSG concentrations. The treatment-by-time effect was statistically significant ($P = 0.009$). The treatment and time effects were not statistically significant ($P = 0.18$ and 0.74 , respectively). Figure 2c Plasma $E_h\text{GSSG}$. The time effect was statistically significant ($P = 0.009$). The treatment and treatment-by-time effects were not statistically significant ($P = 0.18$ and 0.74 , respectively)

was associated with lower plasma GSH and GSSG and a more reduced plasma $E_h\text{CySS}$ [24]. In the current study in mechanically ventilated adults, we have measured, for the first time, the effect of a high-dose vitamin D₃ intervention on plasma thiol/disulfide redox. Our data show that vitamin D₃ supplementation decreases GSSG, the oxidized disulfide form of GSH, relative to the placebo group, although there were no effects on other measured redox outcomes. Experiments performed in vitro support our clinical finding in that vitamin D, in its active hormonal form (calcitriol), upregulates glutathione reductase, which catalyzes the reduction of GSSG to GSH [28]. Administration of vitamin D₃ in patients with type 2 diabetes and in vitamin D insufficient women revealed an improvement in oxidative stress as measured by increased GSH, total antioxidant capacity, and malondialdehyde [15, 16]. Our primary study found a decrease in hospital length of stay [21]; this secondary analysis provides a potential mechanism by which vitamin D may impact clinical outcomes.

The major strength of this study was the use of a randomized, placebo-controlled design in a well-characterized cohort. A limitation of this pilot study was the small sample size, particularly in the number of subjects undergoing bronchoscopy. The primary outcome of our primary study was change in plasma total 25(OH)D [21], therefore this study may not have been adequately powered to detect changes in the secondary outcomes reported here. Despite our small sample size, we were able to determine that systemic oxidative stress improved over time with high-dose vitamin D₃ administration. Additional markers of oxidative stress or antioxidants were not measured. An additional limitation is that we only assessed macrophage phagocytosis with *S. aureus* and no other pathogens, which could have different responses. We chose a bacterial model since that is still the most common pathogen in the lungs of hospitalized critically ill patients. Finally, many subjects were liberated from the ventilator before repeat bronchoscopy on day 7, thus limiting our ability to conduct longitudinal BALF analyses.

Conclusion

We showed that a more oxidized plasma Cys/CySS redox pool was associated with alveolar macrophage phagocytic function, while systemic GSH-related oxidative stress decreased during clinical recovery. Finally, in a randomized, placebo-controlled trial, high-dose vitamin D₃ decreased plasma GSSG concentrations, which suggests that vitamin D can possibly improve the oxidative stress environment, which may provide potential mechanism by which vitamin D may impact clinical outcomes.

Acknowledgements We would like to acknowledge Mona Brown, Guatam Hebbbar, Jennifer Jones, Frank Harris, and Janine Ward for their contribution to the study.

Funding This project was supported, in part, by National Institutes of Health grants: NIH R21 HL110044 (GSM, TRZ, LAB, VT, JEH), K24 DK096574 (TRZ), UL1 TR000454 (JEH, GSM, TRZ, VT), K01 DK102851 (JAA), T32 HL076118 (BS), and T32 AA013528 (JEH).

Conflict of interest The authors declare that they have no conflict of interest.

References

1. Gunaydin B, Sancak B, Candan S, Sariahmetoglu M, Ozcagli G, Tunctan B, et al. Temporal variation of oxidant stress in critically ill patients. *Minerva Anesthesiol.* 2007;73(5):261–6.
2. Moriarty-Craige SE, Jones DP. Extracellular thiols and thiol/disulfide redox in metabolism. *Annu Rev Nutr.* 2004;24:481–509.
3. Grunwell JR, Gillespie SE, Ward JM, Fitzpatrick AM, Brown LA, Gauthier TW, et al. Comparison of glutathione, cysteine, and their redox potentials in the plasma of critically ill and healthy children. *Front Pediatr.* 2015;3:46.
4. Bunnell E, Pacht ER. Oxidized glutathione is increased in the alveolar fluid of patients with the adult respiratory distress syndrome. *Am Rev Respir Dis.* 1993;148(5):1174–8.
5. Pacht ER, Timmerman AP, Lykens MG, Merola AJ. Deficiency of alveolar fluid glutathione in patients with sepsis and the adult respiratory distress syndrome. *Chest.* 1991;100(5):1397–403.
6. Brune B, Dehne N, Grossmann N, Jung M, Namgaladze D, Schmid T, et al. Redox control of inflammation in macrophages. *Antioxid Redox Signal.* 2013;19(6):595–637.
7. Metukuri MR, Namas R, Gladstone C, Clermont T, Jefferson B, Barclay D, et al. Activation of latent transforming growth factor-beta1 by nitric oxide in macrophages: role of soluble guanylate cyclase and MAP kinases. Wound repair and regeneration: official publication of the Wound Healing Society [and] the European Tissue Repair Society. 2009;17(4):578–88.
8. Heinloth A, Brune B, Fischer B, Galle J. Nitric oxide prevents oxidized LDL-induced p53 accumulation, cytochrome c translocation, and apoptosis in macrophages via guanylate cyclase stimulation. *Atherosclerosis.* 2002;162(1):93–101.
9. Duan J, Kodali VK, Gaffrey MJ, Guo J, Chu RK, Camp DG, et al. Quantitative profiling of protein S-glutathionylation reveals redox-dependent regulation of macrophage function during nanoparticle-induced oxidative stress. *ACS Nano.* 2016;10(1):524–38.
10. Liang Y, Yeligar SM, Brown LA. Chronic-alcohol-abuse-induced oxidative stress in the development of acute respiratory distress syndrome. *Sci World J.* 2012;2012:740308.
11. Liang Y, Harris FL, Brown LA. Alcohol induced mitochondrial oxidative stress and alveolar macrophage dysfunction. *Bio Med Res Int.* 2014;2014:371593.
12. Ke CY, Yang FL, Wu WT, Chung CH, Lee RP, Yang WT, et al. Vitamin D₃ reduces tissue damage and oxidative stress caused by exhaustive exercise. *Int J Med Sci.* 2016;13(2):147–53.
13. Sales de Almeida JP, Liberatti LS, Nascimento Barros FE, Kallaur AP, Batisti Lozovoy MA, Scavuzzi BM, et al. Profile of oxidative stress markers is dependent on vitamin D levels in patients with chronic hepatitis C. *Nutrition.* 2016;32(3):362–7.
14. Kanikarla-Marie P, Jain SK. 1,25(OH)₂D₃ inhibits oxidative stress and monocyte adhesion by mediating the upregulation of GCLC and GSH in endothelial cells treated with acetoacetate (ketosis). *J Steroid Biochem Mol Biol.* 2016;159:94–101.
15. de Medeiros Cavalcante IG, Silva AS, Costa MJ, Persuhn DC, Issa CI, de Luna Freire TL, et al. Effect of vitamin D₃ supplementation and influence of BsmI polymorphism of the VDR gene of the inflammatory profile and oxidative stress in elderly women with vitamin D insufficiency: Vitamin D₃ megadose reduces inflammatory markers. *Exp Gerontol.* 2015;66:10–6.
16. Shab-Bidar S, Neyestani TR, Djazayeri A. The interactive effect of improvement of vitamin D status and VDR FokI variants on oxidative stress in type 2 diabetic subjects: a randomized controlled trial. *Eur J Clin Nutr.* 2015;69(2):216–22.
17. Jeng L, Yamshchikov AV, Judd SE, Blumberg HM, Martin GS, Ziegler TR, et al. Alterations in vitamin D status and antimicrobial peptide levels in patients in the intensive care unit with sepsis. *J Transl Med.* 2009;7:28.
18. Mata-Granados JM, Vargas-Vassero J, Ferreira-Vera C, Luque de Castro MD, Pavon RG, Quesada Gomez JM. Evaluation of vitamin D endocrine system (VDES) status and response to treatment of patients in intensive care units (ICUs) using an on-line SPE-LC-MS/MS method. *J Steroid Biochem Mol Biol.* 2010;121(1–2):452–5.
19. Higgins DM, Wischmeyer PE, Queensland KM, Sillau SH, Sufit AJ, Heyland DK. Relationship of vitamin D deficiency to clinical outcomes in critically ill patients. *J Parenter Enter Nutr.* 2012;36(6):713–20.
20. Nair P, Lee P, Reynolds C, Nguyen ND, Myburgh J, Eisman JA, et al. Significant perturbation of vitamin D-parathyroid-calcium axis and adverse clinical outcomes in critically ill patients. *Intensive Care Med.* 2013;39(2):267–74.
21. Han JE, Jones JL, Tangpricha V, Brown MA, Brown LA, Hao L, et al. High dose vitamin D administration in ventilated intensive care unit patients: A pilot double blind randomized controlled trial. *J Clin Transl Endocrinol.* 2016;4:59–65.
22. Amrein K, Schnedl C, Holl A, Riedl R, Christopher KB, Pachler C, et al. Effect of high-dose vitamin D₃ on hospital length of stay in critically ill patients with vitamin D deficiency: the VITdAL-ICU randomized clinical trial. *JAMA.* 2014;312(15):1520–30.
23. Alvarez JA, Chowdhury R, Jones DP, Martin GS, Brigham KL, Binongo JN, et al. Vitamin D status is independently associated with plasma glutathione and cysteine thiol/disulfide redox status in adults. *Clin Endocrinol.* 2014;81(3):458–66.
24. Alvarez JA, Grunwell JR, Gillespie SE, Tangpricha V, Hebbbar KB. Vitamin D deficiency is associated with an oxidized plasma cysteine redox potential in critically ill children. *J Steroid Biochem Mol Biol.* 2018;175:164–9.
25. Gauthier TW, Ping XD, Harris FL, Wong M, Elbahesh H, Brown LA. Fetal alcohol exposure impairs alveolar macrophage function via decreased glutathione availability. *Pediatr Res.* 2005;57(1):76–81.
26. Yeh MY, Burnham EL, Moss M, Brown LA. Chronic alcoholism alters systemic and pulmonary glutathione redox status. *Am J Respir Crit Care Med.* 2007;176(3):270–6.

27. Jones DP, Liang Y. Measuring the poise of thiol/disulfide couples in vivo. *Free Radic Biol Med.* 2009;47(10):1329–38.
28. Jain SK, Micinski D. Vitamin D upregulates glutamate cysteine ligase and glutathione reductase, and GSH formation, and decreases ROS and MCP-1 and IL-8 secretion in high-glucose exposed U937 monocytes. *Biochem Biophys Res Commun.* 2013;437(1):7–11.