

COMMENT



Genetic counselling and testing for neurodegenerative disorders using a proposed standard of practice for ALS/MND: diagnostic testing comes first

Ashley Crook ^{1,2}✉ and Alison McEwen¹

© The Author(s), under exclusive licence to European Society of Human Genetics 2021

European Journal of Human Genetics (2022) 30:394–395; <https://doi.org/10.1038/s41431-021-01036-z>

Amyotrophic lateral sclerosis (ALS, commonly referred to as motor neurone disease (MND) in some countries) is a neurodegenerative disorder with an incidence of ~2.31 per 100,000 in Europe [1]. About 20% of cases are caused by a pathogenic variant in one of several causative genes [2], and family history cannot be relied upon to confirm the presence of all pathogenic variants [1, 3]. This finding, along with the emergence of genotype-driven antisense oligonucleotide (ASO) therapies, is driving increased interest in diagnostic genetic testing for ALS/MND and associated frontotemporal dementia (FTD). Consequently, access to genetic counselling and testing is now considered a fundamental right of all people diagnosed with ALS/MND [4]. While the initial diagnostic testing may be facilitated by specialists involved in the diagnosis and care of people with neurodegenerative disorders, such as neurologists [5], ensuring adequate exploration of the implications for other members of the family is also important. Genetic counsellors (where available) are well placed to discuss and explore the familial implications of testing as part of the multi-disciplinary team [6, 7]. As genetics and genomics is increasingly implemented into routine clinical care [8], a standard of practice for genetic counselling and testing for ALS/MND/FTD offers a model that may also be relevant for genetic counselling and testing for other neurodegenerative disorders.

Diagnostic genetic testing for ALS/MND/FTD can occur during a period of significant stress, coinciding with the disease diagnosis and/or progression. For some patients and their carers, the genetic counselling that accompanies the testing may form a minor part of the overall care. For others, the context within which genetic testing is offered forms a significant aspect of their care. The confirmation of a causative pathogenic variant may impact both the patient and their family members as they adjust to the diagnosis and the implications for family members who are at risk of carrying the same genetic variant [9–11]. The potential for increased uptake of diagnostic genetic testing if treatment trials are successful, along with changing views on patient confidentiality, the potential for non-disclosure to at-risk family members, fears of discrimination or stigmatisation, and associated legal implications [11, 12] all contribute to the need for a standard of practice for diagnostic genetic counselling and testing for ALS/MND/FTD.

Realising the potential options and benefits that arise following the identification of a causative pathogenic variant in an individual relies on this information being shared with at-risk family members. While diagnostic genetic counselling and testing may focus on the implications of the testing for the patient, the potential impact of the results for family members and the need for family communication should be explored [9, 10]. Family communication is complex, with many studies across a diverse range of speciality areas demonstrating that ongoing support may be required to ensure information reaches the at-risk members of the extended family [13–15]. Genetic counselling support improves communication of genetic information in some families [16], so developing a standard of care that embeds ongoing support and facilitates appropriate onward referral when needed, is crucial if we are to realise the familial benefits of diagnostic genetic counselling and testing.

Genetic counsellors are allied health professionals with post-graduate training in genetics and genomics, counselling and communication. Genetic counsellors are registered healthcare professionals in a number of countries [7, 17]. They are working in specialist multi-disciplinary teams, including in neurology, in growing numbers. When delivered by a genetic counsellor, genetic counselling integrates interpretation of family and medical history, education about the disorder, counselling to promote informed choice and support to facilitate adaptation [18]. Exploring and supporting family communication is central to genetic counselling conducted by genetic counsellors. Recent research has found that communication of genetic information to family members may not be adequately discussed when consenting patients to genomic sequencing, with only one-sixth of consent forms reviewed specifically addressing the communication of genetic information with family members, despite statements on consent forms noting that genetics is familial [19]. Our work has demonstrated that when genetic counselling is conducted by other healthcare professionals in a diagnostic context, insufficient time may be available to adequately engage patients and carers in a discussion about family communication. Following identification of a causative variant, healthcare professionals and patients and family members may be unsure or unaware of the availability of referral to genetic counselling,

¹Graduate School of Health, University of Technology Sydney, Chippendale, NSW, Australia. ²Centre for MND research and Department of Clinical Medicine, Faculty of Medicine, Health and Human Sciences, Macquarie University, Sydney, NSW, Australia. ✉email: Ashley.K.Crook@student.uts.edu.au

Received: 16 December 2021 Accepted: 20 December 2021

Published online: 5 January 2022

further hindering family communication and access to predictive and/or reproductive genetic counselling and testing for at-risk family members.

As genomics is integrated into routine clinical care, standards of practice that guide all healthcare professionals involved in genetic counselling and testing must be developed then evaluated. One such standard of practice is being developed by our team focusing on ALS/MND/FTD, and may inform other areas of neurogenetics. A standard of practice may include items for pre- and post-test discussion, including possible results, implications for the patient and their family and a plan to support family communication and adaptation to the result. Alongside a standard of practice, the development of online tools and resources to support informed consent are needed [10, 20]. Education and support for the healthcare professionals who conduct diagnostic genetic counselling and testing is also required [10]. Genetic counsellors are equipped with the training and experience to work alongside healthcare professionals in multi-disciplinary teams, providing education, accepting referrals for predictive and reproductive genetic testing, and discussing and exploring communication within the family. Embedding genetic counsellors in the multi-disciplinary care of clients and families whenever possible improves outcomes for patients and families by supporting family communication and adaptation, and contributes to the realisation of the benefits of integrating genomics into clinical care.

REFERENCES

1. Brown CA, Lally C, Kupelian V, Flanders WD. Estimated prevalence and incidence of amyotrophic lateral sclerosis and SOD1 and C9orf72 genetic variants. *Neuroepidemiology*. 2021. <https://doi.org/10.1159/000516752>.
2. Shephard SR, Parker MD, Cooper-Knock J, Verber NS, Tuddenham L, Heath P, et al. Value of systematic genetic screening of patients with amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry*. 2021;92:510–8.
3. Müller K, Oh K-W, Nordin A, Panthi S, Kim SH, Nordin F, et al. De novo mutations in SOD1 are a cause of ALS. *J Neurol Neurosurg Psychiatry*. 2021:jnnp-2021-327520. <https://doi.org/10.1136/jnnp-2021-327520>.
4. International Alliance of ALS/MND Associations T. Fundamental rights for people living with ALS/MND 2021 [updated April 2021]. <https://www.als-mnd.org/support-for-pals-cals/advocacy/pals-and-cals-rights/>.
5. Crook A, Jacobs C, Newton-John T, O'Shea R, McEwen A. Genetic counseling and testing practices for late-onset neurodegenerative disease: a systematic review. *J Neurol*. 2021. <https://doi.org/10.1007/s00415-021-10461-5>.
6. Salmon K, Anoja N, Breiner A, Chum M, Dionne A, Dupré N, et al. Genetic testing for amyotrophic lateral sclerosis in Canada—an assessment of current practices. *Amyotroph Lateral Scler Frontotemporal Degener*. 2021:1–8. <https://doi.org/10.1080/21678421.2021.1980890>.
7. Abacan M, Alsubaie L, Barlow-Stewart K, Caanen B, Cordier C, Courtney E, et al. The global state of the genetic counseling profession. *Eur J Hum Genet*. 2019;27:183–97.
8. Gaille M, Horn R, Bale M, Bourgain C, Buchanan J, Cambon-Thomsen A, et al. The ethics of genomic medicine: redefining values and norms in the UK and France. *Eur J Hum Genet*. 2021;29:780–8.

9. Howard J, Mazanderani F, Locock L. Life 'on high alert': how do people with a family history of motor neurone disease make sense of genetic risk? insights from an online forum. *Health Risk Soc*. 2021:1–17. <https://doi.org/10.1080/13698575.2021.1946488>.
10. Crook A, Jacobs C, Newton-John T, Richardson E, McEwen A. Patient and relative experiences and decision-making about genetic testing and counseling for familial ALS and FTD: a systematic scoping review. *Alzheimer Dis Assoc Disord*. 2021;35:374–85.
11. Wauters A, Van Hoyweghen I. Normalising life at risk of Huntington's disease. A qualitative study of backgrounds and coping strategies of fears of genetic discrimination. *Eur J Hum Genet*. 2021;29:940–8.
12. Middleton A, Patch C, Roberts J, Milne R, Costa A, Roberts L, et al. Professional duties are now considered legal duties of care within genomic medicine. *Eur J Hum Genet*. 2020;28:1301–4.
13. Oliveira CR, Mendes A, Sequeiros J, Sousa L. Management of information within Portuguese families with Huntington disease: a transgenerational process for putting the puzzle together. *Eur J Hum Genet*. 2020;28:1210–7.
14. Srinivasan S, Won NY, Dotson WD, Wright ST, Roberts MC. Barriers and facilitators for cascade testing in genetic conditions: a systematic review. *Eur J Hum Genet*. 2020;28:1631–44.
15. Pierron L, Hennessy J, Tezenas du Montcel S, Coarelli G, Heinzmann A, Schaerer E, et al. Informing about genetic risk in families with Huntington disease: comparison of attitudes across two decades. *Eur J Hum Genet*. 2021;29:672–9.
16. Forrest LE, Burke J, Bacic S, Amor DJ. Increased genetic counseling support improves communication of genetic information in families. *Genet Med*. 2008;10:167–72.
17. Hoskins C, Gaff C, McEwen A, Macciocia I, Pearn A, Shalhoub C, et al. Professional regulation for Australasian genetic counselors. *J Genet Couns*. 2021;30:361–9.
18. Resta R, Biesecker BB, Bennett RL, Blum S, Estabrooks Hahn S, Strecker MN, et al. A new definition of genetic counseling: National Society of Genetic Counselors' Task Force report. *J Genet Couns*. 2006;15:77–83.
19. Phillips A, Niemiec E, Howard HC, Kagkelari K, Borry P, Vears DF. Communicating genetic information to family members: analysis of consent forms for diagnostic genomic sequencing. *Eur J Hum Genet*. 2020;28:1160–7. <https://doi.org/10.1038/s41431-020-0627-7>.
20. Lewis C, Sanderson SC, Hammond J, Hill M, Searle B, Hunter A, et al. Development and mixed-methods evaluation of an online animation for young people about genome sequencing. *Eur J Hum Genet*. 2020;28:896–906.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Correspondence and requests for materials should be addressed to Ashley Crook.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.