




# Infographic: Collaborative Normal-Tension Glaucoma Study (CNTGS)

R. Mercer<sup>1</sup> · R. G. Mathew<sup>1</sup> · C. Henein<sup>1</sup>

Received: 6 December 2020 / Revised: 16 December 2020 / Accepted: 21 January 2021 / Published online: 5 March 2021  
© The Author(s), under exclusive licence to The Royal College of Ophthalmologists 2021



✉ C. Henein  
c.henein@ucl.ac.uk

<sup>1</sup> UCL Institute of Ophthalmology, London, UK

◀ **Fig. 1** The Collaborative Normal-Tension Glaucoma Study (CNTGS) showed that a 30% reduction of intraocular pressure in patients with normal tension glaucoma slowed the rate of visual field progression compared to eyes in which no effort was made to lower intraocular pressure. IOP intraocular pressure, NTG normal tension glaucoma, RCT randomised controlled trial, Sx Surgery, VF visual field.

**Reference to original study:** Collaborative Normal-Tension Glaucoma Study Group. The effectiveness of

intraocular pressure reduction in the treatment of normal-tension glaucoma. *Am J Ophthalmol.* 1998;126:498–505.

### **Compliance with ethical standards**

**Conflict of interest** The authors declare no competing interests.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.