




Infographic: Collaborative Initial Glaucoma Treatment Study (CIGTS)

A. Mohamed¹ · J. E. Han¹ · Y. Galiyev¹ · C. Henein¹ · R. G. Mathew¹

Received: 15 December 2020 / Revised: 18 December 2020 / Accepted: 21 January 2021 / Published online: 5 March 2021
© The Author(s), under exclusive licence to The Royal College of Ophthalmologists 2021



Icons made by Freepik from www.flaticon.com

Fig. 1 The Collaborative Initial Glaucoma Treatment Study (CIGTS) showed early surgical intervention in patients with POAG reduced visual field progression at 6 years. IOP intraocular pressure, POAG Primary open angle glaucoma, RR relative risk, NNT number needed to treat, MD mean deviation.

✉ R. G. Mathew
r.mathew@ucl.ac.uk

¹ UCL Institute of Ophthalmology, London, UK

Reference to original study: Musch DC, Gillespie BW, Niziol LM, Lichter PR, Varma R, Group CS. Intraocular pressure control and long-term visual field loss in the Collaborative Initial Glaucoma Treatment Study. *Ophthalmology* 2011;118(9):1766–1773.

Compliance with ethical standards

Conflict of interest The authors declare no competing interests.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.