



## CORRECTION



# Correction: Expanded spectrum of focal scleral nodule: focal scleral nodules can be bifocal

Hyo Song Park, Yong Joon Kim, Eun Young Choi, Sung Chul Lee, Suk Ho Byeon , Sung Soo Kim and Christopher Seungkyu Lee 

© The Author(s), under exclusive licence to The Royal College of Ophthalmologists 2022

*Eye* (2023) 37:792–793; <https://doi.org/10.1038/s41433-022-02203-2>

Correction to: *Eye* <https://doi.org/10.1038/s41433-022-02029-y>, published online 12 April 2022

The authors have found out that the “Neurosensory layer status, no. (%)” section of Table 2 had unnecessary typos which should have been removed (Normal = 1 Absent = 0 Thinning = 2 Thickening = 3 of the previous table), and that the some of the numbers of this section did not add up to 13. The table was revised, and the result section of the manuscript describing numbers of Table 2 was changed accordingly. Two additional sentences that were found to be including faulty information of the Table 2 were revised based on the information of Table 2.

Furthermore, the authors would like the figures of the paper to be printed in color rather than black-and-white as it is now since the figure depicts lesions of which color, shape, and size can be important factors.

The authors are truly sorry for the fact that we were unable to detect the errors during the publication process and the inconvenience this might cause. We thank the editors for the time to be spent on this correction process.

The original article has been corrected.

**Table 2.** Multimodal imaging findings in patients with focal scleral nodule.

Imaging method features	Results			
OCT ( <i>n</i> = 13)				
Mean horizontal length (mm)	2.6 ± 1.2			
Mean thickness (μm)	697 ± 174			
Configuration, no. (%)				
Dome	11 (85)			
Nodule	2 (15)			
Volcanic	0 (0)			
Neurosensory layers status, no. (%)				
	Normal	Absent	Thinning	Thickening
Internal limiting membrane	13 (100)	0 (0)	0 (0)	0 (0)
Nerve fiber layer	12 (92.3)	0 (0)	0 (0)	1 (7.7)
Ganglion cell layer	12 (92.3)	0 (0)	1 (7.7)	0 (0)
Inner plexiform layer	11 (84.6)	0 (0)	2 (15.4)	0 (0)
Inner nuclear layer	9 (69.2)	0 (0)	4 (30.8)	0 (0)
Outer plexiform layer	8 (61.5)	0 (0)	5 (38.5)	0 (0)
Outer nuclear layer	7 (53.8)	1 (7.7)	5 (38.5)	0 (0)
External limiting membrane	8 (61.5)	4 (30.8)	1 (7.7)	0 (0)
Ellipsoid layer	0 (0)	11 (84.6)	0 (0)	2 (15.4)
Retinal pigment epithelium, no. (%)				
	Regular	Irregular		
Regularity	10 (76.9)	3 (23.1)		
	Normal	Absent	Thinning	Thickening
Thickness	5 (38.5)	0 (0)	1 (7.7)	7 (53.8)
Choroid above the lesion				
	Normal	Absent	Thinning	Thickening
Qualitative thickness, no. (%)	1 (7.7)	9 (69.2)	3 (23.1)	0 (0)
Thickness (mm)	Mean, 36 ± 75; range, 0–265; median, 0			
Presence of subretinal fluid				
	No	Yes		
	9 (69.2)	4 (30.8)		
OCT angiography ( <i>n</i> = 5)				
	Normal flow	Decreased flow	Increased flow	
Retinal circulation	5 (100.0)	0 (0)	0 (0)	
Choroidal circulation	0 (0)	5 (100.0)	0 (0)	
Fundus autofluorescence no. (%) ( <i>n</i> = 12)				
	Normal	Hyperautofluorescent	Hypoautofluorescent	Stippled
Mass	2 (16.7)	4 (33.3)	4 (33.3)	2 (16.7)
Halo	6 (50.0)	3 (25.0)	3 (25.0)	0 (0)
Near infrared reflectance, no. (%)				
	Isoreflective	Bright	Dark	
Mass	6 (40.0)	6 (40.0)	3 (20.0)	
Halo	2 (13.3)	8 (53.3)	5 (33.3)	
B-scan ultrasonography ( <i>n</i> = 10)				
	Solid, 3 (30.0)	Hollow, 7 (70.0)		
Thickness (mm), mean ± SD	1.8 ± 0.7			
Fluorescein angiography, no. (%) ( <i>n</i> = 10)				
	Hypofluorescence	Hyperfluorescence	Normal fluorescence	
Early	7 (70.0)	2 (20.0)	1 (10.0)	
Late	6 (60.0)	4 (40.0)	0 (0)	
Indocyanine green angiography, no. (%) ( <i>n</i> = 10)				
	Hypofluorescence	Hyperfluorescence	Normal fluorescence	
Early	10 (100.0)	0 (0)	0 (0)	
Late	10 (100.0)	0 (0)	0 (0)	

OCT optical coherence tomography, SD standard deviation.