NEWS

Mapping the clinical phenome

Genome-wide association studies (GWAS) endeavor to identify genetic variants associated with selected diseases or traits, usually by evaluating one disease or trait at a time. In contrast, phenome-wide association studies (PheWAS) investigate cross-phenotype associations by seeking connections between genetic variants and multiple phenotypes

simultaneously. Among the possible outcomes of PheWAS are identification of pleiotropic effects of individual genetic variants and illumination of the genetic bases of comorbidities. PheWAS also have the potential to uncover previously unrecognized genetic links between seemingly unrelated diseases or traits. In a recent article in the *American Journal of Human Genetics* (<https://doi.org/10.1016/j.ajhg.2018.11.006>), Verma et al. describe utilization of the PheWAS approach to develop a disease–disease network (DDN) in which the existence of a shared genetic variant served as the basis for the formulation of disease–disease pairs. To develop their DDN, the research team extracted data on genetic variants and diagnostic codes from the Geisinger electronic health record (EHR) and analyzed associations between 632,574 common genetic variants and 541 International Classification of Diseases, ninth revision (ICD-9) diagnostic codes from 38,682 unrelated individuals. By using a single-source EHR for their PheWAS, the authors were able to reduce inconsistencies in phenotype descriptions across the study population and control for analytical variability by employing a single genotyping platform for variant identification.

Associations with P values $< 1 \times 10^{-4}$ were used to construct the DDN. ICD-9 codes were grouped into 15 disease classes to permit assessment of disease–disease connections within and across disease classes. The DDN developed by the research team included 385 disease diagnoses (nodes) that were connected by 1398 edges. Interconnections between diseases, which suggest shared genetic foundations and common pathophysiological mechanisms, were observed across multiple disease classes, with the strongest connections found among autoimmune disorders. The average number of direct connections for each disease was 7 (range 1–32). Larger, hub nodes—nodes with significantly more edges than other nodes—were identified across different disease classes, with the highest number of hub nodes found among endocrine disorders. The team also applied community detection methods to characterize subnetworks within the DDN and aid investigation of biological explanations for disease connections; this analysis identified nine communities composed of 2–102 diseases. The authors note the discovery of novel and unexpected disease–disease connections, including, for example, a link between hyperlipidemia and immune disorders. They conclude that PheWAS-based community detection can identify cross-phenotype associations and reveal previously unknown links between diseases. The authors also assert that this approach can be adapted to publicly available databases and suggest that perspectives gained from PheWAS could impact research and clinical practice by advancing knowledge and understanding of disease mechanisms. —Raye Alford, News Editor

Mining patterns of genetic variation to identify coding regions subject to purifying selection

Constrained coding regions (CCRs) are regions of the genome that exhibit notably less DNA sequence variation than would normally be expected. The assumption is that these regions encode essential functions that are intolerant to changes in protein sequence. To date, the analytical methods used to identify CCRs have examined patterns of genetic variation at the resolution of whole genes. As a result, these methods fail to detect constrained variation within genes. In a recent article in *Nature Genetics* (<https://doi.org/10.1038/s41588-018-0294-6>), Havrilla et al. sought to measure this unexplored genetic variability, or lack thereof, and to develop a detailed CCR map of the human exome. The research team used the Genome Aggregation Database (gnomAD) to search for variation in protein-coding sequences. The gnomAD v2.0.1 resource is based on 123,136 exomes and includes nearly 4.8 million missense or loss-of-function variants, which occur at an average density of 1 variant every 7 coding base pairs. Coding regions considered to be constrained were those that showed a greater than average number of base pairs between protein-altering variants. Controlling for confounding factors such as sequencing coverage, segmental duplications, and high-identity repeats, the team quantified constraint across 88% and 82% of the autosomal exome and X chromosome, respectively. Adjusting for CpG dinucleotide density, the team established percentile ranks for constrained regions and computed median CCR lengths of 52 base pairs for the 95th percentile and 94 base pairs for the 99th percentile. The most highly constrained CCRs—those at or above the 95th percentile—were enriched for ClinVar-listed pathogenic variants, while the least constrained CCRs—those below the 20th percentile—were depleted of pathogenic variants. This trend was observed for all autosomal genes taken as a group as well as for autosomal dominant disease genes separately. For de novo missense variants identified in probands with neurodevelopmental disorders, CCRs at or above the 95th percentile demonstrated a 7.1-fold enrichment for pathogenic variants compared with the unaffected siblings of probands with autism spectrum disorder, while CCRs below the 20th percentile demonstrated a 4.0-fold depletion of pathogenic variants. More than 22,000 CCRs in the 95th percentile or higher lacked a pathogenic variant in ClinVar, suggesting that these CCRs were subject to extreme selective pressure. The authors conclude that CCRs define structurally or functionally important genetic domains that are subject to strong purifying selection. They assert that the most highly constrained CCRs represent regions in which variation results in severe developmental disorders or embryonic lethality. They also suggest that CCRs can aid interpretation of de novo and rare variants in cases affected by dominant phenotypes. —Raye Alford, News Editor